



Time Series Analysis and Forecasting

Chapter 2: ARMA Models



Daniel Traian PELE

Bucharest University of Economic Studies

IDA Institute Digital Assets

Blockchain Research Center

AI4EFin Artificial Intelligence for Energy Finance

Romanian Academy, Institute for Economic Forecasting

MSCA Digital Finance

Learning Objectives

By the end of this chapter, you will be able to:


1. **Define** and simulate $AR(p)$, $MA(q)$, and $ARMA(p, q)$ processes
2. **Verify** stationarity and invertibility conditions
3. **Identify** orders p and q through ACF/PACF analysis
4. **Estimate** parameters via Yule-Walker, MLE, and information criteria (AIC, BIC)
5. **Diagnose** the model through residual analysis and the Ljung-Box test
6. **Forecast** using ARMA models with confidence intervals
7. **Apply** the Box-Jenkins methodology to real data (sunspots)

Outline

- Motivation
- Introduction and the Lag Operator
- Autoregressive (AR) Models
- Moving Average (MA) Models
- ARMA Models
- Model Identification
- Parameter Estimation
- Model Diagnostics
- Forecasting with ARMA
- Practical Implementation
- Case Study: Real Data
- Summary
- Quiz




Why ARMA Models?



- **AR processes:** Current value depends on past values \Rightarrow mean-reverting behavior
- **MA processes:** Current value depends on past shocks \Rightarrow short memory
- **ARMA:** Combines both mechanisms for flexible modeling

Model Identification Through ACF Patterns

Distinct ACF patterns for different models



ACF Reflects Model Structure

- ▣ **Distinct patterns:** AR: exponential decay; MA: sharp cutoff; ARMA: mixed decay
- ▣ **Identification:** Visual analysis of ACF/PACF guides the selection of orders p and q

Recap: Stationarity

From Chapter 1

- A process $\{X_t\}$ is **weakly stationary** if:
 1. $\mathbb{E}[X_t] = \mu$ (constant mean)
 2. $\text{Var}(X_t) = \sigma^2 < \infty$ (constant, finite variance)
 3. $\text{Cov}(X_t, X_{t+h}) = \gamma(h)$ (covariance depends only on lag h)


Why Stationarity Matters for ARMA

- ARMA models assume stationarity
 - ▶ Parameters remain stable over time
 - ▶ Autocorrelation structure is maintained
- Non-stationary data \Rightarrow difference first (ARIMA, Ch. 3)

Chapter Objective

- Parametric models for stationary series \Rightarrow combining dependence on past observations (AR) with the influence of random shocks (MA)

The Lag Operator: Visual Illustration



Role of the Lag Operator

- ▣ **Notation foundation:** Enables compact writing of difference equations
- ▣ **Utility:** Facilitates algebraic manipulation of ARMA models

The Lag Operator (Backshift Operator)

Definition 1 (Lag Operator)

- The **lag operator** L (or backshift operator B) shifts a time series back by one period: $LX_t = X_{t-1}$


Properties

- $L^k X_t = X_{t-k}$ (shift back by k periods)
- $L^0 X_t = X_t$ (identity)
- $(1 - L)X_t = X_t - X_{t-1} = \Delta X_t$ (first difference)
- $(1 - L)^d X_t = \Delta^d X_t$ (difference of order d)

Lag Polynomials

- **AR polynomial:** $\phi(L) = 1 - \phi_1 L - \phi_2 L^2 - \dots - \phi_p L^p$
- **MA polynomial:** $\theta(L) = 1 + \theta_1 L + \theta_2 L^2 + \dots + \theta_q L^q$

White Noise: Visual Illustration



Key Characteristics

- **Left:** Random fluctuations, no patterns, constant variance
- **Right:** ACF only a spike at lag 0; others within significance bounds \Rightarrow no linear dependence

The White Noise Process

Definition 2 (White Noise)


- A process $\{\varepsilon_t\}$ is **white noise**, denoted $\varepsilon_t \sim WN(0, \sigma^2)$, if:
 1. $\mathbb{E}[\varepsilon_t] = 0$ for all t
 2. $\text{Var}(\varepsilon_t) = \sigma^2$ for all t
 3. $\text{Cov}(\varepsilon_t, \varepsilon_s) = 0$ for all $t \neq s$

Properties

- **Building block:** White noise underlies all ARMA models
- **ACF:** $\rho(0) = 1$, $\rho(h) = 0$ for $h \neq 0$; PACF: same pattern
- **Gaussian white noise:** $\varepsilon_t \sim N(0, \sigma^2)$ i.i.d.
- **Unpredictable:** White noise is *not* predictable \Rightarrow it is purely random

AR(1): Visual Illustration

AR(1): different behavior for positive vs negative ϕ



Visual Interpretation

- ▣ **Positive ϕ :** Persistent fluctuations, gradual mean reversion
- ▣ **Negative ϕ :** Oscillating behavior, alternating around the mean
- ▣ **Larger $|\phi|$ \Rightarrow greater persistence, slower reversion**

The AR(1) Model: Definition

Definition 3 (AR(1) Process)

- An **autoregressive process of order 1** is: $X_t = c + \phi X_{t-1} + \varepsilon_t$
- $\varepsilon_t \sim WN(0, \sigma^2)$ and $|\phi| < 1$ for stationarity

Interpretation

- c : constant (intercept)
- ϕ : autoregressive coefficient
 - ▶ Measures the persistence of the series
- ε_t : innovation (shock)

Lag Operator Notation

- $(1 - \phi L)X_t = c + \varepsilon_t$
- $\phi(L)X_t = c + \varepsilon_t$
- $\phi(L) = 1 - \phi L$

AR(1) Stationarity Condition

Necessary and Sufficient Condition: $|\phi| < 1$

- The root of the characteristic equation must lie outside the unit circle

- Shocks diminish over time
 - ▶ Process reverts to the mean
 - ▶ Finite, stable variance

Non-stationary ($|\phi| \geq 1$)

- $|\phi| = 1$: random walk
 - ▶ Unit root, variance $\rightarrow \infty$
- $|\phi| > 1$: explosive process

Characteristic Equation

- $\phi(z) = 1 - \phi z = 0 \implies z = 1/\phi$
- Stationarity \Leftrightarrow root outside the unit circle ($|z| > 1$)

AR(1) Properties

Stationary AR(1) with $|\phi| < 1$

□ Moment properties:

Mean: $\mu = \mathbb{E}[X_t] = \frac{c}{1-\phi}$

Variance: $\gamma(0) = \text{Var}(X_t) = \frac{\sigma^2}{1-\phi^2}$


Autocovariance: $\gamma(h) = \frac{\phi^h \sigma^2}{1-\phi^2}$

Autocorrelation (ACF): $\rho(h) = \phi^h$

Key Observation

□ **AR(1) signature:** ACF decays exponentially with factor ϕ


- ▶ $\phi > 0$: monotone decay towards zero
- ▶ $\phi < 0$: damped oscillations (alternating signs)

AR(1) Simulations: Effect of ϕ 

Interpretation

- Different values of ϕ produce distinct behaviors: larger $|\phi| \Rightarrow$ more persistence
- Positive ϕ creates smooth trajectories; negative ϕ creates oscillations
- As $|\phi| \rightarrow 1$, the process approaches non-stationarity

Theoretical AR(1) ACF




ACF Pattern

- Formula: $\rho(h) = \phi^h \Rightarrow$ exponential decay
- $\phi > 0$: monotone decay; $\phi < 0$: alternating signs

AR(1) ACF and PACF: Theory vs Sample

ACF and PACF for AR(1): theory vs sample



Interpretation

- ACF: Exponential decay with factor ϕ ; formula: $\rho(h) = \phi^h$
- PACF: A single spike at lag 1, then cuts off \Rightarrow identifies AR(1)
- Sample estimates fluctuate around theoretical values

Proof: AR(1) Mean

Claim

- For AR(1): $X_t = c + \phi X_{t-1} + \varepsilon_t$, the mean is $\mu = \frac{c}{1-\phi}$

Proof

- Take expectations of both sides: $\mathbb{E}[X_t] = c + \phi\mathbb{E}[X_{t-1}] + \mathbb{E}[\varepsilon_t]$
- By stationarity, $\mathbb{E}[X_t] = \mathbb{E}[X_{t-1}] = \mu$, and $\mathbb{E}[\varepsilon_t] = 0$: $\mu = c + \phi\mu$
- Solving: $\mu - \phi\mu = c \implies \mu(1 - \phi) = c \implies \boxed{\mu = \frac{c}{1 - \phi}}$

Requirement

- **Necessary condition:** $\phi \neq 1$ for the mean to be defined
 - ▶ If $\phi = 1$ (unit root), the mean is undefined
 - ▶ The process becomes a random walk (non-stationarity)

Proof: AR(1) Variance

Claim

$$\square \text{Var}(X_t) = \frac{\sigma^2}{1-\phi^2}$$

Proof

\square Assume $c = 0$. Take the variance of $X_t = \phi X_{t-1} + \varepsilon_t$:


$$\square \text{Var}(X_t) = \phi^2 \text{Var}(X_{t-1}) + \text{Var}(\varepsilon_t) + 2\phi \underbrace{\text{Cov}(X_{t-1}, \varepsilon_t)}_{=0}$$

\square By stationarity, $\text{Var}(X_t) = \text{Var}(X_{t-1}) = \gamma(0)$:

$$\square \gamma(0) = \phi^2 \gamma(0) + \sigma^2 \implies \gamma(0)(1 - \phi^2) = \sigma^2 \implies \boxed{\gamma(0) = \frac{\sigma^2}{1 - \phi^2}}$$

Note

\square Requires $|\phi| < 1$ for positive variance. When $|\phi| \rightarrow 1$, variance $\rightarrow \infty$

AR(1) Variance as a Function of ϕ 

Observations

- As $|\phi| \rightarrow 1$, the variance explodes \Rightarrow non-stationarity
- For $\phi = 0$: $\gamma(0) = \sigma^2$ (white noise); variance increases monotonically with $|\phi|$

Proof: AR(1) Autocorrelation Function

Claim: $\rho(h) = \phi^h$ for $h \geq 0$

- Find the autocovariance $\gamma(h) = \text{Cov}(X_t, X_{t-h})$

Proof

- Multiply $X_t = \phi X_{t-1} + \varepsilon_t$ by X_{t-h} and take expectations:
- $\mathbb{E}[X_t X_{t-h}] = \phi \mathbb{E}[X_{t-1} X_{t-h}] + \mathbb{E}[\varepsilon_t X_{t-h}]$
- For $h \geq 1$: $\mathbb{E}[\varepsilon_t X_{t-h}] = 0 \Rightarrow \gamma(h) = \phi \gamma(h-1)$
- Recursive relation from $\gamma(0)$: $\gamma(1) = \phi \gamma(0)$, $\gamma(2) = \phi^2 \gamma(0)$, ... $\gamma(h) = \phi^h \gamma(0)$
- ACF: $\rho(h) = \frac{\gamma(h)}{\gamma(0)} = \frac{\phi^h \gamma(0)}{\gamma(0)} = \boxed{\phi^h}$

Proof: AR(1) Stationarity Condition

Claim

- AR(1) is stationary if and only if $|\phi| < 1$

Proof

- Recursive substitution: $X_t = \phi X_{t-1} + \varepsilon_t = \phi(\phi X_{t-2} + \varepsilon_{t-1}) + \varepsilon_t = \dots$
- After n steps: $X_t = \phi^n X_{t-n} + \sum_{j=0}^{n-1} \phi^j \varepsilon_{t-j}$
- If $|\phi| < 1$: $\phi^n \rightarrow 0$ as $n \rightarrow \infty$, so $X_t = \sum_{j=0}^{\infty} \phi^j \varepsilon_{t-j}$
- Finite variance: $\text{Var}(X_t) = \sigma^2 \sum_{j=0}^{\infty} \phi^{2j} = \frac{\sigma^2}{1-\phi^2} < \infty$ (geometric series)

Conclusion

- Converges $\iff |\phi| < 1$. For $|\phi| \geq 1$, the term $\phi^n X_{t-n}$ does not vanish \Rightarrow infinite variance

The Partial Autocorrelation Function (PACF)

Definition 4 (PACF)

- The **partial autocorrelation** of order k , denoted π_k , measures the correlation between X_t and X_{t-k} **after removing** the linear effects of the intermediate variables $X_{t-1}, \dots, X_{t-k+1}$

Formal Definition

- $\pi_1 = \rho(1)$
- For $k \geq 2$: π_k is the last coefficient in:
$$X_t = \alpha_1 X_{t-1} + \dots + \alpha_k X_{t-k} + e_t$$
- $\pi_k = \alpha_k$ (coefficient of X_{t-k})

Computation via Yule-Walker

- Solve the Yule-Walker equations of order k
- π_k = last element of the solution vector

Utility

- **Identification:** PACF determines the order p of an AR model
 - ▶ PACF cuts off after lag p

Durbin-Levinson Algorithm for PACF

Durbin-Levinson Recursion

▣ Computes PACF (π_k) recursively, without inverting the Toeplitz matrix:

1. **Initialize:** $\pi_1 = \hat{\rho}(1)$, $v_1 = \hat{\gamma}(0)(1 - \pi_1^2)$

2. **Recursion** ($k = 2, 3, \dots$):

$$\pi_k = \frac{\hat{\rho}(k) - \sum_{j=1}^{k-1} \phi_{k-1,j} \hat{\rho}(k-j)}{1 - \sum_{j=1}^{k-1} \phi_{k-1,j} \hat{\rho}(j)}$$

3. **Update:** $\phi_{k,j} = \phi_{k-1,j} - \pi_k \phi_{k-1,k-j}$ for $j = 1, \dots, k-1$; $\phi_{k,k} = \pi_k$

4. **Prediction variance:** $v_k = v_{k-1}(1 - \pi_k^2)$

Complexity


- ▣ $O(k^2)$ vs $O(k^3)$ (direct inversion)
- ▣ Exploits Toeplitz structure of Γ_k
- ▣ Guarantees $v_k > 0$ if stationary

AR(p) Identification

- ▣ $\pi_k = 0$ for $k > p \Rightarrow$ order is p
- ▣ CI: $|\pi_k| > 1.96/\sqrt{T} \Rightarrow$ significant
- ▣ Equivalent to t -test on last OLS coeff.

AR(p): Visual Illustration

AR(2) Process: pseudo-cyclic behavior



Observations

- AR(2) can exhibit pseudo-cyclic behavior (complex roots); damped sinusoidal ACF
- PACF cuts off after lag 2 \Rightarrow key identification pattern

The AR(p) Model: General Form

Definition 5 (AR(p) Process)

- ▣ An **autoregressive process of order p** is: $X_t = c + \phi_1 X_{t-1} + \phi_2 X_{t-2} + \cdots + \phi_p X_{t-p} + \varepsilon_t$
- ▣ **Lag operator**: $\phi(L)X_t = c + \varepsilon_t$, where $\phi(L) = 1 - \phi_1 L - \phi_2 L^2 - \cdots - \phi_p L^p$



Stationarity Condition

- ▣ All roots of $\phi(z) = 0$ must lie **outside** the unit circle
- ▣ Equivalently: all roots have modulus > 1

PACF Pattern

- ▣ PACF cuts off after lag p
- ▣ ACF decays (exponentially or with damped oscillations)


AR(2) Stationarity: Unit Circle Visualization



Characteristic Polynomial and Unit Circle Condition

- **Characteristic polynomial** of $AR(p)$: $\phi(z) = 1 - \phi_1 z - \phi_2 z^2 - \dots - \phi_p z^p$
- All roots of $\phi(z) = 0$ must lie **outside** the unit circle ($|z| > 1$)
- Roots on the circle: non-stationary; roots inside: explosive process


The AR(2) Stationarity Triangle



Stationarity Region

- The triangular region defines the stationary AR(2) parameter combinations
- **Boundaries:** $\phi_1 + \phi_2 < 1$, $\phi_2 - \phi_1 < 1$ and $|\phi_2| < 1$
- Points outside the region \Rightarrow non-stationary or explosive processes

Characteristic Polynomial Roots



Types of Roots

- Real roots: exponential decay in ACF
- Complex roots: damped oscillations (pseudo-cycles)
- All roots must lie outside the unit circle

The AR(2) Model

Definition 6 (AR(2) Process)

$$\square X_t = c + \phi_1 X_{t-1} + \phi_2 X_{t-2} + \varepsilon_t$$

Stationarity Conditions


$$\square \phi_1 + \phi_2 < 1; \quad \phi_2 - \phi_1 < 1; \quad |\phi_2| < 1$$

ACF Behavior

- \square **Real roots:** mixture of two exponential decays
- \square **Complex roots:** damped sinusoidal pattern (pseudo-cycles)
- \square **PACF:** Cuts off after lag 2 ($\pi_k = 0$ for $k > 2$)

MA(1): Visual Illustration

MA(1): short memory series with ACF cutoff



Visual Interpretation

- Left panel: MA(1) series \Rightarrow rapid mean reversion
- Right panel: ACF with **cutoff** after lag 1; PACF exponential decay

The MA(1) Model: Definition

Definition 7 (MA(1) Process)

- ▣ A **moving average process of order 1** is: $X_t = \mu + \varepsilon_t + \theta\varepsilon_{t-1}$
- ▣ $\varepsilon_t \sim WN(0, \sigma^2)$

Interpretation

- ▣ μ : process mean
- ▣ θ : MA coefficient
 - Measures the impact of the past shock
- ▣ Depends on ε_t and ε_{t-1}

Lag Operator Notation

- ▣ $X_t = \mu + \theta(L)\varepsilon_t$
- ▣ $\theta(L) = 1 + \theta L$

Key Property

- ▣ **Guaranteed stationarity:** MA processes are always stationary
 - Does not depend on the value of θ

MA(1) Properties

$$\text{MA}(1): X_t = \mu + \varepsilon_t + \theta\varepsilon_{t-1}$$

- **Mean:** $\mathbb{E}[X_t] = \mu$; **Variance:** $\gamma(0) = \sigma^2(1 + \theta^2)$
- **Autocovariance:** $\gamma(1) = \theta\sigma^2$, $\gamma(h) = 0$ ($h > 1$)
- **ACF:** $\rho(1) = \frac{\theta}{1+\theta^2}$, $\rho(h) = 0$ ($h > 1$)

Key Observation

- **MA(1) signature:** ACF cuts off after lag 1
 - ▶ $\rho(1) \neq 0$, but $\rho(h) = 0$ for $h > 1$; opposite pattern to AR(1)

Proof: MA(1) Variance and Autocovariance

Starting point: $X_t = \varepsilon_t + \theta\varepsilon_{t-1}$ (assuming $\mu = 0$)

▣ **Variance:**

$$\gamma(0) = \text{Var}(\varepsilon_t + \theta\varepsilon_{t-1}) = \sigma^2 + \theta^2\sigma^2 + 0 = \boxed{\sigma^2(1 + \theta^2)}$$

Autocovariance at lag 1

$$\begin{aligned}\square \gamma(1) &= \text{Cov}(\varepsilon_t + \theta\varepsilon_{t-1}, \varepsilon_{t-1} + \theta\varepsilon_{t-2}) \\ \square &= \text{Cov}(\varepsilon_t, \varepsilon_{t-1}) + \theta\text{Cov}(\varepsilon_t, \varepsilon_{t-2}) + \theta\text{Cov}(\varepsilon_{t-1}, \varepsilon_{t-1}) + \theta^2\text{Cov}(\varepsilon_{t-1}, \varepsilon_{t-2}) \\ \square &= 0 + 0 + \theta\sigma^2 + 0 = \boxed{\theta\sigma^2}\end{aligned}$$

Autocovariance at lag $h \geq 2$

▣ No common ε terms $\Rightarrow \gamma(h) = 0$

Proof: Maximum ACF for MA(1)


Claim: $|\rho(1)| \leq 0.5$ for any value of θ

- ACF at lag 1: $\rho(1) = \frac{\theta}{1+\theta^2}$
- Differentiate: $\frac{d\rho(1)}{d\theta} = \frac{1-\theta^2}{(1+\theta^2)^2} = 0 \Rightarrow \theta = \pm 1$
- At these values: $\rho(1)|_{\theta=1} = \frac{1}{2}$, $\rho(1)|_{\theta=-1} = -\frac{1}{2}$

Implication

- **Practical test:** If $|\hat{\rho}(1)| > 0.5$ from data, the process is **not** MA(1)
 - ▶ The maximum $|\rho(1)| = 0.5$ is reached at $\theta = \pm 1$
 - ▶ Consider AR or ARMA models as alternatives

MA(1) Simulations: Effect of θ



Interpretation

- MA(1) is always stationary regardless of $\theta \Rightarrow$ finite memory of only one lag
- Positive θ smooths the series; negative θ creates faster fluctuations
- Unlike AR(1), MA(1) shocks affect the process for only one period

Proof: ACF for MA(1)

Claim: $\rho(1) = \frac{\theta}{1+\theta^2}$ and $\rho(h) = 0$ for $h > 1$

- MA(1) has non-zero autocorrelation **only** at lag 1

Proof


- Let $X_t = \varepsilon_t + \theta\varepsilon_{t-1}$. Autocorrelation at lag 1:
- $\rho(1) = \frac{\gamma(1)}{\gamma(0)} = \frac{\theta\sigma^2}{\sigma^2(1+\theta^2)} = \frac{\theta}{1+\theta^2}$
- For $h > 1$: $\gamma(h) = \text{Cov}(\varepsilon_t + \theta\varepsilon_{t-1}, \varepsilon_{t-h} + \theta\varepsilon_{t-h-1})$
- The terms $\varepsilon_t, \varepsilon_{t-1}$ do not overlap with $\varepsilon_{t-h}, \varepsilon_{t-h-1}$ when $h > 1$, so $\gamma(h) = 0$

Practical Consequence

- ACF cuts off sharply after lag 1 \Rightarrow distinctive signature of MA(1) processes

MA(1) ACF and PACF: Visual Comparison

ACF and PACF for MA(1): opposite pattern to AR(1)





Interpretation

- ACF: A single spike at lag 1, then cuts off \Rightarrow key MA(1) signature
- PACF: Exponential decay \Rightarrow opposite pattern to AR(1)
- This reversal differentiates MA processes from AR processes

Invertibility: Visual Illustration

Invertibility of MA models



Interpretation

- Left: invertibility requires roots outside the unit circle
- Right: $AR(\infty)$ weights decay only when $|\theta| < 1$

Invertibility of MA Models

Definition 8 (Invertibility)

- An MA process is **invertible** if it can be written as an infinite AR process:
- $X_t = \mu + \sum_{j=1}^{\infty} \pi_j (X_{t-j} - \mu) + \varepsilon_t$

Invertibility Conditions

- **MA(1)**: Invertible if $|\theta| < 1$
- **MA(q)**: Roots of $\theta(z) = 0$ outside the unit circle

Why Invertibility Matters

- Ensures unique representation (without invertibility, multiple MA models describe the same data)
- Necessary for forecasting and estimation
- **Stationarity** \Rightarrow AR; **Invertibility** \Rightarrow MA

Proof: MA(1) Invertibility

Claim

- MA(1) is invertible if and only if $|\theta| < 1$

Proof

- From $X_t = \varepsilon_t + \theta\varepsilon_{t-1}$, isolate: $\varepsilon_t = X_t - \theta\varepsilon_{t-1}$
- Recursive back-substitution: $\varepsilon_t = X_t - \theta(X_{t-1} - \theta\varepsilon_{t-2}) = X_t - \theta X_{t-1} + \theta^2\varepsilon_{t-2}$
- Continuing: $\varepsilon_t = \sum_{j=0}^n (-\theta)^j X_{t-j} + (-\theta)^{n+1} \varepsilon_{t-n-1}$


- If $|\theta| < 1$: $(-\theta)^{n+1} \rightarrow 0$, so
$$\varepsilon_t = \sum_{j=0}^{\infty} (-\theta)^j X_{t-j}$$

Conclusion

- Geometric series converges $\iff |\theta| < 1 \Rightarrow$ MA(1) can be written as AR(∞)

MA(q): Visual Illustration

MA(q) Process: ACF signature cuts off after lag q



Observation

- MA(3) process: key signature \Rightarrow ACF cuts off after lag q ($\rho(h) = 0$ for $h > 3$)

The MA(q) Model: General Form

Definition 9 (MA(q) Process)


- ▣ A **moving average process of order q**: $X_t = \mu + \varepsilon_t + \theta_1\varepsilon_{t-1} + \cdots + \theta_q\varepsilon_{t-q}$
- ▣ **Lag operator**: $X_t = \mu + \theta(L)\varepsilon_t$, where $\theta(L) = 1 + \theta_1L + \cdots + \theta_qL^q$

Properties

- ▣ Always stationary (finite variance)
- ▣ ACF cuts off after lag q : $\rho(h) = 0$ for $h > q$; PACF decays gradually
- ▣ Invertible if all roots of $\theta(z) = 0$ lie outside the unit circle

ARMA: Visual Illustration

ARMA(1,1): neither ACF nor PACF cut off



ARMA(1,1) Interpretation

- Combines AR persistence with MA shock response
- ACF pattern:** Decay after the first lag (lags decay geometrically)
- PACF pattern:** Also decays (no sharp cutoff as in pure AR)
- Neither ACF nor PACF cuts off \Rightarrow key identifier for mixed models

The ARMA(p,q) Model: Definition

Definition 10 (ARMA(p,q) Process)

- $X_t = c + \phi_1 X_{t-1} + \cdots + \phi_p X_{t-p} + \varepsilon_t + \theta_1 \varepsilon_{t-1} + \cdots + \theta_q \varepsilon_{t-q}$
- **Compact form:** $\phi(L)X_t = c + \theta(L)\varepsilon_t$, where $\mu = \frac{c}{1 - \phi_1 - \cdots - \phi_p}$


Key Idea

- **Flexibility:** Combines AR and MA components
 - ▶ AR captures persistence; MA captures shock response
- **Parsimony:** ARMA(1,1) can be better than AR(5) or MA(5)
 - ▶ Fewer parameters, less risk of overfitting

ARMA Model Structure

ARMA(p,q) Model Structure


$$X_t = \phi_1 X_{t-1} + \dots + \phi_p X_{t-p} + \varepsilon_t + \theta_1 \varepsilon_{t-1} + \dots + \theta_q \varepsilon_{t-q}$$



Components

- ▣ **AR component:** influence of past values of the series
- ▣ **MA component:** impact of past random shocks



How ARMA Simulation Works



Steps

- Generate white noise, apply the ARMA equation recursively, obtain simulated series

ARMA Examples



Observation

- Different combinations of orders (p, q) produce distinct behaviors

The ARMA(1,1) Model

Definition 11 (ARMA(1,1) Process)

$$\boxed{\cdot} \quad X_t = c + \phi X_{t-1} + \varepsilon_t + \theta \varepsilon_{t-1}$$

Properties (stationarity and invertibility)

$$\boxed{\cdot} \quad \text{Mean: } \mu = \frac{c}{1-\phi}; \quad \text{Variance: } \gamma(0) = \frac{(1+2\phi\theta+\theta^2)\sigma^2}{1-\phi^2}$$

ACF

$$\boxed{\cdot} \quad \rho(1) = \frac{(1+\phi\theta)(\phi+\theta)}{1+2\phi\theta+\theta^2}; \quad \rho(h) = \phi \cdot \rho(h-1) \text{ for } h \geq 2$$

$\boxed{\cdot}$ ACF decays exponentially after lag 1 (starting point depends on ϕ and θ)

Proof: ARMA(1,1) Variance


Claim

$$\square \gamma(0) = \frac{(1+2\phi\theta+\theta^2)\sigma^2}{1-\phi^2}$$

Proof

- Let $Y_t = X_t - \mu$: $Y_t = \phi Y_{t-1} + \varepsilon_t + \theta \varepsilon_{t-1}$
- Square: $Y_t^2 = \phi^2 Y_{t-1}^2 + \varepsilon_t^2 + \theta^2 \varepsilon_{t-1}^2 + 2\phi Y_{t-1} \varepsilon_t + 2\phi\theta Y_{t-1} \varepsilon_{t-1} + 2\theta \varepsilon_t \varepsilon_{t-1}$
- Take expectations; $\mathbb{E}[\varepsilon_t Y_{t-1}] = 0$, $\mathbb{E}[\varepsilon_t \varepsilon_{t-1}] = 0$:
- $\gamma(0) = \phi^2 \gamma(0) + \sigma^2 + \theta^2 \sigma^2 + 2\phi\theta \mathbb{E}[\varepsilon_{t-1} Y_{t-1}]$
- From $Y_{t-1} = \phi Y_{t-2} + \varepsilon_{t-1} + \theta \varepsilon_{t-2}$: only ε_{t-1}^2 contributes $\Rightarrow \mathbb{E}[\varepsilon_{t-1} Y_{t-1}] = \sigma^2$
- $\gamma(0)(1 - \phi^2) = (1 + 2\phi\theta + \theta^2)\sigma^2 \implies \boxed{\gamma(0) = \frac{(1 + 2\phi\theta + \theta^2)\sigma^2}{1 - \phi^2}}$

ACF/PACF Patterns: AR vs MA vs ARMA



Proof: ARMA(1,1) ACF at Lag 1

Claim

$$\square \quad \rho(1) = \frac{(1+\phi\theta)(\phi+\theta)}{1+2\phi\theta+\theta^2}; \quad \rho(h) = \phi \rho(h-1) \text{ for } h \geq 2$$

Proof

□ Multiply Y_t by Y_{t-1} and take expectations:

$$\square \quad \gamma(1) = \phi\gamma(0) + \underbrace{\mathbb{E}[\varepsilon_t Y_{t-1}]}_{=0} + \theta \underbrace{\mathbb{E}[\varepsilon_{t-1} Y_{t-1}]}_{=\sigma^2} = \phi\gamma(0) + \theta\sigma^2$$

□ Divide by $\gamma(0)$: $\rho(1) = \phi + \frac{\theta\sigma^2}{\gamma(0)}$. Substitute $\gamma(0)$:




$$\square \quad \rho(1) = \phi + \frac{\theta(1-\phi^2)}{1+2\phi\theta+\theta^2} = \frac{\phi(1+2\phi\theta+\theta^2)+\theta(1-\phi^2)}{1+2\phi\theta+\theta^2}$$

$$\square \quad \text{Numerator: } \phi + \theta + \phi^2\theta + \phi\theta^2 = (\phi + \theta)(1 + \phi\theta), \text{ so } \boxed{\rho(1) = \frac{(1 + \phi\theta)(\phi + \theta)}{1 + 2\phi\theta + \theta^2}}$$

Recursion

□ For $h \geq 2$: $\gamma(h) = \phi\gamma(h-1)$, so $\rho(h) = \phi \rho(h-1) \Rightarrow$ exponential decay from lag 1

Impulse Response Functions



Shock Propagation

- ▣ Shows how a unit shock propagates through the system over time
- ▣ **AR**: exponential or oscillating decay; **MA**: effect limited to q periods

Stationarity and Invertibility Summary

Conditions for a Valid ARMA(p,q) Model


▣ Requirements summary:

Condition	Requirement
Stationarity	Roots of $\phi(z) = 0$ outside the unit circle
Invertibility	Roots of $\theta(z) = 0$ outside the unit circle

Implications

- ▣ **Stationarity:** Can be written as $MA(\infty)$: $X_t = \mu + \sum_{j=0}^{\infty} \psi_j \varepsilon_{t-j}$
- ▣ **Invertibility:** Can be written as $AR(\infty)$: $X_t = \mu + \sum_{j=1}^{\infty} \pi_j (X_{t-j} - \mu) + \varepsilon_t$
- ▣ **Causal representation:** X_t depends only on *past* shocks \Rightarrow necessary for forecasting


Wold's Decomposition Theorem



Wold's Theorem

- Any purely non-deterministic stationary process can be written as $MA(\infty)$:
- $X_t = \sum_{j=0}^{\infty} \psi_j \varepsilon_{t-j}$ with $\sum \psi_j^2 < \infty$
- Theoretical justification for ARMA modeling

The Box-Jenkins Methodology



Iterative Approach

□ Identification \Rightarrow estimation \Rightarrow validation; repeat until residuals are white noise



ACF/PACF Identification Rules


Theoretical Patterns for Stationary Processes

Model	ACF Pattern	PACF Pattern
AR(1)	Exponential decay	Spike at lag 1, then 0
AR(2)	Exp./damped sinusoid	Spikes at lags 1-2, then 0
AR(p)	Gradual decay	Cuts off after lag p
MA(1)	Spike at lag 1, then 0	Exponential decay
MA(2)	Spikes at lags 1-2, then 0	Exp./damped sinusoid
MA(q)	Cuts off after lag q	Gradual decay
ARMA(p,q)	Decays	Decays

Model Identification: ACF/PACF Patterns

Model	ACF Pattern	PACF Pattern
AR(p)	Exponential decay or damped sinusoid	Cuts off after lag p
MA(q)	Cuts off after lag q	Exponential decay or damped sinusoid
ARMA(p,q)	Exponential decay after lag q	Exponential decay after lag p

AIC vs BIC: Model Selection



Interpretation

- White square marks the best model; lower values (green) are better

Information Criteria

AIC (Akaike)

- $AIC = -2 \ln(\hat{L}) + 2k$
- Moderate penalty
 - ▶ Tends to select larger models
 - ▶ Optimal for forecasting

BIC (Bayesian)


- $BIC = -2 \ln(\hat{L}) + k \ln(n)$
- Stronger penalty
 - ▶ Prefers parsimonious models
 - ▶ Consistent for identification

where: \hat{L} = maximum of the likelihood function, k = number of estimated parameters, n = sample size

Rules

- Lower values = better model. Compare models on the *same data*

Parsimony Principle: Bias-Variance Trade-off



Bias-Variance Trade-off

- Too simple model \Rightarrow high bias (underfitting)
- Too complex model \Rightarrow high variance (overfitting)
- The optimum lies at the intersection of the two curves

Automatic Model Selection

Grid Search Approach

- ▣ Estimate ARMA(p, q) for $p = 0, \dots, p_{max}$ and $q = 0, \dots, q_{max}$
- ▣ Select the model with the lowest AIC or BIC; verify with validation tests


In Python

- ▣ `pm.auto_arima()` from the `pmdarima` package
- ▣ Automatically tests stationarity, iterates over orders (p, q) , returns the best model

Caution

- ▣ Automatic selection is not the final answer \Rightarrow verify model validity
- ▣ Full Auto-ARIMA (including selection of d) \Rightarrow Chapter 3


Estimation Methods



The Three Main Approaches

- **Yule-Walker**: closed-form, AR only; equates sample autocorrelations with theoretical values
- **MLE**: most efficient and consistent; requires distributional assumption (Gaussian)
- **Conditional Least Squares**: compromise; minimizes sum of squared residuals

The Yule-Walker Equations for AR(p)



Yule-Walker Equations

$$\rho(1) = \phi_1 + \phi_2 \rho(1)$$

$$\rho(2) = \phi_1 \rho(1) + \phi_2$$

$$\text{Matrix form: } R \cdot \phi = \rho$$

R = autocorrelation matrix

$$\text{Solution: } \hat{\phi} = R^{-1} \rho$$

Main Idea

- Linear relationship between autocorrelations and AR parameters
- Allows closed-form estimation (no numerical optimization)

The Yule-Walker Equations: Matrix Form

Yule-Walker Equations for AR(p)

$$\square \quad \rho(k) = \phi_1 \rho(k-1) + \phi_2 \rho(k-2) + \cdots + \phi_p \rho(k-p), \quad k = 1, 2, \dots, p$$

Matrix Form

$$\square \quad \begin{pmatrix} \rho(0) & \rho(1) & \cdots & \rho(p-1) \\ \rho(1) & \rho(0) & \cdots & \rho(p-2) \\ \vdots & \vdots & \ddots & \vdots \\ \rho(p-1) & \rho(p-2) & \cdots & \rho(0) \end{pmatrix} \begin{pmatrix} \phi_1 \\ \phi_2 \\ \vdots \\ \phi_p \end{pmatrix} = \begin{pmatrix} \rho(1) \\ \rho(2) \\ \vdots \\ \rho(p) \end{pmatrix}$$

\square **Estimation:** Replace $\rho(k)$ with $\hat{\rho}(k)$; the Toeplitz matrix is symmetric and positive definite

Numerical Example: Yule-Walker for AR(2)

Sample Data ($T = 100$)

▣ **Estimated autocorrelations:** $\hat{\rho}(1) = 0.75$, $\hat{\rho}(2) = 0.65$

▶ Estimated variance: $\hat{\gamma}(0) = 4.0$

Step 1: Matrix System

▣ **Yule-Walker:** $R\hat{\phi} = \rho$

▶
$$\begin{pmatrix} 1 & 0.75 \\ 0.75 & 1 \end{pmatrix} \begin{pmatrix} \hat{\phi}_1 \\ \hat{\phi}_2 \end{pmatrix} = \begin{pmatrix} 0.75 \\ 0.65 \end{pmatrix}$$

Step 2: Solution (Cramer's Rule)

▣ $\det(R) = 1 - 0.75^2 = 0.4375$

▣ $\hat{\phi}_1 = \frac{0.75 \times 1 - 0.75 \times 0.65}{0.4375} = \frac{0.2625}{0.4375} = \boxed{0.600}$ $\hat{\phi}_2 = \frac{0.65 \times 1 - 0.75 \times 0.75}{0.4375} = \frac{0.0875}{0.4375} = \boxed{0.200}$

Step 3: Noise Variance

▣ $\hat{\sigma}^2 = \hat{\gamma}(0)(1 - \hat{\phi}_1\hat{\rho}(1) - \hat{\phi}_2\hat{\rho}(2)) = 4.0(1 - 0.45 - 0.13) = \boxed{1.68}$

Stationarity check: $\hat{\phi}_1 + \hat{\phi}_2 = 0.8 < 1 \checkmark$ $|\hat{\phi}_2| = 0.2 < 1 \checkmark$ $\hat{\phi}_2 - \hat{\phi}_1 = -0.4 > -1 \checkmark$

Proof: The Yule-Walker Equations

Goal: Derive $\rho(k) = \phi_1\rho(k-1) + \cdots + \phi_p\rho(k-p)$

- Start from AR(p): $X_t = \phi_1X_{t-1} + \cdots + \phi_pX_{t-p} + \varepsilon_t$
- Multiply by X_{t-k} and take expectations:
- $\mathbb{E}[X_tX_{t-k}] = \phi_1\mathbb{E}[X_{t-1}X_{t-k}] + \cdots + \phi_p\mathbb{E}[X_{t-p}X_{t-k}] + \mathbb{E}[\varepsilon_tX_{t-k}]$
- For $k \geq 1$: $\mathbb{E}[\varepsilon_tX_{t-k}] = 0 \Rightarrow \gamma(k) = \phi_1\gamma(k-1) + \cdots + \phi_p\gamma(k-p)$
- Dividing by $\gamma(0)$: $\rho(k) = \phi_1\rho(k-1) + \phi_2\rho(k-2) + \cdots + \phi_p\rho(k-p)$

Special Case AR(1)

- $\rho(k) = \phi_1\rho(k-1) = \phi_1^k$ (using $\rho(0) = 1$)

Maximum Likelihood Estimation

ARMA(p,q) Log-Likelihood (Gaussian errors: $\varepsilon_t \sim N(0, \sigma^2)$)

- ▣ $\ell(\phi, \theta, \sigma^2) = -\frac{n}{2} \ln(2\pi) - \frac{n}{2} \ln(\sigma^2) - \frac{1}{2\sigma^2} \sum_{t=1}^n \varepsilon_t^2$
- ▣ ε_t are the innovations computed recursively

Estimation Procedure

- ▣ Initialization: use method of moments or OLS for starting values
- ▣ Optimization: numerical methods (BFGS, Newton-Raphson)
- ▣ Iterate until convergence

In Practice

- ▣ `statsmodels.tsa.arima.model.ARIMA` \Rightarrow implements exact MLE with automatic initialization

Standard Errors and Inference

Asymptotic Distribution of MLE

- ▣ $\hat{\theta} \xrightarrow{d} N(\theta_0, \frac{1}{n}I(\theta_0)^{-1})$, where $I(\theta)$ is the **Fisher information matrix**
- ▣ $I(\theta) = -E\left[\frac{\partial^2 \ln L(\theta)}{\partial \theta \partial \theta'}\right] \Rightarrow$ average curvature of the log-likelihood
- ▣ Estimated variance-covariance matrix: $\hat{V} = \frac{1}{n}\hat{I}^{-1}$


What is the Standard Error (SE)?

- ▣ $SE(\hat{\theta}_j) = \sqrt{\hat{V}_{jj}} = \sqrt{\text{diag}_j\left(\frac{1}{n}\hat{I}^{-1}\right)} \Rightarrow$ measures estimation uncertainty
- ▣ **Example AR(1):** $SE(\hat{\phi}) \approx \sqrt{(1 - \hat{\phi}^2)/n}$; for $\hat{\phi} = 0.8$, $n = 100$: $SE \approx 0.06$
- ▣ **Interpretation:** small SE \Rightarrow parameter is estimated with high precision

Testing Parameter Significance

- ▣ $H_0 : \theta_j = 0$ Statistic: $z = \frac{\hat{\theta}_j}{SE(\hat{\theta}_j)} \sim N(0, 1)$ asymptotically
- ▣ Reject if $|z| > 1.96$ at 5% \Rightarrow **CI:** $\hat{\theta}_j \pm 1.96 \cdot SE(\hat{\theta}_j)$

Residual Diagnostics



If the model is correctly specified, residuals must be white noise


- ▣ **Residual plot:** random fluctuations around zero, constant variance
- ▣ **Residual ACF:** no significant spikes \Rightarrow white noise
- ▣ **Q-Q plot:** points on diagonal \Rightarrow normal; heavy tails \Rightarrow non-normal errors

Decision

- ▣ ✓ All checks OK \Rightarrow adequate model
- ✗ Not satisfied \Rightarrow return to identification

The Ljung-Box Test: Visual Illustration

Ljung-Box Test: good model vs inadequate model



Interpretation

- Left: good model \Rightarrow white noise residuals
- Right: inadequate model \Rightarrow residual autocorrelation \Rightarrow re-specification needed

The Ljung-Box Test

Definition 12 (Ljung-Box Test)

- ▣ Tests whether residuals are independently distributed (no autocorrelation)
- ▣ **Statistic:** $Q(m) = n(n+2) \sum_{k=1}^m \frac{\hat{\rho}_k^2}{n-k}$

Hypotheses and Distribution

- ▣ H_0 : Residuals are white noise; H_1 : Residuals are autocorrelated
- ▣ Under H_0 , $Q(m) \sim \chi^2(m-p-q)$ approximately

Decision

- ▣ **p-value** $> 0.05 \Rightarrow$ do not reject $H_0 \Rightarrow$ residuals are white noise
- ▣ **p-value** $< 0.05 \Rightarrow$ residual autocorrelation \Rightarrow inadequate model

Model Checklist

A Good ARMA Model Should Satisfy

- ▣ **Stationarity:** AR roots outside the unit circle (arroots)
- ▣ **Invertibility:** MA roots outside the unit circle (maroots)
- ▣ **White noise residuals:** No significant ACF (Ljung-Box test)
- ▣ **Normal residuals:** Q-Q plot, Jarque-Bera test
- ▣ **No heteroscedasticity:** Constant variance (ARCH test)
- ▣ **Simple:** Lowest AIC/BIC among adequate models

If Checks Are Not Satisfied

- ▣ Return to identification, try different orders

Point Forecasts

Optimal Forecast: $\hat{X}_{n+h|n} = \mathbb{E}[X_{n+h}|X_n, X_{n-1}, \dots]$

- The conditional expectation minimizes MSE

AR(1): $X_t = c + \phi X_{t-1} + \varepsilon_t$

- $\hat{X}_{n+1|n} = c + \phi X_n$; $\hat{X}_{n+h|n} = \mu + \phi^h(X_n - \mu)$
- Forecasts converge to the mean μ as $h \rightarrow \infty$ (mean reversion)

MA(1): $X_t = \mu + \varepsilon_t + \theta \varepsilon_{t-1}$

- $\hat{X}_{n+1|n} = \mu + \theta \varepsilon_n$; $\hat{X}_{n+h|n} = \mu$ for $h > 1$

Forecast Uncertainty

Mean Square Forecast Error (MSFE)

- **Error:** $e_{n+h|n} = X_{n+h} - \hat{X}_{n+h|n}$
- **MSFE:** $\text{MSFE}(h) = \sigma^2 \sum_{j=0}^{h-1} \psi_j^2$, where ψ_j are the $\text{MA}(\infty)$ coefficients


For AR(1): $\psi_j = \phi^j$

- $\text{MSFE}(h) = \sigma^2 \frac{1-\phi^{2h}}{1-\phi^2} \rightarrow \frac{\sigma^2}{1-\phi^2} = \text{Var}(X_t)$

Key Observation

- Forecast uncertainty increases with the horizon
- Converges to the unconditional variance $\text{Var}(X_t)$

ARMA Forecast with Confidence Intervals



Observation

- The confidence band widens with the horizon \Rightarrow convergence to the unconditional interval

Proof: MSFE for AR(1)

Claim

$$\square \text{ MSFE}(h) = \sigma^2 \frac{1 - \phi^{2h}}{1 - \phi^2} \quad \text{and} \quad \text{MSFE}(\infty) = \gamma(0)$$


Proof

- \square Forecast error at horizon h : $e_{n+h|n} = X_{n+h} - \hat{X}_{n+h|n}$
- \square By recursive substitution: $e_{n+h|n} = \sum_{j=0}^{h-1} \phi^j \varepsilon_{n+h-j}$
- \square $\text{MSFE}(h) = \mathbb{E}[e_{n+h|n}^2] = \sigma^2 \sum_{j=0}^{h-1} \phi^{2j} = \boxed{\sigma^2 \frac{1 - \phi^{2h}}{1 - \phi^2}}$
- \square Limit: $\text{MSFE}(\infty) = \frac{\sigma^2}{1 - \phi^2} = \gamma(0) \Rightarrow$ forecast converges to unconditional mean

Interpretation

- \square At long horizons, we do no better than the unconditional mean: $\text{CI} \rightarrow 2 \times 1.96 \sqrt{\gamma(0)}$



AR(1) Forecast: Mean Reversion



Properties

- ▣ Forecasts converge to the unconditional mean μ as the horizon increases
- ▣ Larger $|\phi| \Rightarrow$ slower reversion; CIs widen with the horizon

Forecast Error Variance by Horizon



Observation

- MSFE increases monotonically with horizon $h \Rightarrow$ convergence to $\text{Var}(X_t)$ (predictability limit)

Confidence Intervals for Forecasts


Formulas

- $X_{n+h}|X_n, \dots \sim N(\hat{X}_{n+h|n}, \text{MSFE}(h))$
- **CI** $(1 - \alpha)$: $\hat{X}_{n+h|n} \pm z_{\alpha/2} \cdot \sqrt{\text{MSFE}(h)}$, where $z_{\alpha/2} = 1.96$ for 95%

Properties

- Intervals widen as the horizon increases
 - ▶ Converge to the unconditional interval: $\mu \pm z_{\alpha/2} \sigma_X$
- Width depends on model parameters
 - ▶ Larger AR coefficients \Rightarrow wider intervals
- **Python**: `model.get_forecast(h).conf_int()`

Train/Validation/Test Forecast Example



Best Practice

- Always evaluate forecasts on data not used for estimation (train/validation/test split)

Forecast Evaluation


Out-of-Sample Testing

- Split data: training + test
- Generate forecasts on test
- Compare with actual values
- **Rolling window**: re-estimate as new data arrives

Error Metrics

- **MAE** = $\frac{1}{n} \sum |e_t|$
 - ▶ Robust to outliers
- **RMSE** = $\sqrt{\frac{1}{n} \sum e_t^2}$
 - ▶ Penalizes large errors
- **MAPE** = $\frac{100}{n} \sum \left| \frac{e_t}{X_t} \right|$
 - ▶ Percentage-based, interpretable

Rolling Window Forecast




Rolling Forecast Methodology

- **Fixed window** (last w obs.) vs **expanding** (all data); generate 1-step forecast, repeat

Rolling vs Multi-Step Forecast


Comparison: Rolling vs Multi-step vs Recursive



Key Differences

- Rolling 1-step (accurate); Multi-step direct (separate model/horizon); Recursive (error accumulation)

Real Data Application: Forecast Comparison



Practical Considerations


- Real data: non-stationarity, structural breaks; compare models; use rolling window validation

Workflow Summary

Box-Jenkins Methodology Steps

- 1. **Data preparation:** Check for missing values, outliers; transform if necessary
- 2. **Stationarity check:** Visual inspection, formal tests (ADF, KPSS); difference if non-stationary
- 3. **Model identification:** ACF/PACF patterns; grid search with information criteria
- 4. **Estimation and validation:** Estimate model, check significance; residual analysis, Ljung-Box test
- 5. **Forecasting:** Point forecasts with confidence intervals; out-of-sample validation

Case Study: Sunspots





Data Description

- Annual sunspots (1700–2008): stationary series with ~ 11 -year cycles; Box-Jenkins methodology

Step 1: ACF/PACF Analysis


ACF/PACF analysis for sunspots



Identification

- Sinusoidal ACF (AR); PACF with spikes at lags 1, 2, 9 \Rightarrow AR(2) or AR(9); stationary series ($d = 0$)


Step 2: Model Comparison



Model Selection

- Compare multiple candidate models using AIC; the **AR(9)** has the lowest AIC, capturing the 11-year solar cycle


Step 3: Model Diagnostics



AR(9) Diagnostics

- Residuals: white noise, zero mean, constant variance, ACF without structure, \approx normal

Step 4: Forecasting



Results

- AR(9) captures the cyclicity; 95% CI covers actual values; RMSE ≈ 30

Key Takeaways

Chapter Summary

- ▣ **AR(p)**: Depends on p past values; stationarity: roots outside the unit circle; PACF cuts off at lag p
- ▣ **MA(q)**: Depends on q past shocks; always stationary; ACF cuts off at lag q
- ▣ **ARMA(p,q)**: Combines AR and MA; both ACF and PACF decay
- ▣ **Box-Jenkins**: Identification \Rightarrow Estimation \Rightarrow Validation \Rightarrow Forecasting
- ▣ **Validation**: Residuals must be white noise
- ▣ **Forecasts**: Converge to the mean; uncertainty increases with the horizon

Next Chapter Preview

Chapter 3: ARIMA Models for Non-Stationary Data

- ▣ Non-stationarity: types, unit root tests (ADF, PP, KPSS)
- ▣ Differencing and the difference operator
- ▣ ARIMA(p,d,q): integrated models for non-stationary data
- ▣ The Auto-ARIMA algorithm: automatic model selection
- ▣ Case study: US GDP Forecasting

Reading

- ▣ Hyndman & Athanasopoulos, *Forecasting: Principles and Practice*, Ch. 9
- ▣ Box, Jenkins, Reinsel & Ljung, *Time Series Analysis*, Ch. 3-4

AI Exercise: Critical Thinking

Prompt to test in ChatGPT / Claude / Copilot

"Download monthly US Industrial Production Index from FRED (series INDPRO) for 2010-01 to 2024-12 (180 observations). Compute monthly log-differences (growth rates). Estimate an ARMA model, perform residual diagnostics, and forecast 12 months ahead. Give me complete Python code with plots."

Exercise:

1. Run the prompt in an LLM of your choice and critically analyze the response.
2. Does it verify stationarity *before* estimating ARMA? Justify.
3. How does it choose the orders p and q ? Does it use ACF/PACF or AIC/BIC?
4. Are residuals tested correctly? (Ljung-Box, Q-Q plot, heteroscedasticity)
5. Do forecast confidence intervals converge to the unconditional mean?

Warning: AI-generated code may run without errors and look professional. *That does not mean it is correct.*

Question 1


Question

□ For which value of ϕ is the AR(1) process $X_t = c + \phi X_{t-1} + \varepsilon_t$ stationary?

Answer Choices

- (A) $\phi = 1.2$
- (B) $\phi = 1.0$
- (C) $\phi = -0.8$
- (D) $\phi = -1.5$

Question 1: Answer



Answer: (C)

- ☐ AR(1) is stationary if and only if $|\phi| < 1$
- ☐ Only $|-0.8| = 0.8 < 1$

Question 2

Question

☐ You observe: ACF has a spike at lag 1, then cuts off. PACF decays gradually. What model?

Answer Choices


(A) AR(1)

(B) MA(1)

(C) ARMA(1,1)

(D) White noise

Question 2: Answer



Answer: (B)

- ACF cuts off \Rightarrow MA process
- PACF decays \Rightarrow confirms MA(1)

Question 3

Question



□ Is the MA(1) $X_t = \varepsilon_t + 1.5\varepsilon_{t-1}$ invertible?

Answer Choices

- (A) Yes, MA processes are always invertible
- (B) Yes, because $1.5 > 0$
- (C) No, because $|\theta| = 1.5 > 1$
- (D) No, MA processes are never invertible

Question 3: Answer

Invertibility of MA models



Answer: (C)

- Invertibility requires $|\theta| < 1$
- Here $|\theta| = 1.5 > 1$, so it is not invertible

Question 4

Question




□ The compact form $\phi(L)X_t = \theta(L)\varepsilon_t$ represents which model?

Answer Choices

- (A) Pure AR model
- (B) Pure MA model
- (C) ARMA model
- (D) None of the above

Question 4: Answer

ARMA(1,1): neither ACF nor PACF cut off



Answer: (C)

□ $\phi(L)$ is the AR polynomial, $\theta(L)$ is the MA polynomial \Rightarrow ARMA(p,q)

Question 5


Question

□ What is $(1 - L)^2 X_t$?

Answer Choices

- (A) $X_t - X_{t-1}$
- (B) $X_t - 2X_{t-1} + X_{t-2}$
- (C) $X_t + X_{t-1} + X_{t-2}$
- (D) $X_t - X_{t-2}$

Question 5: Answer



Answer: (B)

- ☐ $(1 - L)^2 = 1 - 2L + L^2$
- ☐ $(1 - L)^2 X_t = X_t - 2X_{t-1} + X_{t-2}$

Question 6


Question

□ Comparing ARMA(1,1) vs ARMA(2,1) using BIC, which is correct?

Answer Choices

- (A) Lower BIC always means better forecasts
- (B) BIC penalizes complexity less than AIC
- (C) The model with lower BIC is preferred
- (D) BIC can only compare models with the same number of parameters

Question 6: Answer



Answer: (C)

- Lower BIC indicates a better balance between estimation quality and complexity
- BIC penalizes complexity *more* than AIC

Question 7


Question

- ☐ After estimating an ARMA model, you run the Ljung-Box test on residuals and obtain $p\text{-value} = 0.03$. What does this mean?

Answer Choices

- (A) The model is adequate, residuals are white noise
- (B) The model is inadequate, residuals have autocorrelation
- (C) You need to increase the sample size
- (D) The test is inconclusive

Question 7: Answer



Answer: (B)

- ☐ p-value < 0.05 rejects H_0 (white noise)
- ☐ Indicates remaining residual autocorrelation

Question 8


Question

□ For a stationary AR(1) model, what happens to forecasts as the horizon $h \rightarrow \infty$?

Answer Choices

- (A) Forecasts increase without bound
- (B) Forecasts oscillate indefinitely
- (C) Forecasts converge to the unconditional mean μ
- (D) Forecasts become more precise

Question 8: Answer



Answer: (C)

$$\hat{X}_{n+h|n} = \mu + \phi^h(X_n - \mu) \rightarrow \mu \text{ as } h \rightarrow \infty \text{ (since } |\phi| < 1)$$

Question 9


Question

□ Consider an AR(1) process with $\phi = 0.6$ and $\sigma^2 = 4$. What is $\text{Var}(X_t)$?

Answer Choices

- (A) 4.0
- (B) 5.56
- (C) 6.25
- (D) 10.0

Question 9: Answer



Answer: (C)

- ▣ $\text{Var}(X_t) = \frac{\sigma^2}{1 - \phi^2} = \frac{4}{1 - 0.36} = \frac{4}{0.64} = 6.25$
- ▣ The process variance exceeds σ^2 due to persistence

Question 10

Question

□ Consider an MA(1) process with $\theta = 0.5$. What is $\rho(1)$?

Answer Choices



(A) 0.50

(B) 0.40

(C) 0.25

(D) 0.33

Question 10: Answer



Answer: (B)

- ☐ $\rho(1) = \frac{\theta}{1 + \theta^2} = \frac{0.5}{1 + 0.25} = \frac{0.5}{1.25} = 0.40$
- ☐ Note that $\rho(1) < \theta$ — the autocorrelation is **always** attenuated

Question 11

Question


Which statement about the ACF of an ARMA(1,1) process is **true**?

Answer Choices

- (A) It cuts off after lag 1
- (B) Exponential decay starting from lag 1, with $\rho(1) \neq \phi$
- (C) It is zero for all lags
- (D) It exactly follows the pattern ϕ^h for all $h \geq 0$

Question 11: Answer

ACF Comparison: AR(1) vs MA(1) vs ARMA(1,1)



Answer: (B)

- ☐ $\rho(1) = \frac{(1+\phi\theta)(\phi+\theta)}{1+2\phi\theta+\theta^2} \neq \phi$ (the MA component modifies lag 1)
- ☐ For $h \geq 2$: $\rho(h) = \phi \rho(h-1)$ — exponential decay as in AR(1)

Question 12


Question

□ An AR(2) process has $\phi_1 = 0.8$ and $\phi_2 = 0.3$. Is it stationary?

Answer Choices

- (A) Yes, it is stationary
- (B) No, because $\phi_1 + \phi_2 = 1.1 > 1$
- (C) Cannot be determined without data
- (D) Depends on the value of σ^2

Question 12: Answer



Answer: (B)

- Necessary conditions for AR(2) stationarity: $\phi_1 + \phi_2 < 1$, $\phi_2 - \phi_1 < 1$, $|\phi_2| < 1$
- Here $0.8 + 0.3 = 1.1 > 1 \Rightarrow$ the first condition is violated

Question 13


Question

□ What does the Wold decomposition theorem guarantee?

Answer Choices

- (A) Any time series is an AR process
- (B) Any stationary process can be written as $MA(\infty)$: $X_t = \sum_{j=0}^{\infty} \psi_j \varepsilon_{t-j}$
- (C) Any process has finite variance
- (D) ARMA models are always invertible

Question 13: Answer



Answer: (B)

- Wold's theorem: $X_t = \sum_{j=0}^{\infty} \psi_j \varepsilon_{t-j} + D_t$, where D_t is the deterministic component
- This justifies ARMA models: they are parsimonious approximations of $MA(\infty)$

Question 14


Question

□ AR(1) with $\phi = 0.9$, $\sigma^2 = 1$. What happens to the CI width as $h \rightarrow \infty$?

Answer Choices

- (A) It remains constant
- (B) It decreases to zero
- (C) It grows toward $2 \times 1.96 \times \sqrt{1/(1 - 0.81)} \approx 9.0$
- (D) It grows to infinity

Question 14: Answer



Answer: (C)

- ▣ $\text{MSFE}(\infty) = \frac{\sigma^2}{1-\phi^2} = \frac{1}{1-0.81} = \frac{1}{0.19} \approx 5.26$
- ▣ $\text{CI width} = 2 \times 1.96 \sqrt{5.26} \approx 2 \times 1.96 \times 2.29 \approx 9.0$

Data Sources and Software

Software Packages

- ▣ `statsmodels` \Rightarrow Statistical models for Python, including ARIMA
- ▣ `pmdarima` \Rightarrow Automatic ARIMA selection for Python
- ▣ `scipy` \Rightarrow Optimization and statistical functions
- ▣ `numpy`, `pandas` \Rightarrow Data manipulation
- ▣ `matplotlib` \Rightarrow Visualization

Data and Examples

- ▣ Simulated time series for illustrations
- ▣ Examples based on Hyndman & Athanasopoulos (2021)

Bibliography I

Fundamental ARMA Works

- Box, G.E.P., & Jenkins, G.M. (1970). *Time Series Analysis: Forecasting and Control*, Holden-Day.
- Akaike, H. (1974). A New Look at the Statistical Model Identification, *IEEE Transactions on Automatic Control*, 19(6), 716–723.
- Schwarz, G. (1978). Estimating the Dimension of a Model, *The Annals of Statistics*, 6(2), 461–464.

Diagnostics and Validation

- Ljung, G.M., & Box, G.E.P. (1978). On a Measure of Lack of Fit in Time Series Models, *Biometrika*, 65(2), 297–303.
- Brockwell, P.J., & Davis, R.A. (2016). *Introduction to Time Series and Forecasting*, 3rd ed., Springer.

Bibliography II

Textbooks and Additional References

- ▣ Hamilton, J.D. (1994). *Time Series Analysis*, Princeton University Press.
- ▣ Shumway, R.H., & Stoffer, D.S. (2017). *Time Series Analysis and Its Applications*, 4th ed., Springer.
- ▣ Hyndman, R.J., & Athanasopoulos, G. (2021). *Forecasting: Principles and Practice*, 3rd ed., OTexts.


Online Resources and Code

- ▣ **Quantlet**: <https://quantlet.com> – Code platform for quantitative methods
- ▣ **Quantinar**: <https://quantinar.com> – Learning platform for quantitative methods
- ▣ **GitHub TSA**: https://github.com/QuantLet/TSA/tree/main/TSA_ch2 – Python code for this chapter

Thank You!

Questions?

Course materials available at: <https://danpele.github.io/Time-Series-Analysis/>



Quantlet



Quantinar