



# Time Series Analysis and Forecasting

## Chapter 5: GARCH and Volatility



Daniel Traian PELE

Bucharest University of Economic Studies

IDA Institute Digital Assets

Blockchain Research Center

AI4EFin Artificial Intelligence for Energy Finance

Romanian Academy, Institute for Economic Forecasting

MSCA Digital Finance

## Learning Objectives

By the end of this chapter, you will be able to:

- Understand volatility clustering and its importance in financial data
- Model conditional heteroskedasticity using ARCH and GARCH models
- Estimate GARCH models and interpret their parameters
- Forecast volatility and apply it to risk management

## Outline

- Introduction to Volatility Modeling
- The ARCH Model
- The GARCH Model
- Asymmetric GARCH Models
- Model Selection and Diagnostics
- Volatility Forecasting
- Case Study: S&P 500
- Summary

## Why Model Volatility?


### Empirical Observations in Financial Series

- Financial returns exhibit **volatility clustering** — periods of high volatility tend to be followed by periods of high volatility
- The distribution of returns has **fat tails** (leptokurtosis)
- Return correlation is nearly zero, but correlation of squares is significant
- Volatility responds **asymmetrically** to shocks (leverage effect)

### Limitation of ARIMA Models


ARIMA models assume **constant variance** (homoskedasticity), which is not realistic for financial series!

## Volatility Clustering



- High volatility periods are followed by high volatility; calm by calm
- This suggests that **conditional variance** is predictable

## Stylized Facts of Financial Returns



### Observed Properties

1. **No autocorrelation** in returns
2. **Autocorrelation** in  $r_t^2$ ,  $|r_t|$
3. **Fat tails** (kurtosis  $> 3$ )
4. **Leverage effect**
5. **Volatility clustering**

## Conditional Heteroskedasticity

### Definition 1 (Conditional Variance)

For return series  $\{r_t\}$ , the **conditional variance** at time  $t$  is:  $\sigma_t^2 = \text{Var}(r_t | \mathcal{F}_{t-1}) = \mathbb{E}[(r_t - \mu_t)^2 | \mathcal{F}_{t-1}]$  where  $\mathcal{F}_{t-1}$  is the information available up to time  $t - 1$ .

### General Model

$$r_t = \mu_t + \varepsilon_t, \quad \varepsilon_t = \sigma_t z_t, \quad z_t \sim \text{i.i.d.}(0, 1)$$

- $\mu_t$  = conditional mean (ARMA);  $\sigma_t^2$  = conditional variance (GARCH)
- $z_t$  = standardized innovations (Normal, Student-t, GED)

## The ARCH( $q$ ) Model — Engle (1982)

### Definition 2 (ARCH( $q$ ))

The **Autoregressive Conditional Heteroskedasticity** model of order  $q$ :

$$\varepsilon_t = \sigma_t z_t, \quad z_t \sim \text{i.i.d.}(0, 1), \quad \sigma_t^2 = \omega + \sum_{i=1}^q \alpha_i \varepsilon_{t-i}^2$$

### Stationarity Restrictions


- ▣  $\omega > 0$  (positive base variance),  $\alpha_i \geq 0$  (non-negativity)
- ▣  $\sum_{i=1}^q \alpha_i < 1$  (stationarity)

### Remark 1

Robert Engle received the **Nobel Prize in Economics** in 2003 for developing the ARCH model!



## ARCH(1) Simulation: Effect of $\alpha$ Parameter



Higher  $\alpha$  means volatility reacts more strongly to recent shocks.

## Properties of the ARCH(1) Model

$$\text{ARCH}(1): \sigma_t^2 = \omega + \alpha_1 \varepsilon_{t-1}^2$$

- ▣ **Unconditional variance:**  $\mathbb{E}[\varepsilon_t^2] = \frac{\omega}{1 - \alpha_1}$  (if  $\alpha_1 < 1$ )
- ▣ **Kurtosis:**  $\kappa = 3 \cdot \frac{1 - \alpha_1^2}{1 - 3\alpha_1^2}$  (if  $\alpha_1^2 < 1/3$ )
- ▣ Kurtosis  $> 3$  for  $\alpha_1 > 0 \Rightarrow$  **fat tails!**

### Numerical Example

If  $\omega = 0.0001$  and  $\alpha_1 = 0.3$ :

- ▣ Unconditional variance:  $\sigma^2 = \frac{0.0001}{1-0.3} = 0.000143$
- ▣ Kurtosis:  $\kappa = 3 \cdot \frac{1-0.09}{1-0.27} = 3.74 > 3$

## Derivation: Unconditional Variance of ARCH(1)

### Derivation.

Let  $\varepsilon_t = \sigma_t z_t$  with  $z_t \sim N(0, 1)$  and  $\sigma_t^2 = \omega + \alpha_1 \varepsilon_{t-1}^2$ .

**Step 1:** Take unconditional expectation:

$$\mathbb{E}[\varepsilon_t^2] = \mathbb{E}[\sigma_t^2 z_t^2] = \mathbb{E}[\sigma_t^2] \cdot \mathbb{E}[z_t^2] = \mathbb{E}[\sigma_t^2]$$

**Step 2:** Apply expectation to variance equation:

$$\mathbb{E}[\sigma_t^2] = \mathbb{E}[\omega + \alpha_1 \varepsilon_{t-1}^2] = \omega + \alpha_1 \mathbb{E}[\varepsilon_{t-1}^2]$$

**Step 3:** By stationarity,  $\mathbb{E}[\varepsilon_t^2] = \mathbb{E}[\varepsilon_{t-1}^2] = \sigma^2$ :

$$\sigma^2 = \omega + \alpha_1 \sigma^2 \quad \Rightarrow \quad \sigma^2(1 - \alpha_1) = \omega$$

**Result:**  $\boxed{\sigma^2 = \frac{\omega}{1 - \alpha_1}}$  (requires  $\alpha_1 < 1$  for stationarity)



## Derivation: Kurtosis of ARCH(1)

For  $\varepsilon_t = \sigma_t z_t$  with  $z_t \sim N(0, 1)$ :

**Step 1:**  $\mathbb{E}[\varepsilon_t^4] = \mathbb{E}[\sigma_t^4] \cdot \mathbb{E}[z_t^4] = 3\mathbb{E}[\sigma_t^4]$  (since  $\mathbb{E}[z^4] = 3$ )

**Step 2:** Using  $\sigma_t^2 = \omega + \alpha_1 \varepsilon_{t-1}^2$ :

$$\mathbb{E}[\sigma_t^4] = \mathbb{E}[(\omega + \alpha_1 \varepsilon_{t-1}^2)^2] = \omega^2 + 2\omega\alpha_1\sigma^2 + \alpha_1^2\mathbb{E}[\varepsilon_{t-1}^4]$$

**Step 3:** Solving the recursion yields:

$$\kappa = \frac{\mathbb{E}[\varepsilon_t^4]}{(\mathbb{E}[\varepsilon_t^2])^2} = 3 \cdot \frac{1 - \alpha_1^2}{1 - 3\alpha_1^2}$$

### Interpretation

- ▣  $\kappa > 3$  for any  $\alpha_1 > 0 \Rightarrow$  **fat tails** (leptokurtosis)
- ▣ Requires  $\alpha_1 < 0.577$  for finite fourth moment
- ▣ ARCH naturally generates heavy-tailed distributions!

## Testing for ARCH Effects

### Engle's Test for ARCH Effects

#### Procedure:

1. Estimate the mean model and obtain residuals  $\hat{\varepsilon}_t$
2. Calculate  $\hat{\varepsilon}_t^2$
3. Regress  $\hat{\varepsilon}_t^2$  on its lags:

$$\hat{\varepsilon}_t^2 = \beta_0 + \beta_1 \hat{\varepsilon}_{t-1}^2 + \cdots + \beta_q \hat{\varepsilon}_{t-q}^2 + u_t$$

4. Calculate the statistic  $LM = T \cdot R^2 \sim \chi^2(q)$

### Hypotheses

- $H_0$ : No ARCH effects ( $\alpha_1 = \cdots = \alpha_q = 0$ )
- $H_1$ : ARCH effects present (at least one  $\alpha_i \neq 0$ )

## Limitations of the ARCH Model

### Practical Problems

1. **High order** — many lags are usually needed (large  $q$ )
2. **Many parameters** — estimation difficulties
3. **Non-negativity constraints** — difficult to impose for large  $q$
4. **Does not capture persistence** — observed volatility is very persistent

### The Solution

**The GARCH Model** — introduces lags of conditional variance to capture persistence with fewer parameters!

## The GARCH(p,q) Model — Bollerslev (1986)

### Definition 3 (GARCH(p,q))

The **Generalized ARCH** model:

$$\varepsilon_t = \sigma_t z_t, \quad z_t \sim \text{i.i.d.}(0, 1)$$
$$\sigma_t^2 = \omega + \sum_{i=1}^q \alpha_i \varepsilon_{t-i}^2 + \sum_{j=1}^p \beta_j \sigma_{t-j}^2$$


### Interpretation

- $\omega$  = base level of volatility
- $\alpha_i$  = reaction to recent shocks (news coefficients)
- $\beta_j$  = volatility persistence (memory)
- $\alpha + \beta$  = total persistence

## The GARCH(1,1) Model

### The Most Popular Volatility Model

$$\sigma_t^2 = \omega + \alpha \varepsilon_{t-1}^2 + \beta \sigma_{t-1}^2$$




### Restrictions & Properties

□  $\omega > 0, \alpha \geq 0, \beta \geq 0; \quad \alpha + \beta < 1$  (stationarity)

□  $\bar{\sigma}^2 = \frac{\omega}{1-\alpha-\beta}; \quad \text{Half-life: } HL = \frac{\ln(0.5)}{\ln(\alpha+\beta)}$



## GARCH(1,1) Simulation: Persistence Effect



Parameter  $\alpha$  controls reaction to shocks,  $\beta$  controls persistence. The sum  $\alpha + \beta$  determines mean-reversion speed.

## Derivation: Unconditional Variance of GARCH(1,1)

### Derivation.

For  $\sigma_t^2 = \omega + \alpha \varepsilon_{t-1}^2 + \beta \sigma_{t-1}^2$ :

**Step 1:** Take unconditional expectation:  $\mathbb{E}[\sigma_t^2] = \omega + \alpha \mathbb{E}[\varepsilon_{t-1}^2] + \beta \mathbb{E}[\sigma_{t-1}^2]$

**Step 2:** By stationarity,  $\mathbb{E}[\sigma_t^2] = \mathbb{E}[\sigma_{t-1}^2] = \bar{\sigma}^2$  and  $\mathbb{E}[\varepsilon_t^2] = \bar{\sigma}^2$ :  $\bar{\sigma}^2 = \omega + (\alpha + \beta)\bar{\sigma}^2$

**Step 3:** Solve:  $\bar{\sigma}^2(1 - \alpha - \beta) = \omega \Rightarrow \bar{\sigma}^2 = \frac{\omega}{1 - \alpha - \beta}$



### Stationarity Condition

Requires  $\alpha + \beta < 1$  for finite unconditional variance.



## GARCH(1,1) as ARMA for $\varepsilon_t^2$

### ARMA(1,1) Representation

Define  $\nu_t = \varepsilon_t^2 - \sigma_t^2$  (variance shock). Then:

$$\varepsilon_t^2 = \omega + (\alpha + \beta)\varepsilon_{t-1}^2 + \nu_t - \beta\nu_{t-1}$$

This is an **ARMA(1,1)** for  $\varepsilon_t^2$ !

### Implications

- ACF of  $\varepsilon_t^2$  decays exponentially (like ARMA)
- Persistence is given by  $\alpha + \beta$
- PACF can help identify the order

## Derivation: ARMA Representation of GARCH(1,1)

### Derivation.

**Step 1:** Define variance shock:  $\nu_t = \varepsilon_t^2 - \sigma_t^2$

□  $\mathbb{E}[\nu_t | \mathcal{F}_{t-1}] = \mathbb{E}[\varepsilon_t^2 | \mathcal{F}_{t-1}] - \sigma_t^2 = \sigma_t^2 - \sigma_t^2 = 0$

□  $\nu_t$  is a martingale difference sequence

**Step 2:** Substitute  $\sigma_t^2 = \varepsilon_t^2 - \nu_t$  into GARCH equation:

$$\varepsilon_t^2 - \nu_t = \omega + \alpha \varepsilon_{t-1}^2 + \beta (\varepsilon_{t-1}^2 - \nu_{t-1})$$

**Step 3:** Rearrange:

$$\varepsilon_t^2 = \omega + (\alpha + \beta) \varepsilon_{t-1}^2 + \nu_t - \beta \nu_{t-1}$$

**Result:** ARMA(1,1) with AR coefficient  $\phi = \alpha + \beta$  and MA coefficient  $\theta = -\beta$ . □

## Volatility Persistence and Half-Life

### Persistence

$\alpha + \beta$  measures mean reversion speed:

- $\approx 1$ : very persistent
- $\ll 1$ : quick reversion

### Half-Life Formula

$$HL = \frac{-0.693}{\ln(\alpha + \beta)}$$

### Example: S&P 500

$$\alpha = 0.09, \beta = 0.90$$

$$\alpha + \beta = 0.99$$

$$HL = \frac{-0.693}{\ln(0.99)} \approx 69$$

Shock halves in  $\sim 69$  trading days!

## Estimation of GARCH Models

### Maximum Likelihood Estimation (MLE)

Log-likelihood (normal):  $\ell(\theta) = -\frac{T}{2} \ln(2\pi) - \frac{1}{2} \sum_{t=1}^T \left[ \ln(\sigma_t^2) + \frac{\varepsilon_t^2}{\sigma_t^2} \right]$

### Alternative Distributions for $z_t$

- **Student-t**: captures fat tails — most common choice
- **GED**: flexibility for kurtosis
- **Skewed Student-t**: asymmetry and fat tails

### Practical Note

Student-t distribution typically provides better fit for financial returns due to fat tails (kurtosis  $> 3$ ).

## Typical Values for GARCH(1,1)

Series	$\alpha$	$\beta$	$\alpha + \beta$
S&P 500 daily	0.05–0.10	0.85–0.95	0.95–0.99
EUR/USD daily	0.03–0.08	0.90–0.95	0.95–0.99
Bitcoin daily	0.10–0.20	0.75–0.85	0.90–0.98
Bonds	0.02–0.05	0.90–0.97	0.95–0.99

## Observations

- $\alpha + \beta$  close to 1  $\Rightarrow$  **very persistent volatility**
- Small  $\alpha$ , large  $\beta \Rightarrow$  slow reaction to shocks, long memory
- Bitcoin: larger  $\alpha \Rightarrow$  faster reaction to news

## IGARCH — Integrated GARCH

### Definition 4 (IGARCH(1,1))

When  $\alpha + \beta = 1$ :

$$\sigma_t^2 = \omega + \alpha \varepsilon_{t-1}^2 + (1 - \alpha) \sigma_{t-1}^2$$

### Properties


- Unconditional variance does not exist (infinite)
- Shocks have **permanent** effect on volatility
- Used for series with extreme persistence
- Useful for **RiskMetrics** (J.P. Morgan):  $\alpha = 0.06$ ,  $\beta = 0.94$

### Remark 2

IGARCH is analogous to a unit root in variance!




## GARCH vs IGARCH: Persistence Comparison



Standard GARCH reverts to unconditional mean, while IGARCH has no finite mean and shocks persist indefinitely.

## Leverage Effect



### Definition

**Leverage effect:** Negative shocks increase volatility **more** than positive shocks of the same magnitude.

### Problem with GARCH

Standard GARCH:  $\sigma_t^2 = \omega + \alpha \varepsilon_{t-1}^2 + \beta \sigma_{t-1}^2$  — only  $\varepsilon_{t-1}^2$  matters, sign is lost! Economic intuition: Bad news  $\Rightarrow$  stock price falls  $\Rightarrow$  debt/equity ratio rises  $\Rightarrow$  volatility increases.

## The EGARCH Model — Nelson (1991)

### Definition 5 (EGARCH(1,1))

**Exponential GARCH:**


$$\ln(\sigma_t^2) = \omega + \alpha (|z_{t-1}| - \mathbb{E}[|z_{t-1}|]) + \gamma z_{t-1} + \beta \ln(\sigma_{t-1}^2)$$

where  $z_t = \varepsilon_t / \sigma_t$ .

### EGARCH Advantages


- **No non-negativity constraints required** — models  $\ln(\sigma_t^2)$
- **Captures leverage effect** through parameter  $\gamma$ 
  - ▶  $\gamma < 0$ : negative shocks  $\Rightarrow$  higher volatility
  - ▶  $\gamma = 0$ : symmetric effect (like GARCH)
- Persistence is given by  $\beta$

## EGARCH Simulation: Symmetric vs Asymmetric



When  $\gamma < 0$ , negative shocks (bad news) increase volatility more than positive shocks of the same magnitude.

## News Impact Curve — EGARCH



### Definition

**News Impact Curve:**  $\sigma_{t+1}^2$  as function of  $\varepsilon_t$ , holding  $\sigma_t^2$  constant.

- **GARCH:** Symmetric V-shape (parabola); **EGARCH:** Asymmetric — steeper for negative shocks; **GJR:** Piecewise linear with kink at zero
- The asymmetry captures the leverage effect: bad news has larger impact on volatility than good news.

## The GJR-GARCH Model

### Definition 6 (GJR-GARCH(1,1))

Glosten, Jagannathan & Runkle (1993):  $\sigma_t^2 = \omega + \alpha \varepsilon_{t-1}^2 + \gamma \varepsilon_{t-1}^2 \cdot I_{t-1} + \beta \sigma_{t-1}^2$  where  $I_{t-1} = 1$  if  $\varepsilon_{t-1} < 0$ , else 0.

### Interpretation

- Positive shocks: impact =  $\alpha$ ;    Negative shocks: impact =  $\alpha + \gamma$
- Leverage effect present if  $\gamma > 0$
- Stationarity:  $\alpha + \gamma/2 + \beta < 1$

## TGARCH — Threshold GARCH


### Definition 7 (TGARCH(1,1))

Zakoian (1994) models standard deviation:  $\sigma_t = \omega + \alpha^+ \varepsilon_{t-1}^+ + \alpha^- \varepsilon_{t-1}^- + \beta \sigma_{t-1}$

### Comparison of Asymmetric Models

Model	Specification	Leverage
GARCH	$\sigma_t^2$	No
EGARCH	$\ln(\sigma_t^2)$	Yes ( $\gamma < 0$ )
GJR-GARCH	$\sigma_t^2$ with indicator	Yes ( $\gamma > 0$ )
TGARCH	$\sigma_t$	Yes ( $\alpha^- > \alpha^+$ )


## GJR-GARCH/TGARCH Simulation



GJR-GARCH adds an indicator term to capture asymmetric response to negative shocks.




## News Impact Curves Comparison



Standard GARCH is symmetric, while EGARCH and GJR-GARCH capture asymmetry (leverage effect).

## GARCH Family Comparison



All models capture volatility clustering, but differ in how they model asymmetry.

## GARCH-M: GARCH-in-Mean Model

### Definition 8 (GARCH-M)

The **GARCH-in-Mean** model:  $r_t = \mu + \lambda\sigma_t + \varepsilon_t$ , where  $\lambda$  is the **risk premium**.

### Interpretation

- ▣  $\lambda > 0$ : Higher risk  $\Rightarrow$  higher expected return
- ▣  $\lambda = 0$ : Reduces to standard GARCH
- ▣  $\lambda < 0$ : Higher risk  $\Rightarrow$  lower return (rare)

### Financial Intuition

Investors demand compensation for bearing risk — GARCH-M captures this **risk-return tradeoff**.

## GARCH-M: Alternative Specifications

### Common Specifications

Risk premium can enter in different forms: (1)  $r_t = \mu + \lambda\sigma_t + \varepsilon_t$ ; (2)  $r_t = \mu + \lambda\sigma_t^2 + \varepsilon_t$ ; (3)  $r_t = \mu + \lambda \ln(\sigma_t^2) + \varepsilon_t$

### Typical Results for Equity Markets

- ▣ Estimated  $\lambda$  often positive but small (0.01–0.10)
- ▣ Significance varies across markets and periods
- ▣ Variance specification yields larger  $\lambda$  estimates

### Remark 3

GARCH-M is used in asset pricing, portfolio optimization, and CAPM testing.

## Order Selection

### Information Criteria

- **AIC** =  $-2\ell + 2k$
- **BIC** =  $-2\ell + k \ln(T)$
- **HQIC** =  $-2\ell + 2k \ln(\ln(T))$

where  $\ell$  = maximized log-likelihood,  $k$  = number of parameters.

### Practical Recommendations

- GARCH(1,1) is sufficient in **90% of cases**
- Check if asymmetric model significantly improves fit
- Choose innovation distribution that minimizes AIC/BIC

## GARCH Model Diagnostics

### Standardized Residuals


$$\hat{z}_t = \frac{\hat{\varepsilon}_t}{\hat{\sigma}_t}$$

If the model is correctly specified,  $\hat{z}_t$  should be i.i.d.(0,1).

### Diagnostic Checks

1. **Ljung-Box on  $\hat{z}_t$ :** check absence of autocorrelation in mean
2. **Ljung-Box on  $\hat{z}_t^2$ :** check absence of residual ARCH effects
3. **ARCH-LM test on  $\hat{z}_t$ :** confirm absence of heteroskedasticity
4. **Histogram + QQ-plot:** verify assumed distribution

## Diagnostic Example



## Forecasting with GARCH(1,1)

### One-Step-Ahead Forecast

$$\hat{\sigma}_{T+1}^2 = \omega + \alpha \varepsilon_T^2 + \beta \sigma_T^2$$

### Multi-Step Forecast


For  $h > 1$ :  $\mathbb{E}_T[\sigma_{T+h}^2] = \bar{\sigma}^2 + (\alpha + \beta)^{h-1}(\sigma_{T+1}^2 - \bar{\sigma}^2)$  where  $\bar{\sigma}^2 = \frac{\omega}{1-\alpha-\beta}$  = unconditional variance.

### Convergence

$\lim_{h \rightarrow \infty} \mathbb{E}_T[\sigma_{T+h}^2] = \bar{\sigma}^2$  — forecast converges to unconditional variance!



## Volatility Forecast — Visualization



- Forecast converges exponentially to  $\bar{\sigma}^2$ ; speed depends on  $\alpha + \beta$
- The closer  $\alpha + \beta$  is to 1, the slower the convergence

## Applications of Volatility Forecasting

### Value at Risk (VaR)

$$\text{VaR}_\alpha = z_\alpha \cdot \sigma_{T+1}$$

Maximum loss with probability  $1 - \alpha$ .

### Expected Shortfall (ES)


$$\text{ES}_\alpha = \mathbb{E}[-r | r < -\text{VaR}_\alpha]$$

Average loss when VaR is exceeded.

### Other Applications


- Option pricing
- Dynamic hedging
- Portfolio allocation
- Stress testing

## VaR and ES: Graphical Illustration



VaR 1% = loss exceeded only in 1% of cases. Red area = left tail (extreme losses).

## VaR vs Expected Shortfall: Normal vs Student-t



ES (green line) measures average loss when VaR is exceeded. Student-t has heavier tails  $\Rightarrow$  larger VaR and ES.

## Value at Risk — Numerical Example

### VaR Calculation

Portfolio: **1,000,000 EUR**, forecasted volatility  $\hat{\sigma}_{T+1} = 1.5\%$

### VaR with Normal Distribution

Level	$z_{\alpha}$	VaR (%)	VaR (EUR)
5% (1 day)	1.645	2.47%	24,675
1% (1 day)	2.326	3.49%	34,890

### Scaling for Longer Periods

$\text{VaR}_{h \text{ days}} = \text{VaR}_{1 \text{ day}} \cdot \sqrt{h}$  — assumes i.i.d. returns

## Value at Risk — Student-t Distribution

### Why Student-t?

Normal distribution **underestimates** tail risk. Student-t with  $\nu$  degrees of freedom better captures fat tails (kurtosis  $> 3$ ).

VaR 1% (1 day) Comparison:  $\sigma = 1.5\%$ , Portfolio = 1M EUR

Distribution	Quantile	VaR (EUR)
Normal	2.326	34,890
Student-t ( $\nu = 6$ )	3.143	47,145
Student-t ( $\nu = 4$ )	3.747	56,205

### Observation

With  $\nu = 6$  (typical for stocks), VaR is **35% higher** than normal!

## VaR — Complete Example with GARCH

### VaR Calculation Procedure


1. Estimate GARCH(1,1) model with Student-t distribution
2. Obtain volatility forecast:  $\hat{\sigma}_{T+1}$
3. Calculate VaR:  $\text{VaR}_\alpha = t_\alpha(\nu) \cdot \hat{\sigma}_{T+1} \cdot \sqrt{\frac{\nu-2}{\nu}}$

### Example: S&P 500

- ▣ Estimated parameters:  $\alpha = 0.088$ ,  $\beta = 0.900$ ,  $\nu = 6.4$
- ▣ Forecasted volatility:  $\hat{\sigma}_{T+1} = 1.2\%$
- ▣ Portfolio: 10,000,000 EUR

**VaR 1% (1 day):**  $\text{VaR} = 3.05 \times 0.012 \times 10,000,000 = \mathbf{366,000 \text{ EUR}}$


## VaR Backtesting: Visual Overview



Backtesting checks if VaR violations match expected rate (e.g., 2.5 violations/year for VaR 1%).



## Volatility Forecast with Confidence Intervals



Forecast converges to unconditional volatility  $\bar{\sigma}$ . Uncertainty increases with forecast horizon.

## VaR Backtesting: Kupiec Test

### Unconditional Coverage Test

Tests whether the observed violation rate equals the expected rate  $p$  (e.g., 1% for VaR 1%).

Let  $N$  = number of VaR violations,  $T$  = total observations,  $\hat{p} = N/T$ .

**Likelihood Ratio Statistic:**

$$LR_{uc} = -2 \ln \left[ \frac{(1-p)^{T-N} p^N}{(1-\hat{p})^{T-N} \hat{p}^N} \right] \sim \chi^2(1)$$

### Hypotheses

- $H_0: \hat{p} = p$  (VaR model is correctly calibrated)
- $H_1: \hat{p} \neq p$  (VaR model under- or over-estimates risk)

## VaR Backtesting: Christoffersen Test

### Conditional Coverage Test

Tests both **unconditional coverage** and **independence** of violations.  
Violations should be independent — no clustering of exceptions!

### Test Components

- ▣ **Independence test** ( $LR_{ind}$ ): Tests if violations are serially independent
- ▣ **Conditional coverage**:  $LR_{cc} = LR_{uc} + LR_{ind} \sim \chi^2(2)$

### Interpretation

Reject  $LR_{uc}$ : wrong frequency; Reject  $LR_{ind}$ : clustered violations; Reject  $LR_{cc}$ : model fails

## VaR Backtesting: Python Implementation

### Kupiec Test Implementation

```
import numpy as np
from scipy import stats
def kupiec_test(violations, T, p=0.01):
    N = np.sum(violations)
    p_hat = N / T
    if N == 0 or N == T:
        return np.nan, np.nan
    LR = -2 * (np.log((1-p)**(T-N) * p**N) -
               np.log((1-p_hat)**(T-N) * p_hat**N))
    return LR, 1 - stats.chi2.cdf(LR, df=1)
```

### Usage

```
LR, pval = kupiec_test(violations, T=250, p=0.01)
```

## ARMA-GARCH: Joint Mean and Variance Modeling

### Why Joint Modeling?

**Serial correlation**  $\Rightarrow$  ARMA for mean; **Volatility clustering**  $\Rightarrow$  GARCH for variance.

### Definition 9 (ARMA(p,q)-GARCH(r,s))

**Mean equation:**  $r_t = \mu + \sum_{i=1}^p \phi_i (r_{t-i} - \mu) + \sum_{j=1}^q \theta_j \varepsilon_{t-j} + \varepsilon_t$

**Variance equation:**  $\varepsilon_t = \sigma_t z_t, \quad \sigma_t^2 = \omega + \sum_{i=1}^r \alpha_i \varepsilon_{t-i}^2 + \sum_{j=1}^s \beta_j \sigma_{t-j}^2$

## ARMA-GARCH: Model Selection Strategy

### Step-by-Step Approach

1. **Identify mean model:** Check ACF/PACF of returns for ARMA structure
2. **Test for ARCH effects:** Apply ARCH-LM test to residuals
3. **Specify variance model:** Usually GARCH(1,1) is sufficient
4. **Joint estimation:** Estimate both equations via MLE
5. **Diagnostic checking:** Standardized residuals should be i.i.d.

### Common Specifications

- ☐ **Stock returns:** AR(1)-GARCH(1,1) or ARMA(1,1)-GARCH(1,1)
- ☐ **Exchange rates:** Often just GARCH(1,1) (no mean dynamics)
- ☐ **Interest rates:** AR(1)-EGARCH(1,1) for leverage effects

## ARMA-GARCH: Python Implementation

### Using the arch Package

```
from arch import arch_model
model = arch_model(returns,
                   mean='ARX',
                   lags=1,
                   vol='Garch',
                   p=1, q=1,
                   dist='t')
result = model.fit(dispen='off')
print(result.summary())
```

### Parameters

mean='ARX': ARMA mean; lags=1: AR(1); dist='t': Student-t

## ARMA-GARCH: Complete Example


```
from arch import arch_model
model = arch_model(returns,
                   mean='ARX', lags=[1],
                   vol='EGARCH', p=1, q=1,
                   dist='skewt')
result = model.fit(update_freq=0)
cond_mean = result.conditional_mean
cond_vol = result.conditional_volatility
forecasts = result.forecast(horizon=5)
```

### Note

For MA terms, use mean='HARX' or pre-filter with statsmodels ARIMA.




## S&P 500 Volatility Analysis



- S&P 500 daily returns (2000–2024) — volatility clustering visible
- Crisis periods: 2008 (financial), 2020 (COVID-19), 2022 (inflation)

## GARCH(1,1) Estimation — S&amp;P 500




## Estimation Results

Parameter	Value
$\omega$	0.0108
$\alpha$	0.0883
$\beta$	0.9002
$\alpha + \beta$	0.9885
$\nu$ (df)	6.42

Very persistent; Half-life  $\approx 60$  days

Time Series Analysis and Forecasting

## GARCH vs EGARCH Comparison — S&P 500



### Leverage Effect Confirmed

EGARCH:  $\gamma = -0.12$  (significantly negative) — negative shocks amplify volatility more than positive shocks. Both models capture volatility clustering, but EGARCH better fits crisis periods (2008, 2020).

## Key Formulas

### Volatility Models

- ▣ **ARCH(q):**  $\sigma_t^2 = \omega + \sum_{i=1}^q \alpha_i \varepsilon_{t-i}^2$
- ▣ **GARCH(1,1):**  $\sigma_t^2 = \omega + \alpha \varepsilon_{t-1}^2 + \beta \sigma_{t-1}^2$
- ▣ **EGARCH:**  $\ln(\sigma_t^2) = \omega + \alpha(|z_{t-1}| - \mathbb{E}[|z|]) + \gamma z_{t-1} + \beta \ln(\sigma_{t-1}^2)$
- ▣ **GJR-GARCH:**  $\sigma_t^2 = \omega + \alpha \varepsilon_{t-1}^2 + \gamma \varepsilon_{t-1}^2 I_{t-1} + \beta \sigma_{t-1}^2$

### Properties and Measures

- ▣ **Unconditional variance:**  $\bar{\sigma}^2 = \frac{\omega}{1-\alpha-\beta}$      **Half-life:**  $HL = \frac{\ln(0.5)}{\ln(\alpha+\beta)}$
- ▣ **VaR:**  $\text{VaR}_\alpha = z_\alpha \cdot \sigma_{T+1}$      **Stationarity:**  $\alpha + \beta < 1$
- ▣ **ARCH-LM:**  $LM = T \cdot R^2 \sim \chi^2(q)$

## Summary — Chapter 5: Volatility Models

### Key Concepts

- ▣ **ARCH( $q$ )**: conditional variance depends on past squared errors
- ▣ **GARCH( $p,q$ )**: adds variance lags for persistence
- ▣ **EGARCH/GJR-GARCH**: capture leverage effect (asymmetric response)






### Applications

Risk measurement (VaR, ES), derivative pricing, portfolio management

### Practical Tip

Start with GARCH(1,1), check for leverage, choose distribution minimizing AIC/BIC!

## References

-  Engle, R.F. (1982). *Autoregressive Conditional Heteroscedasticity with Estimates of the Variance of United Kingdom Inflation*. *Econometrica*, 50(4), 987-1007.
-  Bollerslev, T. (1986). *Generalized Autoregressive Conditional Heteroskedasticity*. *Journal of Econometrics*, 31(3), 307-327.
-  Nelson, D.B. (1991). *Conditional Heteroskedasticity in Asset Returns: A New Approach*. *Econometrica*, 59(2), 347-370.
-  Glosten, L.R., Jagannathan, R., & Runkle, D.E. (1993). *On the Relation between the Expected Value and the Volatility of the Nominal Excess Return on Stocks*. *The Journal of Finance*, 48(5), 1779-1801.
-  Tsay, R.S. (2010). *Analysis of Financial Time Series*. 3rd Edition, Wiley.