



Time Series Analysis

Seminar 1: Stochastic processes and stationarity



Daniel Traian PELE

Bucharest University of Economic Studies

IDA Institute Digital Assets

Blockchain Research Center

AI4EFin Artificial Intelligence for Energy Finance

Romanian Academy, Institute for Economic Forecasting

MSCA Digital Finance

Acronyms used in this material

- ▣ ACF: Autocorrelation Function
- ▣ AI: Artificial Intelligence
- ▣ AR: AutoRegressive (model); in event studies: Abnormal Return
- ▣ BET: Bucharest Exchange Trading (index)
- ▣ BNR: Banca Națională a României (the National Bank of Romania)
- ▣ EODHD: EOD Historical Data (market-data provider)
- ▣ GDP: Gross Domestic Product
- ▣ HICP: Harmonised Index of Consumer Prices
- ▣ MA: Moving Average
- ▣ PACF: Partial AutoCorrelation Function

Today's question and route

- **Question:** is a series stationary, is it white noise, and how do the ACF and the Ljung–Box test answer these questions?
 - ▶ this seminar comes **before** Lecture 1: the section “What you need today” gives every definition the tasks use
- **Route**
 - ▶ Part A: autocovariances of moving averages and of a random walk, a sample ACF and a Ljung–Box statistic by hand
 - ▶ Part B: BET, S&P 500 and EUR/RON returns; Romanian GDP and inflation, each with an interpretation question
 - ▶ Part C: an open question for a project and an AI answer to audit
- Notebook for today: [open the seminar notebook in Google Colab](#); each task names its notebook section

Exercise map

Task	Question	Type	Model
A1, A2	mean, autocovariances and ACF of MA(1) and MA(2) processes	Solved, Proposed	A1
A3, A4	a random walk; a trend-stationary series and its first difference	Solved, Proposed	A3
A5, A6	sample ACF, bands, Box–Pierce and Ljung–Box by hand	Solved, Proposed	A5
B1, B2	prices and returns: BET; S&P 500 and EUR/RON	Solved, Proposed	B1
B3, B4	transformations: Romanian real GDP; Romanian HICP inflation	Solved, Proposed	B3
C1, C2	has inflation become more persistent? what is wrong in an AI answer?	Proposed	B1, B4

- **[Solved]**: full solution in the slides and in the notebook, a model to follow; **[Proposed]**: you solve it, following the model

Data used

Series	Source	Frequency	Period
BET, S&P 500	EODHD	daily, close	2000–2026
EUR/RON	BNR reference rate	daily	2005–2026
Real GDP, Romania	Eurostat (namq_10_gdp)	quarterly, not adjusted	1995–2026
HICP, Romania	Eurostat (prc_hicp_minr)	monthly, 2015 = 100	1996–2026

- Daily log returns in %: $r_t = 100 (\ln P_t - \ln P_{t-1})$; weekends and repeated holiday closes are dropped; the last day is 18 September 2026
- Growth and inflation rates in %: $100 \Delta \ln Y_t$ (one period), $100 \Delta_4 \ln Y_t$ and $100 \Delta_{12} \ln P_t$ (one year)
- In the notebook: `b1_returns('bet')`, `sample_acf(x)`, `ljung_box(x, 10)`; no account or key is needed

What you need today (1/4): processes and moments

- ▣ **Stochastic process** $\{X_t\}$: one random variable for each date t ; a time series $x_1, \dots, x_T$ is one observed path
 - ▶ mean $\mu_t = E[X_t]$; autocovariance $\gamma(t, s) = \text{Cov}(X_t, X_s)$
- ▣ **Weakly stationary**: $E[X_t^2] < \infty$, $E[X_t] = \mu$ and $\text{Cov}(X_t, X_{t+h}) = \gamma(h)$ for every t
 - ▶ **ACF**: $\rho(h) = \gamma(h)/\gamma(0)$, $\rho(0) = 1$, $|\rho(h)| \leq 1$, $\rho(-h) = \rho(h)$
- ▣ Rules for covariances (with constants a, b, c):
 - ▶ $\text{Cov}(aX + bY, Z) = a \text{Cov}(X, Z) + b \text{Cov}(Y, Z)$; $\text{Cov}(X + c, Y) = \text{Cov}(X, Y)$
 - ▶ $\text{Var}(\sum_i a_i X_i) = \sum_i \sum_j a_i a_j \text{Cov}(X_i, X_j)$

What you need today (2/4): white noise, moving averages, random walk

- ▣ **White noise** $\varepsilon_t \sim \text{WN}(0, \sigma^2)$: mean 0, variance σ^2 , $\text{Cov}(\varepsilon_t, \varepsilon_s) = 0$ for $t \neq s$
 - ▶ weak (uncorrelated), i.i.d. (independent), Gaussian (i.i.d. Normal)
- ▣ **MA(q)** (moving average): $X_t = \mu + \varepsilon_t + \theta_1 \varepsilon_{t-1} + \cdots + \theta_q \varepsilon_{t-q}$
 - ▶ with $\theta_0 = 1$: $\gamma(h) = \sigma^2 \sum_{j=0}^{q-h} \theta_j \theta_{j+h}$ for $0 \leq h \leq q$, and $\gamma(h) = 0$ for $h > q$
- ▣ **Random walk**: $X_t = X_{t-1} + \varepsilon_t$, $X_0 = 0$, so $X_t = \varepsilon_1 + \cdots + \varepsilon_t$; with drift: $X_t = c + X_{t-1} + \varepsilon_t$
 - ▶ **difference**: $\Delta X_t = X_t - X_{t-1}$; **seasonal difference**: $\Delta_s X_t = X_t - X_{t-s}$

What you need today (3/4): sample ACF and its bands

- Sample mean $\bar{x}$, **sample autocovariance** and **sample ACF** (divisor T):
 - ▶ $\hat{\gamma}(h) = \frac{1}{T} \sum_{t=1}^{T-h} (x_t - \bar{x})(x_{t+h} - \bar{x})$, $\hat{\rho}(h) = \hat{\gamma}(h)/\hat{\gamma}(0)$
- If the series is i.i.d.: $\hat{\rho}(h) \approx N(0, 1/T)$; **95% band** $\pm 1.96/\sqrt{T}$
 - ▶ a bar outside the band: $\rho(h) \neq 0$ at 5%, for that lag alone; with 20 lags, about 1 bar crosses by chance
- Typical patterns
 - ▶ white noise: no bar outside; MA(q): bars up to lag q , then nothing; AR(1): $\rho(h) = \phi^h$, geometric decay
 - ▶ random walk or trend: very slow, almost linear decay; seasonal data: peaks at lags $s, 2s, \dots$

What you need today (4/4): portmanteau tests and transformations

- $H_0: \rho(1) = \dots = \rho(m) = 0$; under H_0 both statistics are approximately $\chi^2(m)$
 - ▶ **Box–Pierce:** $Q(m) = T \sum_{h=1}^m \hat{\rho}(h)^2$; **Ljung–Box:** $Q^*(m) = T(T+2) \sum_{h=1}^m \hat{\rho}(h)^2 / (T-h)$
 - ▶ 5% critical values: $\chi_{0.95}^2(2) = 5.99$, $\chi_{0.95}^2(3) = 7.81$, $\chi_{0.95}^2(8) = 15.51$, $\chi_{0.95}^2(10) = 18.31$, $\chi_{0.95}^2(12) = 21.03$
- **Transformations** of a positive series Y_t
 - ▶ log: $\ln Y_t$ stabilises a variance that grows with the level; $100 \Delta \ln Y_t \approx$ growth in %
 - ▶ differences remove trends and random walks; Δ_4 and Δ_{12} remove the season of quarterly and monthly data
 - ▶ Box–Cox: $(Y_t^\lambda - 1)/\lambda$, with $\lambda = 0$ meaning $\ln Y_t$

A1: an MA(1) process [Solved]

Task

- ▣ $X_t = \varepsilon_t + 0.5 \varepsilon_{t-1}$, with $\varepsilon_t \sim \text{WN}(0, 2)$.
- ▣ 1. Compute the mean $E[X_t]$.
- ▣ 2. Compute $\gamma(0)$, $\gamma(1)$ and $\gamma(2)$.
- ▣ 3. Compute $\rho(1)$ and $\rho(2)$.
- ▣ 4. Say whether the process is weakly stationary, and why.
- ▣ Report: five numbers and one sentence.

Solution

- ▣ 1. $E[X_t] = 0 + 0.5 \cdot 0 = 0$
- ▣ 2. $\gamma(0) = \text{Var}(\varepsilon_t) + 0.25 \text{Var}(\varepsilon_{t-1}) = 2 + 0.5 = 2.50$
- ▣ $\gamma(1) = \text{Cov}(\varepsilon_t + 0.5\varepsilon_{t-1}, \varepsilon_{t+1} + 0.5\varepsilon_t) = 0.5 \cdot 2 = 1.00$;
 $\gamma(2) = 0$ (no common shock)
- ▣ 3. $\rho(1) = 1.00/2.50 = 0.40$; $\rho(2) = 0$
- ▣ 4. Yes: the mean and all $\gamma(h)$ are finite and do not depend on t .



A2: an MA(2) process [Proposed]

Solution discussed in class

Task

- ▣ $X_t = 2 + \varepsilon_t + 0.4 \varepsilon_{t-1} - 0.3 \varepsilon_{t-2}$, with $\varepsilon_t \sim \text{WN}(0, 1)$. Model: A1.
- ▣ 1. Compute the mean.
- ▣ 2. Compute $\gamma(0)$, $\gamma(1)$, $\gamma(2)$ and $\gamma(3)$.
- ▣ 3. Compute $\rho(1)$ and $\rho(2)$, and sketch the ACF up to lag 4.
- ▣ 4. Say how the constant 2 changes the autocovariances.
- ▣ Report: six numbers, the sketch and one sentence.



A3: a random walk [Solved]

Task

- ▣ $X_t = X_{t-1} + \varepsilon_t$, $X_0 = 0$, $\varepsilon_t \sim \text{WN}(0, 0.25)$.
- ▣ 1. Compute $E[X_{100}]$, $\text{Var}(X_{100})$ and $\text{Var}(X_{120})$.
- ▣ 2. Compute $\text{Cov}(X_{100}, X_{120})$ and $\text{Corr}(X_{100}, X_{120})$.
- ▣ 3. Say whether X_t and ΔX_t are stationary.
- ▣ 4. With a drift $c = 0.1$, compute $E[X_{100}]$.
- ▣ Report: six numbers and two sentences.

Solution

- ▣ 1. $X_t = \sum_{i=1}^t \varepsilon_i$: $E[X_{100}] = 0$; $\text{Var}(X_{100}) = 100 \cdot 0.25 = 25$; $\text{Var}(X_{120}) = 30$
- ▣ 2. The common shocks are $\varepsilon_1, \dots, \varepsilon_{100}$: $\text{Cov} = 25$; $\text{Corr} = 25 / \sqrt{25 \cdot 30} = \sqrt{100/120} = 0.913$
- ▣ 3. X_t : no, its variance grows with t ; $\Delta X_t = \varepsilon_t$: yes, it is white noise.
- ▣ 4. $E[X_{100}] = 100 \cdot 0.1 = 10$; the variance does not change.

A4: a trend-stationary series [Proposed]

Solution discussed in class

Task

- ▣ $Y_t = 5 + 0.2t + \varepsilon_t$, $\varepsilon_t \sim \text{WN}(0, 1)$. Model: A3.
- ▣ 1. Compute $E[Y_t]$ and $\text{Var}(Y_t)$, and say whether Y_t is stationary.
- ▣ 2. Write $D_t = \Delta Y_t$ and compute its mean, $\gamma(0)$, $\gamma(1)$ and $\rho(1)$.
- ▣ 3. Compare Y_t with a random walk with drift 0.2: what happens to a shock ε_{50} in each?
- ▣ Report: five numbers and two sentences.



A5: a sample ACF by hand [Solved]

Task

- ▣ Eight observations: $x = (4, 6, 5, 8, 7, 9, 6, 7)$.
- ▣ 1. Compute $\bar{x}$ and the deviations $x_t - \bar{x}$.
- ▣ 2. Compute $\hat{\gamma}(0)$, $\hat{\gamma}(1)$, $\hat{\gamma}(2)$ with divisor $T = 8$, then $\hat{\rho}(1)$ and $\hat{\rho}(2)$.
- ▣ 3. Compute the band $\pm 1.96/\sqrt{T}$ and compare.
- ▣ 4. Compute the Ljung–Box $Q^*(2)$ and decide at 5%.
- ▣ Report: seven numbers and one sentence.

Solution

- ▣ 1. $\bar{x} = 52/8 = 6.5$; deviations $(-2.5, -0.5, -1.5, 1.5, 0.5, 2.5, -0.5, 0.5)$
- ▣ 2. sum of squares 18: $\hat{\gamma}(0) = 2.250$; products at lag 1 sum to 0.25: $\hat{\gamma}(1) = 0.0312$; at lag 2 to 7.00: $\hat{\gamma}(2) = 0.875$
- ▣ $\hat{\rho}(1) = 0.014$, $\hat{\rho}(2) = 0.389$
- ▣ 3. Band ± 0.693 : both inside
- ▣ 4. $Q^*(2) = 8 \cdot 10 (\hat{\rho}(1)^2/7 + \hat{\rho}(2)^2/6) = 2.02 < 5.99$ ($p = 0.36$): do not reject white noise; with $T = 8$ the test has almost no power.

A6: Box–Pierce or Ljung–Box? [Proposed]

Solution discussed in class


Task

- ▣ A series of $T = 120$ monthly returns has $\hat{\rho}(1) = 0.21$, $\hat{\rho}(2) = -0.08$, $\hat{\rho}(3) = 0.12$. Model: A5.
- ▣ 1. Compute the band and say which autocorrelations lie outside it.
- ▣ 2. Compute $Q(3)$ and $Q^*(3)$.
- ▣ 3. Compare both with $\chi^2_{0.95}(3) = 7.81$ and decide.
- ▣ 4. Explain why the two tests can disagree, and which one you trust here.
- ▣ Report: four numbers and two sentences.

B1: BET prices and returns [Solved]

- **Question:** is the BET log price stationary, and are BET daily returns white noise?
- **Inputs:** BET closes since 2000; $\ln P_t$ and $r_t = 100 \Delta \ln P_t$
- **Tasks**
 1. Compute the sample ACF of $\ln P_t$ at lags 1, 10 and 50.
 2. Draw the ACF of r_t up to lag 20 with the band $\pm 1.96/\sqrt{T}$, and count the bars outside it.
 3. Compute Ljung–Box $Q^*(10)$ for r_t and for r_t^2 , with their p-values.
 4. Interpretation: can yesterday's return be used to forecast today's return?
- **Report:** three ACF values, the chart, two statistics with p-values and two sentences
- **Notebook:** section B1

B1: solution [Solved]



- $\ln P_t$: $\hat{\rho}(1) = 0.999$, $\hat{\rho}(10) = 0.991$, $\hat{\rho}(50) = 0.953$: almost no decay, a random-walk-like level, not stationary
- r_t ($T = 6670$, band ± 0.024): 7 of 20 bars outside, the largest $\hat{\rho}(1) = 0.109$; $Q^*(10) = 108.0$, $p < 0.001$
- r_t^2 : $\hat{\rho}(1) = 0.33$; $Q^*(10) = 2440$, $p < 0.001$: much stronger dependence
- Interpretation: yesterday's return explains only $\hat{\rho}(1)^2 = 1.2\%$ of today's variance, too little for a trading rule after costs; the size of yesterday's move does predict today's risk


B2: the S&P 500 and EUR/RON [Proposed]

- ▣ **Question:** do the S&P 500 and the EUR/RON rate behave like the BET?
- ▣ **Inputs:** S&P 500 closes since 2000; EUR/RON (BNR reference rate) since July 2005; model: B1
- ▣ **Tasks**
 1. Repeat the three steps of B1 for both series.
 2. Compare the sign and the size of $\hat{\rho}(1)$ of the returns across BET, S&P 500 and EUR/RON.
 3. Compare $Q^*(10)$ of the squared returns across the three series.
 4. Interpretation: which of the three return series is closest to white noise?
- ▣ **Report:** one table (three series, six numbers each) and two sentences
- ▣ **Notebook:** section B2

B3: Romanian real GDP [Solved]

- **Question:** which transformation makes Romanian quarterly GDP closest to stationary?
- **Inputs:** real GDP, chain-linked volumes (2010), not seasonally adjusted, Eurostat, from 1995
- **Tasks**
 1. Compute $\ln Y_t$, $100 \Delta \ln Y_t$ and $100 \Delta_4 \ln Y_t$.
 2. Draw the three series and their ACF up to lag 12.
 3. Report $\hat{\rho}(1)$, $\hat{\rho}(4)$ and $Q^*(8)$ for each.
 4. Interpretation: why is $\hat{\rho}(4)$ of the quarterly growth rate close to 1?
- **Report:** a table of nine numbers, the chart and two sentences
- **Notebook:** section B3

B3: solution [Solved]



	$\hat{\rho}(1)$	$\hat{\rho}(4)$	$Q^*(8)$
$\ln Y_t$	0.68	0.94	495.0
$100 \Delta \ln Y_t$	-0.24	0.96	329.6
$100 \Delta_4 \ln Y_t$	0.63	0.03	74.2

- $\ln Y_t$: trend and season, peaks at lags 4 and 8; $\Delta \ln Y_t$: the season dominates (standard deviation 27.5%); $\Delta_4 \ln Y_t$: mean 2.73%, the ACF dies out after 2–3 quarters
- Interpretation: a q/q rate of unadjusted data repeats the same seasonal swing every year, hence $\hat{\rho}(4) \approx 1$; the annual rate removes the season but reacts late to turning points and overlaps across quarters (positive $\hat{\rho}(1)$)

B4: Romanian inflation [Proposed]

- **Question:** is monthly Romanian inflation since 2005 white noise around its mean?
- **Inputs:** HICP, 2015 = 100, Eurostat, January 2005 to the last month available; model: B3
- **Tasks**
 1. Compute the monthly inflation $100 \Delta \ln P_t$ and the 12-month inflation $100 \Delta_{12} \ln P_t$.
 2. Draw the ACF of both up to lag 36 and report $\hat{\rho}(1)$, $\hat{\rho}(12)$ and $\hat{\rho}(24)$.
 3. Compute Ljung–Box $Q^*(12)$ for the monthly inflation.
 4. Interpretation: what does the ACF of the monthly inflation say about how fast an inflation shock fades?
- **Report:** six ACF values, one statistic with its p-value, the chart and two sentences
- **Notebook:** section B4

C1: has Romanian inflation become more persistent? [Proposed]

- ▣ **Question:** is monthly inflation more persistent after 2020 than in 2005–2019?
- ▣ **Inputs:** monthly HICP inflation of B4, split in January 2020; models: B1, B4
- ▣ **Tasks**
 1. Compute $\hat{\rho}(1)$ and $Q^*(12)$ in each subperiod, with the band of each.
 2. Draw $\hat{\rho}(1)$ on rolling windows of 60 months.
 3. List the events of 2020–2024 that could change the persistence.
 4. Interpretation: why is the difference of two sample autocorrelations hard to judge with 80 observations?
- ▣ **Report:** a table, the chart and a plan for a project
- ▣ **Notebook:** section C1

C2: audit an AI answer [Proposed]

- A student asked an AI assistant to interpret the S&P 500 daily data since 2000 ($T = 6\,714$ returns). The answer:
- (a) The ACF of the log price at lag 1 is 0.999, so the log price is a stationary process with a very long memory.
- (b) The return autocorrelation at lag 1 is -0.099, outside the band of 0.024, so a trader can forecast most of tomorrow's return.
- (c) Ljung-Box $Q(10)$ of the squared returns is 6133, so the returns themselves are autocorrelated.
- (d) A weakly stationary process is always strictly stationary, since its mean and variance are constant.
- (e) Differencing a series that is already stationary does no harm: white noise stays white noise.
- (f) A random walk has $\text{Var}(X(t)) = t * \sigma^2$, so it is not stationary.
- Tasks
 - ▶ 1. For each statement, say whether it is correct; if not, give the correct statement and, where possible, the correct number from the notebook (section C2).
 - ▶ 2. Report: a list of six verdicts with one line of justification each.

Key takeaways

- ▣ Autocovariances of linear processes follow from the covariance rules: only common shocks count
- ▣ A random walk has a growing variance; its difference is white noise; a trend-stationary series becomes an over-differenced MA(1)
- ▣ Look at the level, the changes and the squares: prices wander, returns are almost uncorrelated, squares are strongly correlated
- ▣ For macro data, choose the transformation first ($\log$, Δ , Δ_4 , Δ_{12}), then read the ACF
- ▣ An AI answer is a draft: check what is stationary, what “significant” means, and whether squares or levels were tested

After the seminar

- Lecture 1 develops each topic of today: stationarity, white noise and the random walk, Wold, ergodicity, ACF and PACF, the portmanteau tests, transformations
- Try the [Proposed] tasks in the notebook
- C1 can grow into a team project: inflation persistence in Romania and in the region (Hungary, Poland, Czechia)
- Reading: [Huang and Petukhina \(2022\)](#), Ch. 1–2; [Hyndman and Athanasopoulos \(2021\)](#), Ch. 2–3; [Brockwell and Davis \(2016\)](#), Ch. 1
- **The seminar is for practice and is not graded; the solutions of [Proposed] tasks are discussed in class**

References

- ▣ Box, G. E. P., & Pierce, D. A. (1970). Distribution of residual autocorrelations in autoregressive-integrated moving average time series models. *Journal of the American Statistical Association*, 65(332), 1509–1526. doi:[10.1080/01621459.1970.10481180](https://doi.org/10.1080/01621459.1970.10481180)
- ▣ Brockwell, P. J., & Davis, R. A. (2016). *Introduction to Time Series and Forecasting* (3rd ed.). Springer. doi:[10.1007/978-3-319-29854-2](https://doi.org/10.1007/978-3-319-29854-2)
- ▣ Huang, C., & Petukhina, A. (2022). *Applied Time Series Analysis and Forecasting with Python*. Springer. doi:[10.1007/978-3-031-13584-2](https://doi.org/10.1007/978-3-031-13584-2)
- ▣ Hyndman, R. J., & Athanasopoulos, G. (2021). *Forecasting: Principles and Practice* (3rd ed.). OTexts. otexts.com/fpp3
- ▣ Ljung, G. M., & Box, G. E. P. (1978). On a measure of lack of fit in time series models. *Biometrika*, 65(2), 297–303. doi:[10.1093/biomet/65.2.297](https://doi.org/10.1093/biomet/65.2.297)