

## Appendix: Examples of Equation Forecasts

The examples below show equation-continuation tasks sampled from recent arXiv papers. In each example, the red marker indicates the point where the equation was hidden from the forecasting model. Each display aligns the paper continuation and the model forecast at a shared point before the hidden suffix, so differences after the cut are easier to compare. For each example, the score line reports the clipped judge-score gain from conditioning on the forecast instead of a same-budget excerpt of immediately preceding paper context. Positive values mean the forecast made the true hidden suffix more likely under the judge.

### Example 1

Source: arXiv:2604.24491. Score gain over context baseline: +0.171.

$$\begin{aligned} \forall r \in \mathbb{N}^*, s \in \text{[CUT]} \frac{s_1 + 1}{2} + \mathbb{N}, \quad R_{r,s} = 0. \quad \text{paper} \\ \forall r \in \mathbb{N}^*, s \in \text{[CUT]} \mathbb{N}^*, \quad s \geq \frac{s_1 + 1}{2} \implies R_{r,s} = 0. \quad \text{forecast} \end{aligned}$$


---

### Example 2

Source: arXiv:2604.18952. Score gain over context baseline: +0.338.

$$\begin{aligned} |\ell s| \gtrsim |s \partial_p A_{k-p, \ell-k+p}| \geq |t \partial_p A_{k-p, p}| - \text{[CUT]} |\partial_p \Phi| \geq \theta_0 |kt| - |\partial_p \Phi| \geq \theta_0 |kt|/2, \quad \text{paper} \\ |\ell s| \gtrsim |s \partial_p A_{k-p, \ell-k+p}| \geq |t \partial_p A_{k-p, p}| - \text{[CUT]} |\partial_p \Phi| \geq \frac{\theta_0}{2} |kt|. \quad \text{forecast} \end{aligned}$$


---

### Example 3

Source: arXiv:2604.19881. Score gain over context baseline: +0.136.

$$\begin{aligned} Y_{ab}^\mu Y_\mu^{ab} &= \sum_{i=2,3} (\partial_a \partial_b dx_i - \frac{1}{2} d_{ab} \partial_c \partial^c dx_i)^2 \\ \text{paper} \quad \frac{1}{2} \sum_{i=\text{[CUT]} 2,3} &\left( (\partial_{x_0}^2 dx_i)^2 + (\partial_{x_1}^2 dx_i)^2 + 2 \partial_{x_1}^2 dx_i \cdot \partial_{x_0}^2 dx_i \right) = -\frac{1}{2} J(x_0, x_1), \\ \text{forecast} \quad \frac{1}{2} \sum_{i=\text{[CUT]} 2,3} &\left( (\partial_{x_0}^2 dx_i - \partial_{x_1}^2 dx_i)^2 + 4(\partial_{x_0} \partial_{x_1} dx_i)^2 \right). \end{aligned}$$


---

### Example 4

Source: arXiv:2604.20635. Score gain over context baseline: +0.345.

$$\begin{aligned} E = \frac{\partial l}{\partial u} \cdot u - l, \quad p = \text{[CUT]} l - \rho \frac{\partial l}{\partial \rho}. \quad \text{paper} \\ E = \frac{\partial l}{\partial u} \cdot u - l, \quad p = \text{[CUT]} l - \rho \frac{\partial l}{\partial \rho} \quad \text{forecast} \end{aligned}$$


---

## Example 5

Source: arXiv:2604.19506. Score gain over context baseline: +0.090.

$$N^{(2)}(z) = \begin{cases} \mathcal{E}(z)N^{alg}(z), & z \in \text{[CUT]} \mathbb{C} \setminus \mathcal{D}, \quad \mathcal{E}(z)N^{alg}(z)N^{\tilde{\kappa}}(z), z \in \mathcal{D}, \quad \text{paper} \\ \mathcal{E}(z)N^{alg}(z), & z \in \text{[CUT]} \mathbb{C} \setminus \mathcal{D}, \quad \mathcal{E}(z)N^{loc}(z), z \in \mathcal{D}. \quad \text{forecast} \end{cases}$$


---

## Example 6

Source: arXiv:2604.26927. Score gain over context baseline: +0.684.

$$\Omega^{cl}(d=2, N, k) = \text{[CUT]} \frac{1}{N} \frac{k!}{k^k} \sum_{l=0}^k \frac{k^l}{l!}. \quad \text{paper}$$

$$\Omega^{cl}(d=2, N, k) = \text{[CUT]} \frac{1}{N} \frac{k!}{k^k} \sum_{l=0}^k \frac{k^l}{l!}. \quad \text{forecast}$$


---

## Example 7

Source: arXiv:2604.27074. Score gain over context baseline: -0.076.

$$K_T(x_f, x_i) = \sum_{X: x_i \rightarrow x_f} \mathcal{A}[X] = \text{[CUT]} \sum_{X: x_i \rightarrow x_f} \sqrt{\mathbb{P}[X]} e^{i\Phi[X]}. \quad \text{paper}$$

$$K_T(x_f, x_i) = \sum_{X: x_i \rightarrow x_f} \mathcal{A}[X] = \text{[CUT]} \sum_{X: x_i \rightarrow x_f} \mathcal{A}_0[X] \mathcal{Z}_T[X], \quad \text{forecast}$$


---

## Example 8

Source: arXiv:2604.27015. Score gain over context baseline: -0.026.

$$\begin{aligned} f_0(r) &= r \bmod 2, \\ f_k(r) &= \text{[CUT]} \left\lfloor \frac{r}{2^k} \right\rfloor \bmod 2 \quad (k = 1, \dots, K), \quad \text{paper} \\ f_0(r) &= r \bmod 2, \\ f_k(r) &= \text{[CUT]} \left\lfloor \frac{r}{\Delta_k} \right\rfloor \bmod 2, \quad k = 1, \dots, K. \quad \text{forecast} \end{aligned}$$


---

## Example 9

Source: arXiv:2604.24755. Score gain over context baseline: -0.136.

$$H_{\text{child}}^{\text{eff}} = - \sum_{(E, \mu)} \left( \sum_{P_d} Q_{P_d} - \text{[CUT]} \sum_{P_o} Q_{P_o}^1 \right) - \sum_C F_C = H_{\text{XC}}. \quad \text{paper}$$

$$H_{\text{child}}^{\text{eff}} = - \sum_{(E, \mu)} \left( \sum_{P_d} Q_{P_d} - \text{[CUT]} \sum_{V_d} \frac{1}{2} \left( B_{V_d}^{1\bar{1}} + B_{V_d}^{\psi\bar{\psi}} \right) \right) \quad \text{forecast}$$


---

## Example 10

Source: arXiv:2604.24314. Score gain over context baseline: +0.051.

$$\begin{aligned} v_k(z) &:= \frac{(\eta u_k)(\tilde{z}_k + r_k z) - (\eta u_k)(\tilde{z}_k)}{r_k^\alpha E_k}, \\ \text{paper} \quad w_k(z) &:= \text{[CUT]} \frac{\eta(\tilde{z}_k)(u_k(\tilde{z}_k + r_k z) - u_k(\tilde{z}_k))}{r_k^\alpha E_k}, \\ \text{forecast} \quad w_k(z) &:= \text{[CUT]} \frac{u_k(\tilde{z}_k + r_k z)}{r_k^\alpha E_k}. \end{aligned}$$

---