

# AN EXPLICIT EXAMPLE OF TOMTER'S ALGEBRAIC ANOSOV FLOWS OF THE MIXED TYPE

DANYU ZHANG

We give an example of Tomter's algebraic Anosov flow of mixed type over a geodesic flow of a genus 2 surface, and try to understand the orbits and foliations in the fibres.

## 1 Arithmetic Fuchsian groups

We know that  $\mathcal{A} := (\frac{26, -5}{\mathbb{Q}})$  is a quaternion division algebra that gives rise to an arithmetic Fuchsian group of a genus 2 surface [Mac08]. We compute the action on the torus  $\mathbb{R}^4$ . Then the action of all the integer points passes to an action on  $\mathbb{T}^4$ .

More generally, for a quaternion algebra  $\mathcal{B} = (\frac{a, b}{\mathbb{F}})$  where  $\mathbb{F}$  is a field extension of degree  $n$  over  $\mathbb{Q}$ ,  $\mathcal{B}$  acts on  $\mathbb{R}^{4n}$ .

We can write  $x \in \mathcal{A}$  as  $x = a + bi + cj + dk$  where  $i^2 = 26$  and  $j^2 = -5$ ,  $a, b, c, d \in \mathbb{Q}$ . We also have  $ij = k = -ji, jk = -kj = jij = -ij^2 = 5i, ik = -ki = iij = 26j, k^2 = ijij = -jii = -26j^2 = 130$ .

There is an embedding of  $x = a + bi + cj + dk$  where  $a, b, c, d \in \mathbb{R}$  into  $M_{22}(\mathbb{R})$  by

$$1 \mapsto \begin{bmatrix} 1 & \\ & 1 \end{bmatrix}, i \mapsto \begin{bmatrix} \sqrt{26} & \\ & -\sqrt{26} \end{bmatrix}, j \mapsto \begin{bmatrix} & 1 \\ -5 & \end{bmatrix}, k \mapsto \begin{bmatrix} & \sqrt{26} \\ 5\sqrt{26} & \end{bmatrix}.$$

Then  $x \mapsto \begin{bmatrix} a + b\sqrt{26} & c + d\sqrt{26} \\ -5c + d(5\sqrt{26}) & a - b\sqrt{26} \end{bmatrix}$ . Note that  $|x|^2 = x\bar{x} = (a + bi + cj + dk)(a - bi - cj - dk) = a^2 - 26b^2 + 5c^2 - 130d^2 = \det \left( \begin{bmatrix} a + b\sqrt{26} & c + d\sqrt{26} \\ -5c + d(5\sqrt{26}) & a - b\sqrt{26} \end{bmatrix} \right)$ .

We can also embed  $\mathcal{A}$  into  $M_{44}(\mathbb{Q})$  by  $x \mapsto (M_x : y \mapsto xy)$  for  $x, y \in \mathcal{A}$ . Now  $M_x$  is 4-by-4 matrix with rational coefficients which we can write down explicitly:

$$\begin{bmatrix} a & 26b & -5c & 130d \\ b & a & 5d & 5c \\ c & -26d & a & 26b \\ d & -c & b & a \end{bmatrix}.$$

The image of  $\mathcal{A}$  forms a linear subspace of  $M_{44}(\mathbb{Q})$ .

On the other hand, we consider  $\begin{bmatrix} a + b\sqrt{26} & c + d\sqrt{26} \\ -5c + d(5\sqrt{26}) & a - b\sqrt{26} \end{bmatrix}$  acting on  $\left\{ \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 0 \\ 1 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix} \right\}$  (a change of basis of  $1, i, j, k \in \mathcal{A}$  in  $M_{22}(\mathbb{R})$ ) as acting on  $\mathbb{R}^2 \oplus \mathbb{R}^2 = \mathbb{R}^4$ , so the  $\det(M_x) = \det \left( \begin{bmatrix} a + b\sqrt{26} & c + d\sqrt{26} \\ -5c + d(5\sqrt{26}) & a - b\sqrt{26} \end{bmatrix} \right)^2$ .

Now denote  $\mathbb{G}(F) := \{M_x \in M_{44}(F) \cap W : \det(M_x) = 1\}$ , where  $F$  can be  $\mathbb{Z}, \mathbb{Q}, \mathbb{R}$ . We check that  $\mathbb{G}(\mathbb{R}) \cong SL(2, \mathbb{R})$ : the isomorphism is given by  $(x \mapsto M_x)$ . Denote  $\Gamma := \mathbb{G}(\mathbb{Z})/\{\pm I_2\}$ .  $\Gamma$  is a Fuchsian group and it is cocompact in  $PSL(2, \mathbb{R})$  by Mahler compactness criterion [Kat92, Theorem 5.4.1]. The action of  $\Gamma = (\mathbb{Z})/\{\pm I_2\}$  on the torus is well-defined by taking  $\pm x \mapsto \pm M_{\pm x}$ .

## 2 Tomter's example

We compute the eigenvalues and eigenspaces of such matrices  $M_x$ . Let also  $a^2 - 26b^2 + 5c^2 - 130d^2 = 1$ .

For any  $\gamma \in \Gamma$ , there exists  $g \in PSL(2, \mathbb{R})$  such that  $g\gamma g^{-1}$  is a translation along the axis  $(i, i)$  of the upper half plane model  $\mathbb{H}^2$ , which corresponds to  $\begin{bmatrix} a + b\sqrt{26} & \\ & a - b\sqrt{26} \end{bmatrix}$  (in this case  $c = d = 0$ ). The eigenvalues of  $M_{g\gamma g^{-1}}$  are  $\lambda_1 = a + \sqrt{26}b$ ,  $\lambda_2 = a - \sqrt{26}b$ , and the eigenvectors are  $\begin{bmatrix} \sqrt{26} \\ 1 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 0 \\ \sqrt{26} \\ 1 \end{bmatrix}, \begin{bmatrix} -\sqrt{26} \\ 1 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 0 \\ -\sqrt{26} \\ 1 \end{bmatrix}$ . We have  $\lambda_1 \lambda_2 = 1$ , and  $M_{g\gamma g^{-1}}$  is hyperbolic. Therefore  $\gamma$  acts on the torus also by a hyperbolic matrix.

For any  $(\gamma, n) \in \Gamma \ltimes \mathbb{Z}^4$ , we have  $(g, x)(\gamma, n) = (g\gamma, \gamma^{-1}.x + n)$ . Denote  $M := PSL(2, \mathbb{R}) \ltimes \mathbb{R}^4 / (\Gamma \ltimes \mathbb{Z}^4)$ .  $M$  is a torus bundle over the unit tangent bundle of a surface of genus 2.

Let  $a_t = \begin{bmatrix} e^{-t/2} & \\ & e^{t/2} \end{bmatrix}$  denote the geodesic flow on  $T^1\mathbb{H}^2 \cong PSL(2, \mathbb{R})$  and for any  $g \in PSL(2, \mathbb{R})$  it acts by the right action  $R_{a_t}(g) = ga_t^{-1}$ . For  $(g, x) \in PSL(2, \mathbb{R}) \ltimes \mathbb{R}^4$ , let  $R_{a_t}(g, x) = (ga_t^{-1}, x)$ .

The flow  $\Phi^t : M \rightarrow M$  defined by  $\Phi^t((g, x)(\Gamma \ltimes \mathbb{Z}^4)) = (ga_t^{-1}, x)(\Gamma \ltimes \mathbb{Z}^4)$  is Anosov.

## 3 Orbits of $\Phi^t$

## References

- [Kat92] Svetlana Katok. *Fuchsian groups*. Chicago: The University of Chicago Press, 1992.
- [Mac08] Melissa L. Macasieb. Derived Arithmetic Fuchsian Groups of Genus Two. *Experimental Mathematics*, 17(3):347 – 369, 2008.
- [Tom70] Per Tomter. Anosov flows on infra-homogeneous spaces. *Global Analysis*, 14:299–327, 1970.