

Computing the limit of the Nonlocal Curvature of Curves in \mathbb{R}^n

Dan Zimmerman

AMS 2024 Fall Eastern Sectional Meeting, Oct 19, 2024



Origins of Nonlocal Curvature of Curves

Basic functional:

$$J(u) = \int_{\mathbb{R}^n} |\nabla u|^2 dV$$

Origins of Nonlocal Curvature of Curves

Basic functional:

$$J(u) = \int_{\mathbb{R}^n} |\nabla u|^2 dV$$

Replace ∇u with its fractional counterpart, with $0 < \sigma < 1$

$$\nabla^\sigma u(x) \simeq \int_{\mathbb{R}^n} \frac{(u(y) - u(x))(y - x)}{|y - x|^{n+\sigma+1}} dy$$

Origins of Nonlocal Curvature of Curves

Basic functional:

$$J(u) = \int_{\mathbb{R}^n} |\nabla u|^2 dV$$

Replace ∇u with its fractional counterpart, with $0 < \sigma < 1$

$$\nabla^\sigma u(x) \simeq \int_{\mathbb{R}^n} \frac{(u(y) - u(x))(y - x)}{|y - x|^{n+\sigma+1}} dy$$

to obtain

$$J_\sigma(u) = \int_{\mathbb{R}^n} |\nabla^\sigma u|^2 dV \simeq \int_{\mathbb{R}^n} \int_{\mathbb{R}^n} \frac{|u(x) - u(y)|^2}{|x - y|^{n+\sigma}} dx dy$$

Origins of Nonlocal Curvature of Curves

Basic functional:

$$J(u) = \int_{\mathbb{R}^n} |\nabla u|^2 dV$$

Replace ∇u with its fractional counterpart, with $0 < \sigma < 1$

$$\nabla^\sigma u(x) \simeq \int_{\mathbb{R}^n} \frac{(u(y) - u(x))(y - x)}{|y - x|^{n+\sigma+1}} dy$$

to obtain

$$J_\sigma(u) = \int_{\mathbb{R}^n} |\nabla^\sigma u|^2 dV \simeq \int_{\mathbb{R}^n} \int_{\mathbb{R}^n} \frac{|u(x) - u(y)|^2}{|x - y|^{n+\sigma}} dx dy$$

Consider $u = \chi_E$ for bounded E : $\int_E \int_{E^c} \frac{1}{|x - y|^{n+\sigma}} dx dy$

$$\text{Per}_\sigma(E) := \frac{1}{\alpha_{n-1}} \int_E \int_{E^c} \frac{1}{|x - y|^{n+\sigma}} dx dy$$

For unbounded E , fix Ω bounded, and set

$$\text{Per}_\sigma(E, \Omega) := \frac{1}{\alpha_{n-1}} \left(\int_{E \cap \Omega} \int_{E^c} + \int_{E \cap \Omega^c} \int_{E^c \cap \Omega} \right) \frac{1}{|x - y|^{n+\sigma}} dx dy$$

For unbounded E , fix Ω bounded, and set

$$\begin{aligned} \text{Per}_\sigma(E, \Omega) &:= \frac{1}{\alpha_{n-1}} \left(\int_{E \cap \Omega} \int_{E^c} + \int_{E \cap \Omega^c} \int_{E^c \cap \Omega} \right) \frac{1}{|x - y|^{n+\sigma}} dx dy \\ &= \frac{1}{\alpha_{n-1}} \int_E \int_{E^c} \frac{\max\{\chi_\Omega(x), \chi_\Omega(y)\}}{|x - y|^{n+\sigma}} dx dy \end{aligned}$$

For unbounded E , fix Ω bounded, and set

$$\begin{aligned} \text{Per}_\sigma(E, \Omega) &:= \frac{1}{\alpha_{n-1}} \left(\int_{E \cap \Omega} \int_{E^c} + \int_{E \cap \Omega^c} \int_{E^c \cap \Omega} \right) \frac{1}{|x - y|^{n+\sigma}} dx dy \\ &= \boxed{\frac{1}{\alpha_{n-1}} \int_E \int_{E^c} \frac{\max\{\chi_\Omega(x), \chi_\Omega(y)\}}{|x - y|^{n+\sigma}} dx dy} \end{aligned}$$

For unbounded E , fix Ω bounded, and set

$$\begin{aligned} \text{Per}_\sigma(E, \Omega) &:= \frac{1}{\alpha_{n-1}} \left(\int_{E \cap \Omega} \int_{E^c} + \int_{E \cap \Omega^c} \int_{E^c \cap \Omega} \right) \frac{1}{|x - y|^{n+\sigma}} dx dy \\ &= \boxed{\frac{1}{\alpha_{n-1}} \int_E \int_{E^c} \frac{\max\{\chi_\Omega(x), \chi_\Omega(y)\}}{|x - y|^{n+\sigma}} dx dy} \end{aligned}$$

E minimizer of when $\text{Per}_\sigma(E, \Omega) \leq \text{Per}_\sigma(F, \Omega)$ s.t. $F \setminus \Omega = E \setminus \Omega$,

For unbounded E , fix Ω bounded, and set

$$\begin{aligned} \text{Per}_\sigma(E, \Omega) &:= \frac{1}{\alpha_{n-1}} \left(\int_{E \cap \Omega} \int_{E^c} + \int_{E \cap \Omega^c} \int_{E^c \cap \Omega} \right) \frac{1}{|x - y|^{n+\sigma}} dx dy \\ &= \boxed{\frac{1}{\alpha_{n-1}} \int_E \int_{E^c} \frac{\max\{\chi_\Omega(x), \chi_\Omega(y)\}}{|x - y|^{n+\sigma}} dx dy} \end{aligned}$$

E minimizer of when $\text{Per}_\sigma(E, \Omega) \leq \text{Per}_\sigma(F, \Omega)$ s.t. $F \setminus \Omega = E \setminus \Omega$,

$$\int_{\mathbb{R}^n} \frac{\chi_E(x) - \chi_{E^c}(x)}{|z - x|^{n+\sigma}} dx = 0 \quad z \in \partial E$$

For unbounded E , fix Ω bounded, and set

$$\begin{aligned} \text{Per}_\sigma(E, \Omega) &:= \frac{1}{\alpha_{n-1}} \left(\int_{E \cap \Omega} \int_{E^c} + \int_{E \cap \Omega^c} \int_{E^c \cap \Omega} \right) \frac{1}{|x - y|^{n+\sigma}} dx dy \\ &= \boxed{\frac{1}{\alpha_{n-1}} \int_E \int_{E^c} \frac{\max\{\chi_\Omega(x), \chi_\Omega(y)\}}{|x - y|^{n+\sigma}} dx dy} \end{aligned}$$

E minimizer of when $\text{Per}_\sigma(E, \Omega) \leq \text{Per}_\sigma(F, \Omega)$ s.t. $F \setminus \Omega = E \setminus \Omega$,

$$\int_{\mathbb{R}^n} \frac{\chi_E(x) - \chi_{E^c}(x)}{|z - x|^{n+\sigma}} dx = 0 \quad z \in \partial E$$


Nonlocal mean curvature at $z \in \partial E$

$$H_\sigma(z) := \frac{1}{\omega_{n-1}} \int_{\mathbb{R}^n} \frac{\chi_E(x) - \chi_{E^c}(x)}{|z - x|^{n+\sigma}} dx$$

$$\text{Per}_\sigma(E, \Omega) := \frac{1}{\alpha_{n-1}} \int_E \int_{E^c} \frac{\max\{\chi_\Omega(x), \chi_\Omega(y)\}}{|x - y|^{n+\sigma}} dx dy$$


$$\begin{aligned}\mathrm{Per}_\sigma(E, \Omega) &:= \frac{1}{\alpha_{n-1}} \int_E \int_{E^c} \frac{\max\{\chi_\Omega(x), \chi_\Omega(y)\}}{|x-y|^{n+\sigma}} dx dy \\ &= \frac{1}{2\alpha_{n-1}} \int_{\mathcal{X}(\partial E)} \frac{\max\{\chi_\Omega(x), \chi_\Omega(y)\}}{|x-y|^{n+\sigma}} dx dy\end{aligned}$$

$$\begin{aligned}\text{Per}_\sigma(E, \Omega) &:= \frac{1}{\alpha_{n-1}} \int_E \int_{E^c} \frac{\max\{\chi_\Omega(x), \chi_\Omega(y)\}}{|x-y|^{n+\sigma}} dx dy \\ &= \frac{1}{2\alpha_{n-1}} \int_{\mathcal{X}(\partial E)} \frac{\max\{\chi_\Omega(x), \chi_\Omega(y)\}}{|x-y|^{n+\sigma}} dx dy\end{aligned}$$



$$\mathcal{X}(\partial E) = \{(x, y) \in \mathbb{R}^n \times \mathbb{R}^n \mid \mathcal{H}^0([x, y] \cap \partial E) \text{ is odd}\}$$

$$\begin{aligned}\text{Per}_\sigma(E, \Omega) &:= \frac{1}{\alpha_{n-1}} \int_E \int_{E^c} \frac{\max\{\chi_\Omega(x), \chi_\Omega(y)\}}{|x-y|^{n+\sigma}} dx dy \\ &= \frac{1}{2\alpha_{n-1}} \int_{\mathcal{X}(\partial E)} \frac{\max\{\chi_\Omega(x), \chi_\Omega(y)\}}{|x-y|^{n+\sigma}} dx dy\end{aligned}$$



$$\mathcal{X}(\partial E) = \{(x, y) \in \mathbb{R}^n \times \mathbb{R}^n \mid \mathcal{H}^0([x, y] \cap \partial E) \text{ is odd}\}$$


$$\text{Area}_\sigma(S, \Omega) := \frac{1}{2\alpha_{n-1}} \int_{\mathcal{X}(S)} \frac{\max\{\chi_\Omega(x), \chi_\Omega(y)\}}{|x-y|^{n+\sigma}} dx dy$$

$$\text{Area}_\sigma^{(n=2)}(\mathcal{S}, \Omega) = \frac{1}{4} \int_{\mathcal{X}(\mathcal{S})} \frac{\max\{\chi_\Omega(x), \chi_\Omega(y)\}}{|x - y|^{2+\sigma}} dx dy$$

A fractional notion of length and an associated nonlocal curvature - Seguin (2020)

$$\text{Area}_\sigma^{(n=2)}(\mathcal{S}, \Omega) = \frac{1}{4} \int_{\mathcal{X}(\mathcal{S})} \frac{\max\{\chi_\Omega(x), \chi_\Omega(y)\}}{|x - y|^{2+\sigma}} dx dy$$


Disks! (intersecting \mathcal{S} odd times)



A fractional notion of length and an associated nonlocal curvature - Seguin (2020)

$$\text{Area}_\sigma^{(n=2)}(\mathcal{S}, \Omega) = \frac{1}{4} \int_{\mathcal{X}(\mathcal{S})} \frac{\max\{\chi_\Omega(x), \chi_\Omega(y)\}}{|x - y|^{2+\sigma}} dx dy$$

Disks! (intersecting \mathcal{S} odd times)




$$= \frac{1}{2} \int_{\mathcal{D}(\mathcal{S})} \frac{\max\{\chi_\Omega(p - ru^\perp), \chi_\Omega(p + ru^\perp)\}}{(2r)^{1+\sigma}} d\mathcal{H}^4(p, u, r)$$

A fractional notion of length and an associated nonlocal curvature - Seguin (2020)

$$\text{Area}_\sigma^{(n=2)}(\mathcal{S}, \Omega) = \frac{1}{4} \int_{\mathcal{X}(\mathcal{S})} \frac{\max\{\chi_\Omega(x), \chi_\Omega(y)\}}{|x - y|^{2+\sigma}} dx dy$$

Disks! (intersecting \mathcal{S} odd times)



$$= \frac{1}{2} \int_{\mathcal{D}(\mathcal{S})} \frac{\max\{\chi_\Omega(p - ru^\perp), \chi_\Omega(p + ru^\perp)\}}{(2r)^{1+\sigma}} d\mathcal{H}^4(p, u, r)$$

Generalizes nicely


$$\text{Len}_\sigma(\mathcal{C}, \Omega) := \frac{\Gamma\left(\frac{n+1}{2}\right)^2}{2\pi^{n-1}} \int_{\mathcal{D}(\mathcal{C})} \frac{\sup_{v \in \mathcal{U}_n \cap u^\perp} \chi_\Omega(p + rv)}{r^{n-1+\sigma}} d\mathcal{H}^{2n}(p, u, r)$$

Minimizing over \mathcal{C} with fixed boundary, $z \in \mathcal{C}$

$$\left(\int_{\mathcal{A}_{\text{even}}^+(z)} - \int_{\mathcal{A}_{\text{odd}}^+(z)} \right) \frac{(a \cdot t(z))b - (b \cdot t(z))a}{r^{1+\sigma}} d\mathcal{H}^{2n-2}(a, b, r) = 0$$


Minimizing over \mathcal{C} with fixed boundary, $z \in \mathcal{C}$

$$\left(\int_{\mathcal{A}_{\text{even}}^+(z)} - \int_{\mathcal{A}_{\text{odd}}^+(z)} \right) \frac{(a \cdot t(z))b - (b \cdot t(z))a}{r^{1+\sigma}} d\mathcal{H}^{2n-2}(a, b, r) = 0$$



Minimizing over \mathcal{C} with fixed boundary, $z \in \mathcal{C}$

$$\left(\int_{\mathcal{A}_{\text{even}}^+(z)} - \int_{\mathcal{A}_{\text{odd}}^+(z)} \right) \frac{(a \cdot t(z))b - (b \cdot t(z))a}{r^{1+\sigma}} d\mathcal{H}^{2n-2}(a, b, r) = 0$$



Nonlocal vector curvature

$$\kappa_{\sigma}(z) := \left(\int_{\mathcal{A}_{\text{odd}}^+(z)} - \int_{\mathcal{A}_{\text{even}}^+(z)} \right) \frac{(a \cdot t(z))b - (b \cdot t(z))a}{r^{1+\sigma}} d\mathcal{H}^{2n-2}(a, b, r)$$

Asymptotics of fractional & Nonlocal Quantities

$$\lim_{\sigma \uparrow 1} (1 - \sigma) \operatorname{Per}_\sigma(E, B_r) = \operatorname{Per}(E, B_r)$$

$$\lim_{\sigma \uparrow 1} (1 - \sigma) H_\sigma(z) = H(z)$$

$$\lim_{\sigma \uparrow 1} (1 - \sigma) \operatorname{Per}_\sigma(E, B_r) = \operatorname{Per}(E, B_r)$$

$$\lim_{\sigma \uparrow 1} (1 - \sigma) H_\sigma(z) = H(z)$$

$$\lim_{\sigma \uparrow 1} (1 - \sigma) \operatorname{Len}_\sigma(\mathcal{C}, \Omega) = \operatorname{Len}(\mathcal{C}, \Omega)$$

$$\lim_{\sigma \uparrow 1} (1 - \sigma) \operatorname{Per}_\sigma(E, B_r) = \operatorname{Per}(E, B_r)$$

$$\lim_{\sigma \uparrow 1} (1 - \sigma) H_\sigma(z) = H(z)$$

$$\lim_{\sigma \uparrow 1} (1 - \sigma) \operatorname{Len}_\sigma(\mathcal{C}, \Omega) = \operatorname{Len}(\mathcal{C}, \Omega)$$

$$\lim_{\sigma \uparrow 1} (1 - \sigma) \kappa_\sigma(z) \stackrel{?}{=} \kappa(z)$$

$$\lim_{\sigma \uparrow 1} (1 - \sigma) \operatorname{Per}_\sigma(E, B_r) = \operatorname{Per}(E, B_r)$$

$$\lim_{\sigma \uparrow 1} (1 - \sigma) H_\sigma(z) = H(z)$$

$$\lim_{\sigma \uparrow 1} (1 - \sigma) \operatorname{Len}_\sigma(\mathcal{C}, \Omega) = \operatorname{Len}(\mathcal{C}, \Omega)$$

$$\lim_{\sigma \uparrow 1} (1 - \sigma) \, ? \, \kappa_\sigma(z) \stackrel{?}{=} \kappa(z)$$

Argument Outline

1. For \mathcal{C}_R circle with radius R , in \mathbb{R}^2

$$\lim_{\sigma \uparrow 1} (1 - \sigma) \quad \kappa_\sigma(z) = \kappa(z) \quad z \in \mathcal{C}_R$$

Argument Outline

1. For \mathcal{C}_R circle with radius R , in \mathbb{R}^2

$$\lim_{\sigma \uparrow 1} (1 - \sigma) \frac{1}{K_2} \kappa_\sigma(z) = \kappa(z) \quad z \in \mathcal{C}_R, K_2 \in \mathbb{R}$$

Argument Outline

1. For \mathcal{C}_R circle with radius R , in \mathbb{R}^2

$$\lim_{\sigma \uparrow 1} (1 - \sigma) \frac{1}{K_2} \kappa_\sigma(z) = \kappa(z) \quad z \in \mathcal{C}_R, K_2 \in \mathbb{R}$$

2. Use (1), same \mathcal{C}_R , in \mathbb{R}^n

$$\lim_{\sigma \uparrow 1} (1 - \sigma) \kappa_\sigma(z) = \kappa(z) \quad z \in \mathcal{C}_R$$

Argument Outline

1. For \mathcal{C}_R circle with radius R , in \mathbb{R}^2

$$\lim_{\sigma \uparrow 1} (1 - \sigma) \frac{1}{K_2} \kappa_\sigma(z) = \kappa(z) \quad z \in \mathcal{C}_R, K_2 \in \mathbb{R}$$

2. Use (1), same \mathcal{C}_R , in \mathbb{R}^n

$$\lim_{\sigma \uparrow 1} (1 - \sigma) \frac{1}{K(n)} \kappa_\sigma(z) = \kappa(z) \quad z \in \mathcal{C}_R, K(n) \in \mathbb{R} \text{ s.t. } K(2) = K_2$$

Argument Outline

1. For \mathcal{C}_R circle with radius R , in \mathbb{R}^2

$$\lim_{\sigma \uparrow 1} (1 - \sigma) \frac{1}{K_2} \kappa_\sigma(z) = \kappa(z) \quad z \in \mathcal{C}_R, K_2 \in \mathbb{R}$$

2. Use (1), same \mathcal{C}_R , in \mathbb{R}^n

$$\lim_{\sigma \uparrow 1} (1 - \sigma) \frac{1}{K(n)} \kappa_\sigma(z) = \kappa(z) \quad z \in \mathcal{C}_R, K(n) \in \mathbb{R} \text{ s.t. } K(2) = K_2$$

3. For arbitrary \mathcal{C} and \mathcal{C}_R ($R = 1/\kappa(z; \mathcal{C})$) at $z \in \mathcal{C}$

$$\lim_{\sigma \uparrow 1} (1 - \sigma) |\kappa_\sigma(z; \mathcal{C}) - \kappa_\sigma(z; \mathcal{C}_R)| = 0$$

Argument Outline

1. For \mathcal{C}_R circle with radius R , in \mathbb{R}^2

$$\lim_{\sigma \uparrow 1} (1 - \sigma) \frac{1}{K_2} \kappa_\sigma(z) = \kappa(z) \quad z \in \mathcal{C}_R, K_2 \in \mathbb{R}$$

2. Use (1), same \mathcal{C}_R , in \mathbb{R}^n

$$\lim_{\sigma \uparrow 1} (1 - \sigma) \frac{1}{K(n)} \kappa_\sigma(z) = \kappa(z) \quad z \in \mathcal{C}_R, K(n) \in \mathbb{R} \text{ s.t. } K(2) = K_2$$

3. For arbitrary \mathcal{C} and \mathcal{C}_R ($R = 1/\kappa(z; \mathcal{C})$) at $z \in \mathcal{C}$

$$\lim_{\sigma \uparrow 1} (1 - \sigma) |\kappa_\sigma(z; \mathcal{C}) - \kappa_\sigma(z; \mathcal{C}_R)| = 0$$

4. Combine 2 & 3, show for arbitrary \mathcal{C} , $z \in \mathcal{C}$

$$\lim_{\sigma \uparrow 1} (1 - \sigma) \frac{1}{K(n)} \kappa_\sigma(z) = \kappa(z)$$

$$z \in \mathcal{C}_R, \ n = 2$$

$$\kappa_\sigma = \left(\int_{\mathcal{A}_{\text{even}}^+(z)} - \int_{\mathcal{A}_{\text{odd}}^+(z)} \right) \frac{(a \cdot t)b - (b \cdot t)a}{r^{1+\sigma}} d\mathcal{H}^2(a, b, r)$$

$$z \in \mathcal{C}_R, \ n = 2$$

$$\kappa_\sigma = \left(\int_{\mathcal{A}_{\text{even}}^+(z)} - \int_{\mathcal{A}_{\text{odd}}^+(z)} \right) \frac{(a \cdot t)b - (b \cdot t)a}{r^{1+\sigma}} d\mathcal{H}^2(a, b, r)$$

$$\mathbb{R}^2 = \text{span}\{t, n\}$$

$$z \in \mathcal{C}_R, \quad n = 2$$

$$\kappa_\sigma = \left(\int_{\mathcal{A}_{\text{even}}^+(z)} - \int_{\mathcal{A}_{\text{odd}}^+(z)} \right) \frac{(a \cdot t)b - (b \cdot t)a}{r^{1+\sigma}} d\mathcal{H}^2(a, b, r)$$

$$\mathbb{R}^2 = \text{span}\{t, n\}$$

$$\kappa_\sigma \cdot t = \left(\int_{\mathcal{A}_{\text{even}}^+(z)} - \int_{\mathcal{A}_{\text{odd}}^+(z)} \right) \frac{(a \cdot t)(b \cdot t) - (b \cdot t)(a \cdot t)}{r^{1+\sigma}} d\mathcal{H}^2(a, b, r) = 0$$

$$z \in \mathcal{C}_R, \quad n = 2$$

$$\kappa_\sigma = \left(\int_{\mathcal{A}_{\text{even}}^+(z)} - \int_{\mathcal{A}_{\text{odd}}^+(z)} \right) \frac{(a \cdot t)b - (b \cdot t)a}{r^{1+\sigma}} d\mathcal{H}^2(a, b, r)$$

$$\mathbb{R}^2 = \text{span}\{t, n\}$$

$$\kappa_\sigma \cdot t = \left(\int_{\mathcal{A}_{\text{even}}^+(z)} - \int_{\mathcal{A}_{\text{odd}}^+(z)} \right) \frac{(a \cdot t)(b \cdot t) - (b \cdot t)(a \cdot t)}{r^{1+\sigma}} d\mathcal{H}^2(a, b, r) = 0$$

$$\kappa_\sigma \cdot n = \left(\int_{\mathcal{A}_{\text{even}}^+(z)} - \int_{\mathcal{A}_{\text{odd}}^+(z)} \right) \frac{(a \cdot t)(b \cdot n) - (b \cdot t)(a \cdot n)}{r^{1+\sigma}} d\mathcal{H}^2(a, b, r)$$

$$z \in \mathcal{C}_R, n = 2$$

$$\kappa_\sigma = \left(\int_{\mathcal{A}_{\text{even}}^+(z)} - \int_{\mathcal{A}_{\text{odd}}^+(z)} \right) \frac{(a \cdot t)b - (b \cdot t)a}{r^{1+\sigma}} d\mathcal{H}^2(a, b, r)$$


$$\mathbb{R}^2 = \text{span}\{t, n\}$$

$$\kappa_\sigma \cdot t = \left(\int_{\mathcal{A}_{\text{even}}^+(z)} - \int_{\mathcal{A}_{\text{odd}}^+(z)} \right) \frac{(a \cdot t)(b \cdot t) - (b \cdot t)(a \cdot t)}{r^{1+\sigma}} d\mathcal{H}^2(a, b, r) = 0$$

$$\begin{aligned} \kappa_\sigma \cdot n &= \left(\int_{\mathcal{A}_{\text{even}}^+(z)} - \int_{\mathcal{A}_{\text{odd}}^+(z)} \right) \frac{(a \cdot t)(b \cdot n) - (b \cdot t)(a \cdot n)}{r^{1+\sigma}} d\mathcal{H}^2(a, b, r) \\ &= \int_{\mathcal{A}^+(z)} \bar{\chi}_{\mathcal{A}_{\text{even}}^+(z)}(a, b, r) \frac{(a \cdot t)(b \cdot n) - (b \cdot t)(a \cdot n)}{r^{1+\sigma}} d\mathcal{H}^2(a, b, r) \end{aligned}$$


$$\bar{\chi}_{\mathcal{A}_{\text{even}}^+(z)} = \chi_{\mathcal{A}_{\text{even}}^+(z)} - \chi_{(\mathcal{A}_{\text{even}}^+(z))^c} = \chi_{\mathcal{A}_{\text{even}}^+(z)} - \chi_{\mathcal{A}_{\text{odd}}^+(z)}$$

$$z \in \mathcal{C}_R, n = 2$$



$$z \in \mathcal{C}_R, n = 2$$


$$b \cdot t > 0 \implies b = \operatorname{sgn}(a \cdot n) a^\perp$$



$$z \in \mathcal{C}_R, n = 2$$

$$b \cdot t > 0 \implies b = \operatorname{sgn}(a \cdot n) a^\perp$$


$$\text{Disks: } u := 2ra$$



$$z \in \mathcal{C}_R, n = 2$$

$$b \cdot t > 0 \implies b = \operatorname{sgn}(a \cdot n) a^\perp$$


$$\text{Disks: } u := 2ra$$



$$z \in \mathcal{C}_R, n = 2$$

$$b \cdot t > 0 \implies b = \operatorname{sgn}(a \cdot n) a^\perp$$


$$\text{Disks: } u := 2ra$$



$$z \in \mathcal{C}_R, n = 2$$

$$b \cdot t > 0 \implies b = \operatorname{sgn}(a \cdot n) a^\perp$$

$$\text{Disks: } u := 2ra$$




$$\Pi^+ := \left\{ u \in \mathbb{R}^2 \mid (u \cdot n) < 0 \text{ or } \frac{|u|}{2} < R \left(\frac{u}{|u|} \cdot n \right) \right\}$$

$$z \in \mathcal{C}_R, n = 2$$

$$b \cdot t > 0 \implies b = \operatorname{sgn}(a \cdot n) a^\perp$$

$$\text{Disks: } u := 2ra$$



$$\Pi^+ := \left\{ u \in \mathbb{R}^2 \mid (u \cdot n) < 0 \text{ or } \frac{|u|}{2} < R \left(\frac{u}{|u|} \cdot n \right) \right\}$$

$$\kappa_\sigma \cdot n = -2^{\sigma+1/2} \int_{\mathbb{R}^2} \frac{\bar{\chi}_{\Pi^+}(u) \operatorname{sgn}(u \cdot n)}{|u|^{2+\sigma}} d\mathcal{H}^2(u)$$

$$z \in \mathcal{C}_R, \ n = 2$$

$$\kappa_\sigma \cdot n = -2^{\sigma+1/2} \int_{\mathbb{R}^2} \frac{\bar{\chi}_{\Pi^+}(u) \operatorname{sgn}(u \cdot n)}{|u|^{2+\sigma}} d\mathcal{H}^2(u)$$

$$z \in \mathcal{C}_R, \ n = 2$$

$$\begin{aligned} \kappa_\sigma \cdot n &= -2^{\sigma+1/2} \int_{\mathbb{R}^2} \frac{\bar{\chi}_{\Pi^+}(u) \operatorname{sgn}(u \cdot n)}{|u|^{2+\sigma}} d\mathcal{H}^2(u) \\ &= -2^{\sigma+1/2} \frac{-2^{1-\sigma}}{\sigma R^\sigma} B\left(\frac{1}{2}, \frac{1-\sigma}{2}\right) \end{aligned}$$

$$z \in \mathcal{C}_R, \, n = 2$$

$$\begin{aligned} \kappa_\sigma \cdot n &= -2^{\sigma+1/2} \int_{\mathbb{R}^2} \frac{\bar{\chi}_{\Pi^+}(u) \operatorname{sgn}(u \cdot n)}{|u|^{2+\sigma}} d\mathcal{H}^2(u) \\ &= -2^{\sigma+1/2} \frac{-2^{1-\sigma}}{\sigma R^\sigma} \mathrm{B}\left(\frac{1}{2}, \frac{1-\sigma}{2}\right) \end{aligned}$$

$$z \in \mathcal{C}_R, n = 2$$

$$\kappa_\sigma \cdot n = -2^{\sigma+1/2} \int_{\mathbb{R}^2} \frac{\bar{\chi}_{\Pi^+}(u) \operatorname{sgn}(u \cdot n)}{|u|^{2+\sigma}} d\mathcal{H}^2(u)$$

$$= -2^{\sigma+1/2} \frac{-2^{1-\sigma}}{\sigma R^\sigma} B\left(\frac{1}{2}, \frac{1-\sigma}{2}\right)$$

$$\lim_{\sigma \uparrow 1} (1-\sigma)(\kappa_\sigma \cdot n) = 2^{3/2} \lim_{\sigma \uparrow 1} \frac{(1-\sigma)}{\sigma R^\sigma} B\left(\frac{1}{2}, \frac{1-\sigma}{2}\right)$$

$$z \in \mathcal{C}_R, n = 2$$

$$\kappa_\sigma \cdot n = -2^{\sigma+1/2} \int_{\mathbb{R}^2} \frac{\bar{\chi}_{\Pi^+}(u) \operatorname{sgn}(u \cdot n)}{|u|^{2+\sigma}} d\mathcal{H}^2(u)$$

$$= -2^{\sigma+1/2} \frac{-2^{1-\sigma}}{\sigma R^\sigma} \mathrm{B}\left(\frac{1}{2}, \frac{1-\sigma}{2}\right)$$

$$\lim_{\sigma \uparrow 1} (1-\sigma)(\kappa_\sigma \cdot n) = 2^{3/2} \lim_{\sigma \uparrow 1} \frac{(1-\sigma)}{\sigma R^\sigma} \mathrm{B}\left(\frac{1}{2}, \frac{1-\sigma}{2}\right)$$

$$z \in \mathcal{C}_R, n = 2$$

$$\kappa_\sigma \cdot n = -2^{\sigma+1/2} \int_{\mathbb{R}^2} \frac{\bar{\chi}_{\Pi^+}(u) \operatorname{sgn}(u \cdot n)}{|u|^{2+\sigma}} d\mathcal{H}^2(u)$$

$$= -2^{\sigma+1/2} \frac{-2^{1-\sigma}}{\sigma R^\sigma} \mathrm{B}\left(\frac{1}{2}, \frac{1-\sigma}{2}\right)$$

$$\lim_{\sigma \uparrow 1} (1-\sigma)(\kappa_\sigma \cdot n) = 2^{3/2} \lim_{\sigma \uparrow 1} \frac{(1-\sigma)}{\sigma R^\sigma} \mathrm{B}\left(\frac{1}{2}, \frac{1-\sigma}{2}\right)$$

$$= \frac{2^{3/2} \sqrt{\pi}}{R} \lim_{\sigma \uparrow 1} (1-\sigma) \Gamma\left(\frac{1-\sigma}{2}\right)$$

$$z \in \mathcal{C}_R, n = 2$$

$$\kappa_\sigma \cdot n = -2^{\sigma+1/2} \int_{\mathbb{R}^2} \frac{\bar{\chi}_{\Pi^+}(u) \operatorname{sgn}(u \cdot n)}{|u|^{2+\sigma}} d\mathcal{H}^2(u)$$

$$= -2^{\sigma+1/2} \frac{-2^{1-\sigma}}{\sigma R^\sigma} \mathrm{B}\left(\frac{1}{2}, \frac{1-\sigma}{2}\right)$$

$$\lim_{\sigma \uparrow 1} (1-\sigma)(\kappa_\sigma \cdot n) = 2^{3/2} \lim_{\sigma \uparrow 1} \frac{(1-\sigma)}{\sigma R^\sigma} \mathrm{B}\left(\frac{1}{2}, \frac{1-\sigma}{2}\right)$$

$$= \frac{2^{3/2} \sqrt{\pi}}{R} \lim_{\sigma \uparrow 1} (1-\sigma) \Gamma\left(\frac{1-\sigma}{2}\right) = 2^{5/2} \sqrt{\pi} \frac{1}{R}$$

$$z \in \mathcal{C}_R, n = 2$$

$$\kappa_\sigma \cdot n = -2^{\sigma+1/2} \int_{\mathbb{R}^2} \frac{\bar{\chi}_{\Pi^+}(u) \operatorname{sgn}(u \cdot n)}{|u|^{2+\sigma}} d\mathcal{H}^2(u)$$

$$= -2^{\sigma+1/2} \frac{-2^{1-\sigma}}{\sigma R^\sigma} \mathrm{B}\left(\frac{1}{2}, \frac{1-\sigma}{2}\right)$$

$$\lim_{\sigma \uparrow 1} (1-\sigma)(\kappa_\sigma \cdot n) = 2^{3/2} \lim_{\sigma \uparrow 1} \frac{(1-\sigma)}{\sigma R^\sigma} \mathrm{B}\left(\frac{1}{2}, \frac{1-\sigma}{2}\right)$$

$$= \frac{2^{3/2} \sqrt{\pi}}{R} \lim_{\sigma \uparrow 1} (1-\sigma) \Gamma\left(\frac{1-\sigma}{2}\right) = 2^{5/2} \sqrt{\pi} \frac{1}{R}$$

$$z \in \mathcal{C}_R, n = 2$$

$$\kappa_\sigma \cdot n = -2^{\sigma+1/2} \int_{\mathbb{R}^2} \frac{\bar{\chi}_{\Pi^+}(u) \operatorname{sgn}(u \cdot n)}{|u|^{2+\sigma}} d\mathcal{H}^2(u)$$

$$= -2^{\sigma+1/2} \frac{-2^{1-\sigma}}{\sigma R^\sigma} B\left(\frac{1}{2}, \frac{1-\sigma}{2}\right)$$

$$\lim_{\sigma \uparrow 1} (1-\sigma)(\kappa_\sigma \cdot n) = 2^{3/2} \lim_{\sigma \uparrow 1} \frac{(1-\sigma)}{\sigma R^\sigma} B\left(\frac{1}{2}, \frac{1-\sigma}{2}\right)$$

$$= \frac{2^{3/2} \sqrt{\pi}}{R} \lim_{\sigma \uparrow 1} (1-\sigma) \Gamma\left(\frac{1-\sigma}{2}\right) = \underbrace{2^{5/2} \sqrt{\pi}}_{K_2} \underbrace{\frac{1}{R}}_{\kappa}$$

$$z \in \mathcal{C}_R, \ n \geq 2$$

$$\kappa_\sigma(z) := \left(\int_{\mathcal{A}_{\text{even}}^+(z)} - \int_{\mathcal{A}_{\text{odd}}^+(z)} \right) \frac{(a \cdot t(z))b - (b \cdot t(z))a}{r^{1+\sigma}} d\mathcal{H}^{2n-2}(a, b, r)$$

$$z \in \mathcal{C}_R, n \geq 2$$

$$\kappa_\sigma(z) := \left(\int_{\mathcal{A}_{\text{even}}^+(z)} - \int_{\mathcal{A}_{\text{odd}}^+(z)} \right) \frac{(a \cdot t(z))b - (b \cdot t(z))a}{r^{1+\sigma}} d\mathcal{H}^{2n-2}(a, b, r)$$

1. Slice on intersection with $t \times n$ plane (i.e. u in 2d)

$$z \in \mathcal{C}_R, n \geq 2$$

$$\kappa_\sigma(z) := \left(\int_{\mathcal{A}_{\text{even}}^+(z)} - \int_{\mathcal{A}_{\text{odd}}^+(z)} \right) \frac{(a \cdot t(z))b - (b \cdot t(z))a}{r^{1+\sigma}} d\mathcal{H}^{2n-2}(a, b, r)$$

1. Slice on intersection with $t \times n$ plane (i.e. u in 2d)
2. Recognize $\kappa_\sigma \cdot n$ is only non-zero component

$$z \in \mathcal{C}_R, n \geq 2$$

$$\kappa_\sigma(z) := \left(\int_{\mathcal{A}_{\text{even}}^+(z)} - \int_{\mathcal{A}_{\text{odd}}^+(z)} \right) \frac{(a \cdot t(z))b - (b \cdot t(z))a}{r^{1+\sigma}} d\mathcal{H}^{2n-2}(a, b, r)$$

1. Slice on intersection with $t \times n$ plane (i.e. u in 2d)
2. Recognize $\kappa_\sigma \cdot n$ is only non-zero component

Conjecture Any curve in k dimensional subspace only depends on k -dimensional curvature

$$z \in \mathcal{C}_R, n \geq 2$$

$$\kappa_\sigma(z) := \left(\int_{\mathcal{A}_{\text{even}}^+(z)} - \int_{\mathcal{A}_{\text{odd}}^+(z)} \right) \frac{(a \cdot t(z))b - (b \cdot t(z))a}{r^{1+\sigma}} d\mathcal{H}^{2n-2}(a, b, r)$$

1. Slice on intersection with $t \times n$ plane (i.e. u in 2d)
2. Recognize $\kappa_\sigma \cdot n$ is only non-zero component

Conjecture Any curve in k dimensional subspace only depends on k -dimensional curvature

3. Slice on r , i.e. group disks with common u with common r

$$z \in \mathcal{C}_R, n \geq 2$$

$$\kappa_\sigma(z) := \left(\int_{\mathcal{A}_{\text{even}}^+(z)} - \int_{\mathcal{A}_{\text{odd}}^+(z)} \right) \frac{(a \cdot t(z))b - (b \cdot t(z))a}{r^{1+\sigma}} d\mathcal{H}^{2n-2}(a, b, r)$$

1. Slice on intersection with $t \times n$ plane (i.e. u in 2d)
2. Recognize $\kappa_\sigma \cdot n$ is only non-zero component

Conjecture Any curve in k dimensional subspace only depends on k -dimensional curvature

3. Slice on r , i.e. group disks with common u with common r
4. Some more coarea applications, recognition of spheres & beta functions...

$$z \in \mathcal{C}_R, n \geq 2$$

$$\kappa_\sigma(z) := \left(\int_{\mathcal{A}_{\text{even}}^+(z)} - \int_{\mathcal{A}_{\text{odd}}^+(z)} \right) \frac{(a \cdot t(z))b - (b \cdot t(z))a}{r^{1+\sigma}} d\mathcal{H}^{2n-2}(a, b, r)$$

1. Slice on intersection with $t \times n$ plane (i.e. u in 2d)
2. Recognize $\kappa_\sigma \cdot n$ is only non-zero component

Conjecture Any curve in k dimensional subspace only depends on k -dimensional curvature

3. Slice on r , i.e. group disks with common u with common r
4. Some more coarea applications, recognition of spheres & beta functions...

$$\kappa_\sigma \cdot n = -2^{\sigma-3/2} \omega_{n-3}^2 B\left(\frac{3}{2}, \frac{n-2}{2}\right) B\left(\frac{\sigma+2}{2}, \frac{n-2}{2}\right)$$

$$\int_{\mathbb{R}^2} \frac{\bar{\chi}_{\Pi^+}(u) \operatorname{sgn}(n \cdot u)}{|u|^{2+\sigma}} d\mathcal{H}^2(u)$$

$$z \in \mathcal{C}_R, n \geq 2$$

$$\kappa_\sigma(z) := \left(\int_{\mathcal{A}_{\text{even}}^+(z)} - \int_{\mathcal{A}_{\text{odd}}^+(z)} \right) \frac{(a \cdot t(z))b - (b \cdot t(z))a}{r^{1+\sigma}} d\mathcal{H}^{2n-2}(a, b, r)$$

1. Slice on intersection with $t \times n$ plane (i.e. u in 2d)
2. Recognize $\kappa_\sigma \cdot n$ is only non-zero component

Conjecture Any curve in k dimensional subspace only depends on k -dimensional curvature

3. Slice on r , i.e. group disks with common u with common r
4. Some more coarea applications, recognition of spheres & beta functions...

$$\kappa_\sigma \cdot n = -2^{\sigma-3/2} \omega_{n-3}^2 B\left(\frac{3}{2}, \frac{n-2}{2}\right) B\left(\frac{\sigma+2}{2}, \frac{n-2}{2}\right)$$

$$\int_{\mathbb{R}^2} \frac{\bar{\chi}_{\Pi^+}(u) \operatorname{sgn}(n \cdot u)}{|u|^{2+\sigma}} d\mathcal{H}^2(u)$$

$$z \in \mathcal{C}_R, n \geq 2$$

$$\kappa_\sigma(z) := \left(\int_{\mathcal{A}_{\text{even}}^+(z)} - \int_{\mathcal{A}_{\text{odd}}^+(z)} \right) \frac{(a \cdot t(z))b - (b \cdot t(z))a}{r^{1+\sigma}} d\mathcal{H}^{2n-2}(a, b, r)$$

1. Slice on intersection with $t \times n$ plane (i.e. u in 2d)
2. Recognize $\kappa_\sigma \cdot n$ is only non-zero component

Conjecture Any curve in k dimensional subspace only depends on k -dimensional curvature

3. Slice on r , i.e. group disks with common u with common r
4. Some more coarea applications, recognition of spheres & beta functions...

$$\kappa_\sigma \cdot n = -2^{\sigma-3/2} \omega_{n-3}^2 \mathrm{B}\left(\frac{3}{2}, \frac{n-2}{2}\right) \mathrm{B}\left(\frac{\sigma+2}{2}, \frac{n-2}{2}\right)$$

$$\boxed{\frac{-2^{1-\sigma}}{\sigma R^\sigma} \mathrm{B}\left(\frac{1}{2}, \frac{1-\sigma}{2}\right)}$$

$$z \in \mathcal{C}_R, n \geq 2$$

$$\begin{aligned} & \lim_{\sigma \uparrow 1} (1 - \sigma)(\kappa_\sigma \cdot n) \\ &= \frac{\omega_{n-3}^2}{\sqrt{2}} \mathrm{B}\left(\frac{3}{2}, \frac{n-2}{2}\right) \mathrm{B}\left(\frac{\sigma+2}{2}, \frac{n-2}{2}\right) \frac{(1-\sigma)}{\sigma R^\sigma} \mathrm{B}\left(\frac{1}{2}, \frac{1-\sigma}{2}\right) \end{aligned}$$

$$z \in \mathcal{C}_R, n \geq 2$$

$$\lim_{\sigma \uparrow 1} (1 - \sigma)(\kappa_\sigma \cdot n)$$

$$= \frac{\omega_{n-3}^2}{\sqrt{2}} \mathrm{B}\left(\frac{3}{2}, \frac{n-2}{2}\right) \mathrm{B}\left(\frac{\sigma+2}{2}, \frac{n-2}{2}\right) \boxed{\frac{(1-\sigma)}{\sigma R^\sigma} \mathrm{B}\left(\frac{1}{2}, \frac{1-\sigma}{2}\right)}$$

$$z \in \mathcal{C}_R, n \geq 2$$

$$\lim_{\sigma \uparrow 1} (1 - \sigma)(\kappa_\sigma \cdot n)$$

$$\begin{aligned}
 &= \frac{\omega_{n-3}^2}{\sqrt{2}} \mathrm{B}\left(\frac{3}{2}, \frac{n-2}{2}\right) \mathrm{B}\left(\frac{\sigma+2}{2}, \frac{n-2}{2}\right) \boxed{\frac{(1-\sigma)}{\sigma R^\sigma} \mathrm{B}\left(\frac{1}{2}, \frac{1-\sigma}{2}\right)} \\
 &= \omega_{n-3}^2 \sqrt{2\pi} \mathrm{B}\left(\frac{3}{2}, \frac{n-2}{2}\right) \mathrm{B}\left(\frac{\sigma+2}{2}, \frac{n-2}{2}\right) \frac{1}{R}
 \end{aligned}$$

$$z \in \mathcal{C}_R, n \geq 2$$

$$\begin{aligned} & \lim_{\sigma \uparrow 1} (1 - \sigma)(\kappa_\sigma \cdot n) \\ &= \frac{\omega_{n-3}^2}{\sqrt{2}} \mathrm{B}\left(\frac{3}{2}, \frac{n-2}{2}\right) \mathrm{B}\left(\frac{\sigma+2}{2}, \frac{n-2}{2}\right) \boxed{\frac{(1-\sigma)}{\sigma R^\sigma} \mathrm{B}\left(\frac{1}{2}, \frac{1-\sigma}{2}\right)} \\ &= \overbrace{\omega_{n-3}^2 \sqrt{2\pi} \mathrm{B}\left(\frac{3}{2}, \frac{n-2}{2}\right) \mathrm{B}\left(\frac{\sigma+2}{2}, \frac{n-2}{2}\right)}^{K(n)} \overbrace{\frac{1}{R}}^{\kappa} \end{aligned}$$

2nd order approximation

Want $\lim_{\sigma \uparrow 1} (1 - \sigma) |\kappa_\sigma(z; \mathcal{C}) - \kappa_\sigma(z; \mathcal{C}_R)| = 0$ for arbitrary \mathcal{C} , $\mathcal{C}_R = 1/\kappa$

2nd order approximation


Want $\lim_{\sigma \uparrow 1} (1 - \sigma) |\kappa_\sigma(z; \mathcal{C}) - \kappa_\sigma(z; \mathcal{C}_R)| = 0$ for arbitrary \mathcal{C} , $\mathcal{C}_R = 1/\kappa$

$$|\kappa_\sigma(z; \mathcal{C}) - \kappa_\sigma(z; \mathcal{C}_R)| \lesssim \int_{\mathcal{A}^+} \frac{|\bar{\chi}_{\mathcal{A}^+_{\text{even}}(\mathcal{C})} - \bar{\chi}_{\mathcal{A}^+_{\text{even}}(\mathcal{C}_R)}|}{r^{1+\sigma}}$$

2nd order approximation


Want $\lim_{\sigma \uparrow 1} (1 - \sigma) |\kappa_\sigma(z; \mathcal{C}) - \kappa_\sigma(z; \mathcal{C}_R)| = 0$ for arbitrary \mathcal{C} , $\mathcal{C}_R = 1/\kappa$

$$|\kappa_\sigma(z; \mathcal{C}) - \kappa_\sigma(z; \mathcal{C}_R)| \lesssim \int_{\mathcal{A}^+} \frac{|\bar{\chi}_{\mathcal{A}^+_{\text{even}}}(\mathcal{C}) - \bar{\chi}_{\mathcal{A}^+_{\text{even}}}(\mathcal{C}_R)|}{r^{1+\sigma}}$$




2nd order approximation

Want $\lim_{\sigma \uparrow 1} (1 - \sigma) |\kappa_\sigma(z; \mathcal{C}) - \kappa_\sigma(z; \mathcal{C}_R)| = 0$ for arbitrary \mathcal{C} , $\mathcal{C}_R = 1/\kappa$




2nd order approximation

Want $\lim_{\sigma \uparrow 1} (1 - \sigma) |\kappa_\sigma(z; \mathcal{C}) - \kappa_\sigma(z; \mathcal{C}_R)| = 0$ for arbitrary \mathcal{C} , $\mathcal{C}_R = 1/\kappa$




2nd order approximation

Want $\lim_{\sigma \uparrow 1} (1 - \sigma) |\kappa_\sigma(z; \mathcal{C}) - \kappa_\sigma(z; \mathcal{C}_R)| = 0$ for arbitrary \mathcal{C} , $\mathcal{C}_R = 1/\kappa$




2nd order approximation

Want $\lim_{\sigma \uparrow 1} (1 - \sigma) |\kappa_\sigma(z; \mathcal{C}) - \kappa_\sigma(z; \mathcal{C}_R)| = 0$ for arbitrary \mathcal{C} , $\mathcal{C}_R = 1/\kappa$



2nd order approximation


Want $\lim_{\sigma \uparrow 1} (1 - \sigma) |\kappa_\sigma(z; \mathcal{C}) - \kappa_\sigma(z; \mathcal{C}_R)| = 0$ for arbitrary \mathcal{C} , $\mathcal{C}_R = 1/\kappa$



C1 Disks whose boundary intersects ribbon are all that matter

2nd order approximation

Want $\lim_{\sigma \uparrow 1} (1 - \sigma) |\kappa_\sigma(z; \mathcal{C}) - \kappa_\sigma(z; \mathcal{C}_R)| = 0$ for arbitrary \mathcal{C} , $\mathcal{C}_R = 1/\kappa$



- C1 Disks whose boundary intersects ribbon are all that matter
- C2 With the right area & coarea applications the integral over relevant disks vanishes

Final Result

Arbitrary \mathcal{C} , $K(n) = \omega_{n-3}^2 \sqrt{2\pi} B\left(\frac{3}{2}, \frac{n-2}{2}\right) B\left(\frac{\sigma+2}{2}, \frac{n-2}{2}\right)$, $z \in \mathcal{C}$

$$\lim_{\sigma \uparrow 1} (1 - \sigma) \frac{1}{K(n)} \kappa_{\sigma}(z) = \kappa(z)$$

Final Result

Arbitrary \mathcal{C} , $K(n) = \omega_{n-3}^2 \sqrt{2\pi} B(\frac{3}{2}, \frac{n-2}{2}) B(\frac{\sigma+2}{2}, \frac{n-2}{2})$, $z \in \mathcal{C}$

$$\lim_{\sigma \uparrow 1} (1 - \sigma) \frac{1}{K(n)} \kappa_\sigma(z) = \kappa(z)$$

New definition?

$$\kappa_\sigma(z) := \frac{1}{K(n)} \left(\int_{\mathcal{A}_{\text{even}}^+(z)} - \int_{\mathcal{A}_{\text{odd}}^+(z)} \right) \frac{(a \cdot t(z))b - (b \cdot t(z))a}{r^{1+\sigma}} d\mathcal{H}^{2n-2}(a, b, r)$$

Fin