

RESEARCH ARTICLE

Interim sample size reestimation for adequately powered series of N-of-1 trials

Daphne N. Weemering*

¹Department of Methodology and Statistics,
Utrecht University, Utrecht, The
Netherlands

Correspondence

*Corresponding author name, This is sample
corresponding address. Email:
authorone@gmail.com

Present Address

Padualaan 14, 3584 CH Utrecht, The
Netherlands

Lorem ipsum dolor sit amet, consectetur adipiscing elit. Aenean ut elit odio. Donec fermentum tellus neque, vitae fringilla orci pretium vitae. Fusce maximus finibus facilisis. Donec ut ullamcorper turpis. Donec ut porta ipsum. Nullam cursus mauris a sapien ornare pulvinar. Aenean malesuada molestie erat quis mattis. Praesent scelerisque posuere faucibus. Praesent nunc nulla, ullamcorper ut ullamcorper sed, molestie ut est. Donec consequat libero nisi, non semper velit vulputate et. Quisque eleifend tincidunt ligula, bibendum finibus massa cursus eget. Curabitur aliquet vehicula quam non pulvinar. Aliquam facilisis tortor nec purus finibus, sit amet elementum eros sodales. Ut porta porttitor vestibulum. Integer molestie, leo ut maximus aliquam, velit dui iaculis nibh, eget hendrerit purus risus sit amet dolor. Sed sed tincidunt ex. Curabitur imperdiet egestas tellus in iaculis. Maecenas ante neque, pretium vel nisl at, lobortis lacinia neque. In gravida elit vel volutpat imperdiet. Sed ut nulla arcu. Proin blandit interdum ex sit amet laoreet. Phasellus efficitur, sem hendrerit mattis dapibus, nunc tellus ornare nisi, nec eleifend enim nibh ac ipsum. Aenean tincidunt nisl sit amet facilisis faucibus. Donec odio erat, bibendum eu imperdiet sed, gravida luctus turpis.

KEYWORDS:

N-of-1 trials; Sample size reestimation; Simulation study;

1 | INTRODUCTION

Randomized controlled trials (RCTs) are considered the gold standard in determining treatment efficacy in healthcare. At first glance, these standard RCTs seem to earn their position, as the randomization of patients into a parallel experimental and control condition works quite well in controlling factors that are not under experimental control, creating the possibility to make causal inferences about the usefulness of the intervention. A drawback, however, is that these standard RCTs require a relatively large sample size to establish the effectiveness of treatment with sufficient power. In all instances it is desirable to limit the number of subjects to enter a medical study, as it exposes individuals to potentially inferior treatment. However, for the instances of finding the right intervention for patients with rare diseases, standard RCTs become unfeasible due to the relatively large sample size it requires. As populations of patients with rare diseases only have a limited number of people that could enter such a experimental study, alternatives should be sought to obtain reliable estimates of the treatment effect.

A clinical trial methodology that offers the possibility to reduce the number of subjects necessary to find a treatment effect with sufficient power, is the N-of-1 trial. The N-of-1 trial is a randomized controlled multiple crossover trial where a single patient repeatedly receives the experiment and control intervention in a random order, spread out over multiple cycles¹. As

the experiment is conducted within a single patient, the advantage of the N-of-1 trial is that a patient-specific treatment effect estimate is obtained. Often, clinical evidence that is generated by standard RCTs turns out to have poor generalization and is therefore to a limited amount applicable to patients in ordinary practices². Also, treatments that are shown to be safe on average, may have a disbalance in risks and benefits to individual patients³. With an N-of-1 trial, these issues can be avoided by purely estimating the effect of treatment on a single patient.

To obtain reliable results, N-of-1 trials should be performed under specific clinical circumstances. First of all, the disease under study should be long term, symptomatic and stable over time in order to avoid that the treatment differences will be obscured within and between cycles due to disease progression. Second, the intervention should not modify the course of the disease and should have a rapid on- and offset of biological action of the medication^{4,5}.

A single N-of-1 trial does not compare itself with a standard RCT, as the results of a single trial are specific to an individual patient, and hence, cannot be generalized to the population. However, when the results of several separate N-of-1 trials are combined it is possible to estimate the population treatment effect⁶. Even though every patient is unique, there is also some similarity in the disease progression and the treatment response of patients. In the combined analysis of separate N-of-1 trials, both the magnitude of the treatment effect as well as the heterogeneity in treatment response are taken into account⁶. The combined estimate of the separate trials forms the basis of knowledge about the treatment, whereas the estimate of the heterogeneity in treatment response allows for determining optimal treatment for every single patient by balancing the similarities and differences.

Combined N-of-1 trials, now referred to as series of N-of-1 trials, have been performed in, inter alia, studying the influence of Mexiletine on nondystrophic myotonia⁷, studying the effectiveness of Methylphenidate on fatigue in patients with end-stage cancer⁸, and for investigating the usefulness of Sildenafil on Raynaud-Phenomenon patients⁹.

require a smaller sample size compared to standard RCTs, as every patient serves as its own control.

References

1. Guyatt G, Sackett D, Taylor DW, Ghong J, Roberts R, Pugsley S. Determining optimal therapy—randomized trials in individual patients. *New England Journal of Medicine* 1986; 314(14): 889–892.
2. Greenfield S, Kravitz R, Duan N, Kaplan SH. Heterogeneity of treatment effects: implications for guidelines, payment, and quality assessment. *The American journal of medicine* 2007; 120(4): S3–S9.
3. Duan N, Kravitz RL, Schmid CH. Single-patient (n-of-1) trials: a pragmatic clinical decision methodology for patient-centered comparative effectiveness research. *Journal of clinical epidemiology* 2013; 66(8): S21–S28.
4. Araujo A, Julious S, Senn S. Understanding variation in sets of N-of-1 trials. *PloS one* 2016; 11(12): e0167167.
5. Nikles J, Mitchell GK, Schluter P, et al. Aggregating single patient (n-of-1) trials in populations where recruitment and retention was difficult: the case of palliative care. *Journal of clinical epidemiology* 2011; 64(5): 471–480.
6. Zucker D, Schmid C, McIntosh M, D'agostino R, Selker H, Lau J. Combining single patient (N-of-1) trials to estimate population treatment effects and to evaluate individual patient responses to treatment. *Journal of clinical epidemiology* 1997; 50(4): 401–410.
7. Stunnenberg BC, Raaphorst J, Groenewoud HM, et al. Effect of mexiletine on muscle stiffness in patients with nondystrophic myotonia evaluated using aggregated N-of-1 trials. *Jama* 2018; 320(22): 2344–2353.
8. Mitchell GK, Hardy JR, Nikles CJ, et al. The effect of methylphenidate on fatigue in advanced cancer: an aggregated N-of-1 trial. *Journal of pain and symptom management* 2015; 50(3): 289–296.
9. Roustit M, Gai J, Gaget O, et al. On-demand sildenafil as a treatment for Raynaud phenomenon: a series of N-of-1 trials. *Annals of Internal Medicine* 2018; 169(10): 694–703.

