

# Psychiatric ‘diseases’ *versus* behavioral disorders and degree of genetic influence

O. J. Bienvenu<sup>1\*</sup>, D. S. Davydow<sup>2</sup> and K. S. Kendler<sup>3</sup>

<sup>1</sup> Department of Psychiatry and Behavioral Sciences, Johns Hopkins University School of Medicine, Baltimore, MD, USA

<sup>2</sup> Department of Psychiatry and Behavioral Sciences, University of Washington School of Medicine, Seattle, WA, USA

<sup>3</sup> Department of Psychiatry, Virginia Institute for Psychiatric and Behavioral Genetics, Virginia Commonwealth University, Richmond, VA, USA

**Background.** Psychiatric conditions in which symptoms arise involuntarily (‘diseases’) might be assumed to be more heritable than those in which *choices* are essential (behavioral disorders). We sought to determine whether psychiatric ‘diseases’ (Alzheimer’s disease, schizophrenia, and mood and anxiety disorders) are more heritable than behavioral disorders (substance use disorders and anorexia nervosa).

**Method.** We reviewed the literature for recent quantitative summaries of heritabilities. When these were unavailable, we calculated weighted mean heritabilities from twin studies meeting modern methodological standards.

**Results.** Heritability summary estimates were as follows: bipolar disorder (85%), schizophrenia (81%), Alzheimer’s disease (75%), cocaine use disorder (72%), anorexia nervosa (60%), alcohol dependence (56%), sedative use disorder (51%), cannabis use disorder (48%), panic disorder (43%), stimulant use disorder (40%), major depressive disorder (37%), and generalized anxiety disorder (28%).

**Conclusions.** No systematic relationship exists between the disease-like character of a psychiatric disorder and its heritability; many behavioral disorders seem to be more heritable than conditions commonly construed as diseases. These results suggest an error in ‘common-sense’ assumptions about the etiology of psychiatric disorders. That is, among psychiatric disorders, there is no close relationship between the strength of genetic influences and the etiologic importance of volitional processes.

Received 26 May 2009; Revised 28 March 2010; Accepted 31 March 2010; First published online 12 May 2010

**Key words:** Behavior, disease, genetic epidemiology, genetics.

## Introduction

Psychiatric patients present with a variety of kinds of difficulties. One way to coherently address this diversity is to recognize and use distinct methods of psychiatric reasoning for particular kinds of difficulties (McHugh & Slavney, 1998; Ghaemi, 2003; McHugh, 2005). One of the most intuitive distinctions in reasoning is between that for psychiatric conditions that manifest primarily with involuntary symptoms and that for conditions that manifest primarily as dysfunctional behaviors.

‘Disease’ reasoning in psychiatry generally assumes some primary disruption in brain functioning; thus, as noted by McHugh & Slavney (1998), a disease is something a patient *has*. Some diseases with prominent psychiatric symptoms have a known histopathology (e.g. Alzheimer’s disease), and some even a known etiology (e.g. Huntington’s disease), whereas

in other putative psychiatric diseases these characteristics are less clear. For example, psychiatrists routinely use disease reasoning in conceptualizing problems of patients with schizophrenia or mania; the spontaneous nature of the symptoms and their discontinuity from normal human experience suggest a primary disruption in brain functioning (McHugh & Slavney, 1998). Disease logic is also often used in conceptualizing the emergence of depressive or anxiety syndromes (Sheehan, 1983; McHugh & Slavney, 1998).

By contrast, ‘behavioral’ reasoning assumes that patients are ill because of what they are *doing* (McHugh & Slavney, 1998); thus, behavioral reasoning is strongly tied to individual choices, their antecedents, and their consequences. Some behavioral disorders, such as alcohol and drug use disorders, are usually construed as out-of-control motivated behaviors. Other behavioral disorders, such as anorexia nervosa, are usually construed as being shaped by culture and social learning (McHugh & Slavney, 1998). Clinicians typically treat behavioral disorders by attempting to persuade patients to choose to alter their behaviors (with help in disrupting the

\* Address for correspondence: Dr O. J. Bienvenu, Johns Hopkins University, 600 N. Wolfe St., Meyer 115, Baltimore, MD 21287, USA.  
(Email: jbienven@jhmi.edu)

physiologic drives and other factors that provoke and sustain the behaviors). Thus, behavioral reasoning, in particular, invokes learning principles.

A key distinction between these methods of reasoning is the role of volition. That is, symptoms of psychiatric diseases (memory disturbance, hallucinations and emotional disturbance) are typically understood as arising involuntarily. By contrast, behavioral disorders centrally involve voluntary control (though undoubtedly persons do not choose behavioral disorders). 'Common sense' might suggest that neurobiological and genetic factors should be especially influential in the genesis of psychiatric syndromes that are disease-like in their manifestations, whereas other forces might have greater impact on the genesis of syndromes that manifest with disturbed behaviors. Thus, common sense might suggest that any syndromes that centrally involve human decision making (Do I take this next drink? Do I eat this donut?) should be less genetically influenced than syndromes in which symptoms are unbidden. Such reasoning is consistent with the public's notion that behavioral disorders such as alcoholism are more 'self-inflicted' than schizophrenia, depression or Alzheimer's disease (Schomerus *et al.* 2006).

In recent years, genetic epidemiological studies of psychiatric syndromes have expanded to a sufficient degree to examine this issue empirically. We have reviewed the extant twin study literature to produce the best available estimates of the heritabilities of well-studied adult psychiatric syndromes, with the aim of answering the following simple question: are genetic influences consistently greater in disease-like psychiatric disorders than in syndromes of which the major manifestation is disordered behavior?

## Method

For the purposes of this review, we drew upon recent quantitative summaries of the heritabilities of the disorders of interest, if available. To identify relevant studies of phenotypes not previously summarized, we performed PubMed searches [e.g. 'Alzheimer Disease (MeSH Terms) AND twins (MeSH Terms)'] and we searched reference lists of source and review articles. We selected reports that met modern methodological standards for twin studies; that is, they used systematic ascertainment, data collection and diagnostic procedures, in addition to blinding to co-twin diagnostic status (and/or zygosity). In addition, we limited our focus to categorical phenotypes for which heritability had been estimated in at least three separate samples. If more than one study reported on identical participants, we only included the most informative with regard to heritability. We calculated mean heritability

estimates, weighted by the number of pairs in which at least one twin was affected. If the number of pairs with an affected twin was not reported, we calculated mean heritability estimates weighted by the total number of affected twins in each study. As used in this study, heritability is the proportion of individual differences in liability to an illness (in a particular population) that results from genetic differences between individuals.

## Results

### Diseases

We know of no previous quantitative summary of the heritability of Alzheimer's disease. Table 1 lists the five studies that met our inclusion criteria (Breitner *et al.* 1995; Raiha *et al.* 1996; Bergem *et al.* 1997; Gatz *et al.* 1997, 2005, 2006; Meyer & Breitner, 1998). The most comprehensive investigation, with the largest number of affected twins, is based on the Swedish Twin Registry (Gatz *et al.* 2005, 2006). There was some overlap, in terms of affected subjects, with a smaller previous study based on that registry (Gatz *et al.* 1997). The weighted mean heritability estimate was 75%, whether or not we included the earlier study (Gatz *et al.* 1997).

Sullivan *et al.* (2003) performed a meta-analysis of published twin studies of schizophrenia, incorporating ascertainment corrections. Although only four of the 12 twin studies met modern methodological standards, the results did not vary significantly between studies that did and did not meet these standards. The point estimate of heritability was 81%.

We know of no previous quantitative summary of the heritability of bipolar disorder using twin studies that meet modern methodological standards. Table 2 lists the three studies that met our inclusion criteria (Kendler *et al.* 1993, 1995; McGuffin *et al.* 2003; Kieseppa *et al.* 2004). Note that the affected subjects in the Maudsley study by Cardno *et al.* (1999) were included in the later study by McGuffin *et al.* (2003). The weighted mean heritability estimate was 85%.

Sullivan *et al.* (2000) conducted a meta-analysis of published twin studies of major depressive disorder in 2000; five studies met inclusion criteria. The point estimate of heritability was 37%. In a more recent large Swedish twin study, Kendler *et al.* (2006b) reported a similar heritability estimate for lifetime major depressive disorder (38%).

Hettema *et al.* (2001a) conducted a meta-analysis of published twin studies of panic disorder; three studies met inclusion criteria. The point estimate of heritability was 43%. Although two published twin studies of generalized anxiety disorder were meta-analyzed

**Table 1.** Twin studies of Alzheimer's disease

Study	Country	Age (years)	Basis for diagnosis	Diagnostic criteria	No. of pairs <sup>a</sup>	Heritability (%)
Breitner <i>et al.</i> (1995) Meyer & Breitner (1998)	USA	62–73	Nurse/physician examination, neuropsychological testing and laboratory tests	NINCDS-ADRDA, DSM-III-R or CERAD and NIA criteria for post-mortem diagnosis	36	74 <sup>b</sup>
Raiha <i>et al.</i> (1996) Pedersen <i>et al.</i> (2001)	Finland	≥18	Hospital discharge medical record review	NINCDS-ADRDA, DSM-III-R	84	63 <sup>c</sup>
Bergem <i>et al.</i> (1997) Pedersen <i>et al.</i> (2001)	Norway	≥65	Physician examination, laboratory tests	NINCDS-ADRDA, DSM-III-R	32	58 <sup>c</sup>
Gatz <i>et al.</i> (1997) Pedersen <i>et al.</i> (2001)	Sweden	≥54	Physician examination, neuropsychological testing and laboratory tests	NINCDS-ADRDA, DSM-III-R	40	78 <sup>c</sup>
Gatz <i>et al.</i> (2005, 2006)	Sweden	≥65	Physician examination, neuropsychological testing and laboratory tests	NINCDS-ADRDA, DSM-IV	392	79
Summary						75

NINCDS-ADRDA, National Institute of Neurological and Communicative Disorders and Stroke – Alzheimer's Disease and Related Disorders Association; CERAD, The Consortium to Establish a Registry for Alzheimer's Disease; NIA, National Institute on Aging.

<sup>a</sup> At least one twin affected.

<sup>b</sup> Best-fitting multiple-threshold (age at onset) model by Akaike's Information Criteria (note: models with no additive genetic component, only a shared environmental component, or a smaller additive genetic component plus a shared environmental component, fit almost as well). Models assumed that all subjects would develop Alzheimer's disease if subjects lived long enough.

<sup>c</sup> Estimated using a single-threshold model, not accounting for age (potentially inflates the estimate of a shared environmental component) (Pedersen *et al.* 2001).

**Table 2.** Twin studies of bipolar disorder

Study	Country	Age (years)	Basis for diagnosis	Diagnostic criteria	No. of pairs <sup>a</sup>	Heritability (%)
Kendler <i>et al.</i> (1993, 1995)	Sweden	37–57	Questionnaire	DSM-III-R	35	79
McGuffin <i>et al.</i> (2003)	UK	≥15	Medical record review, ± research interview, outside informants	DSM-IV	67	85
Kiesseppa <i>et al.</i> (2004)	Finland	53 ± 13 <sup>b</sup>	Medical record review, diagnostic interview, ± outside informants	DSM-IV	25	93
Summary						85

<sup>a</sup> At least one twin affected.

<sup>b</sup> Mean ± standard deviation.

by Hettema *et al.* (2001a), the authors excluded one eligible study because of its complex ascertainment scheme (Roy *et al.* 1995), and an additional eligible study has been published since the meta-analysis (Mackintosh *et al.* 2006). Table 3 lists the four studies

that met our inclusion criteria (Roy *et al.* 1995; Scherrer *et al.* 2000; Hettema *et al.* 2001b; Mackintosh *et al.* 2006); one study only included males (Scherrer *et al.* 2000). The weighted mean heritability estimate was 28%.

**Table 3.** Twin studies of generalized anxiety disorder

Study	Country	Age (years)	Basis for diagnosis	Diagnostic criteria	Number affected <sup>a</sup>	Heritability (%)
Roy <i>et al.</i> (1995)	Sweden	53 ± 13 <sup>b</sup>	Questionnaire	DSM-III-R, broadened	128	49
Scherrer <i>et al.</i> (2000)	USA	≥37	Diagnostic interview	DSM-III-R, broadened	827	37
Hettema <i>et al.</i> (2001b)	USA	≥20	Diagnostic interview	DSM-III-R, broadened	1173	22
Mackintosh <i>et al.</i> (2006)	Sweden	≥55	Diagnostic interview	DSM-III-R, broadened	1415	27
Summary						28

<sup>a</sup> Among subjects in complete pairs, estimated.<sup>b</sup> Mean ± standard deviation.**Table 4.** Twin studies of anorexia nervosa

Study	Country	Age (years)	Basis for diagnosis	Diagnostic criteria	No. of pairs <sup>a</sup>	Heritability (%)
Wade <i>et al.</i> (2000)	USA	≥17	Diagnostic interview	DSM-III-R, broadened	71	58 <sup>b</sup>
Klump <i>et al.</i> (2001)	USA	16–18	Diagnostic interview, questionnaire	DSM-IV, broadened	23	74
Bulik <i>et al.</i> (2006)	Sweden	≥40	Diagnostic interview, medical record review, questionnaire	DSM-IV	49	56
Summary						60

<sup>a</sup> At least one twin affected.<sup>b</sup> Estimated in best-fitting bivariate model with major depressive disorder.


### Behavioral disorders

Goldman *et al.* (2005) reviewed large unselected twin studies of substance use disorders. Weighted mean heritability estimates were as follows: cannabis use disorder (abuse or dependence), 43% (five studies); sedative use disorder, 51% (three studies); stimulant use disorder, 40% (three studies); cocaine use disorder, 72% (three studies); and alcohol dependence, 56% (five studies). We identified one more recent eligible study of cannabis use disorder (Kendler *et al.* 2006a), and in this study the heritability estimate was higher than most previous studies (77%); including it (total of six studies), the weighted mean heritability estimate was 48%.

We know of no previous quantitative summary of the heritability of anorexia nervosa. Table 4 lists the three studies that met our inclusion criteria (Wade *et al.* 2000; Klump *et al.* 2001; Bulik *et al.* 2006); these studies only included women. The weighted mean heritability estimate was 60%.

### Summary

Fig. 1 shows heritability summary estimates for psychiatric diseases/disease-like conditions and



**Fig. 1.** Heritability summary estimates for psychiatric diseases/disease-like conditions (■) and behavioral disorders (□).

behavioral disorders. Bipolar disorder, schizophrenia, Alzheimer's disease and cocaine use disorder had the highest heritabilities of the phenotypes addressed in this report: all ≥70%. The phenotypes with the next

highest heritabilities (40–60%) were anorexia nervosa, alcohol dependence, sedative use disorder, cannabis use disorder, panic disorder and stimulant use disorder. The least heritable phenotypes (heritabilities <40%) were major depressive disorder and generalized anxiety disorder.

## Discussion

Our results do not strongly support the prediction that psychiatric 'diseases' are more heritable than behavioral disorders. On the one hand, Alzheimer's disease and two disease-like conditions, schizophrenia and bipolar disorder, have very high heritabilities. On the other hand, panic, major depressive and generalized anxiety disorders have substantially lower heritabilities, lower than those of most of the behavioral disorders examined. Thus, there is little indication of a systematic relationship between the disease-like character of a psychiatric disorder and its heritability. Indeed, idiopathic Parkinson's disease, another neuropsychiatric condition with a known histopathology, does not seem to be particularly heritable (Wirdefeldt *et al.* 2004).

It might be argued that major depression and anxiety disorders should not be considered 'disease-like', as these conditions are frequently accompanied by long-standing personality vulnerabilities (Roth *et al.* 1972; Bienvenu *et al.* 2004, 2009). Major depression is probably a heterogeneous condition, including a more disease-like 'endogenous' form and a less disease-like 'neurotic' form (Roth, 2001; Shorter, 2009). Although some studies suggest that endogenous features are associated with higher heritability (Leckman *et al.* 1984*a,b*; McGuffin *et al.* 1996), aspects of depressive illness that are particularly salient for heritability include the related features recurrence, early onset, and perhaps short duration (Kendler *et al.* 2007). When adjusted for the latter features, endogenous symptoms do not uniquely predict heritability (Weissman *et al.* 1986; Kendler *et al.* 1994; McGuffin *et al.* 1996). In addition, the presence of co-morbid anxiety disorders is found to increase, rather than decrease, the heritability of depressive illness (Weissman *et al.* 1986; Kendler *et al.* 1994).

Our results are inconsistent with the notion that psychiatric problems can be divided accurately into those that are 'biologically' caused and involuntary *versus* those that involve choices. That is, our findings support the proposition that biological/genetic factors influence volitional behavior, although it seems unlikely that genes simply evoke particular problem behaviors. The mechanisms through which genetic factors influence behavior are probably numerous. For example, genes can alter the hedonic and/or adverse

effects of cannabis (Lyons *et al.* 1997) and alcohol (Thomasson & Li, 1993), making it more or less pleasant to consume these substances. Genes can also influence levels of sensation-seeking (Agrawal *et al.* 2004), making some individuals more prone to behaviors such as drug use that are rewarding in the short term but harmful in the long term. Other heritable temperamental traits may similarly influence the development of eating disorders (Wade *et al.* 2008). Although personal decisions are no doubt important in behavioral disorders, genes clearly influence their development. Thus, genes do not respect the boundary between 'free will' and 'determinism'.

One final implication of our findings is noteworthy. When we think about diseases in psychiatry, we often think not only about high heritability but also about low levels of personal responsibility and blame-worthiness. By contrast, when we consider behavioral disorders (especially substance use disorders), we sometimes think not only of reduced genetic influences but also of higher levels of personal responsibility and blame-worthiness. After all, because behavior is under at least partial voluntary control, it seems natural to assume that a behavioral disorder is 'the patient's fault'. Indirectly, our findings challenge this assumption. Given the apparent lack of a consistent relationship between heritability and disease *versus* disturbed behavior, should we also re-examine our conception about the relationship between disease and disturbed behavior on the one hand and moral responsibility and blame-worthiness on the other? Our results emphasize the important distinction between stigmatizing destructive behaviors in a treatment context and stigmatizing patients in a social (especially medical) context. Unfortunately, persons with behavioral disorders such as alcoholism are stigmatized, and not just by the general public (Farrell & Lewis, 1990; Schomerus *et al.* 2006; To & Vega, 2006). Given that vulnerability to behavioral disorders is under substantial genetic control, our results emphasize the inappropriateness of libertarian attitudes towards these devastating conditions. That is, we humans do not seem to be equally free in our decisions.

## Limitations

Three limitations deserve comment. First, it is important to recognize that the genetic influence on a phenotype (heritability) is not immutable; it is dependent upon the environment. Indeed, it is most accurate to characterize heritability of an illness in terms of a specific population at a particular point in time (Kendler, 2005). Because different environmental factors may be relevant for different psychiatric illnesses in particular populations, it is difficult to



address this limitation effectively. Nevertheless, within cohorts, the pattern of results in Fig. 1 seems to be consistent. For example, in studies using Swedish Twin Registry data with the same or overlapping birth cohorts, heritability estimates were as follows: bipolar disorder, 79%; Alzheimer's disease, 78–79%; anorexia nervosa, 56%; major depressive disorder, 38%; and generalized anxiety disorder, 27% (Kendler *et al.* 1995, 2006b; Gatz *et al.* 1997, 2006; Bulik *et al.* 2006; Mackintosh *et al.* 2006). Also, in studies using adult Virginia Twin Registry data, heritability estimates were as follows: drug use disorder, 66%; alcohol dependence, 49%; major depressive disorder, 39%; panic disorder, 37%; and generalized anxiety disorder, 22% (Kendler & Prescott, 1999; Hettema *et al.* 2001b; Kendler *et al.* 2001, 2003).

Second, unreliability, a form of measurement error, is confounded with unique environmental effects in twin models, and some diagnoses are more reliable than others in unselected samples. For example, in studies of adults using the Virginia Twin Registry, the reliabilities of lifetime diagnoses of panic, major depressive and generalized anxiety disorders tended to be lower ( $\kappa = 0.34\text{--}0.56$ ) than those of cocaine, sedative, cannabis and stimulant use disorders ( $\kappa = 0.47\text{--}0.68$ ) (Foley *et al.* 1998; Hettema *et al.* 2001b; Kendler *et al.* 2000, 2001). As no psychiatric diagnosis is made with complete reliability, heritability estimates tend to be underestimates, but the heritabilities of the disease-like conditions above would be expected to be slightly more underestimated than those of the substance use disorders. Nevertheless, it is unlikely that correcting heritability estimates for unreliability would substantially reorder the results in Fig. 1. In fact, the reliability of cocaine dependence ( $\kappa = 0.47$ ) was slightly lower than that of panic disorder ( $\kappa = 0.56$ ) in the studies cited above, yet the heritability estimates from these studies were 0.79 and 0.37 respectively for these diagnoses.

Third, etiologic heterogeneity is likely in psychiatric disorders, so our heritability estimates inevitably reflect an averaging across potentially different etiologies. For example, as noted earlier, certain clinical aspects of major depressive disorder are associated with higher heritability, including recurrence and early age-at-onset (Sullivan *et al.* 2000; Levinson *et al.* 2003). However, it is not clear that problems of etiologic heterogeneity would impact heritability estimates differentially for disease-like *versus* behavioral conditions.

### Acknowledgments

The US National Institutes of Health grant K23-MH64543 (Dr Bienvenu) supported this study. We

thank Drs P. McHugh, P. Slavney and G. Nestadt, and anonymous reviewers for helpful comments regarding this project.

### Declaration of Interest

None.

### References

- Agrawal A, Jacobson KC, Prescott CA, Kendler KS (2004). A twin study of personality and illicit drug use and abuse/dependence. *Twin Research* 7, 72–81.
- Bergem ALM, Engedal K, Kringlen E (1997). The role of heredity in late-onset Alzheimer disease and vascular dementia. *Archives of General Psychiatry* 54, 264–270.
- Bienvenu OJ, Samuels JF, Costa PT, Reti IM, Eaton WW, Nestadt G (2004). Anxiety and depressive disorders and the five-factor model of personality: a higher- and lower-order personality trait investigation in a community sample. *Depression and Anxiety* 20, 92–97.
- Bienvenu OJ, Stein MB, Samuels JF, Onyike CU, Eaton WW, Nestadt G (2009). Personality disorder traits as predictors of subsequent first-onset panic disorder or agoraphobia. *Comprehensive Psychiatry* 50, 209–214.
- Breitner JCS, Welsh KA, Gau BA, McDonald WM, Steffens DC, Saunders AM, Magruder KM, Helms MJ, Plassman BL, Folstein MF, Brandt J, Robinette CD, Page WF (1995). Alzheimer's disease in the National Academy of Sciences-National Research Council Registry of Aging Twin Veterans. III. Detection of cases, longitudinal results, and observations on twin concordance. *Archives of Neurology* 52, 763–771.
- Bulik CM, Sullivan PF, Tozzi F, Furberg H, Lichtenstein P, Pedersen NL (2006). Prevalence, heritability, and prospective risk factors for anorexia nervosa. *Archives of General Psychiatry* 63, 305–312.
- Cardno AG, Marshall EJ, Coid B, Macdonald AM, Ribchester TR, Davies NJ, Venturi P, Jones LA, Lewis SW, Sham PC, Gottesman II, Farmer AE, McGuffin P, Reveley AM, Murray RM (1999). Heritability estimates for psychotic disorders: the Maudsley twin psychosis series. *Archives of General Psychiatry* 56, 162–168.
- Farrell M, Lewis G (1990). Discrimination on the grounds of diagnosis. *British Journal of Addiction* 85, 883–890.
- Foley DL, Neale MC, Kendler KS (1998). Reliability of a lifetime history of major depression: implications for heritability and co-morbidity. *Psychological Medicine* 28, 857–870.
- Gatz M, Fratiglioni L, Johansson B, Berg S, Mortimer JA, Reynolds CA, Fiske A, Pedersen N (2005). Complete ascertainment of dementia in the Swedish Twin Registry: the HARMONY study. *Neurobiology of Aging* 26, 439–447.
- Gatz M, Pedersen NL, Berg S, Johansson B, Johansson K, Mortimer JA, Posner SF, Viitanen M, Winblad B, Ahlbom A (1997). Heritability for Alzheimer's disease: the study of dementia in Swedish twins. *Journals of Gerontology. Series A, Biological Sciences and Medical Sciences* 52, M117–M125.

- Gatz M, Reynolds CA, Fratiglioni L, Johansson B, Mortimer JA, Berg S, Fiske A, Pedersen NL (2006). Role of genes and environments for explaining Alzheimer disease. *Archives of General Psychiatry* **63**, 168–174.
- Ghaemi SN (2003). *The Concepts of Psychiatry*. Johns Hopkins University Press: Baltimore, MD.
- Goldman D, Oroszi G, Ducci F (2005). The genetics of addictions: uncovering the genes. *Nature Reviews Genetics* **6**, 521–532.
- Hettema JM, Neale MC, Kendler KS (2001a). A review and meta-analysis of the genetic epidemiology of anxiety disorders. *American Journal of Psychiatry* **158**, 1568–1578.
- Hettema JM, Prescott CA, Kendler KS (2001b). A population-based twin study of generalized anxiety disorder in men and women. *Journal of Nervous and Mental Disease* **189**, 413–420.
- Kendler KS (2005). Psychiatric genetics: a methodologic critique. *American Journal of Psychiatry* **162**, 3–11.
- Kendler KS, Aggen SH, Tambs K, Reichborn-Kjennerud T (2006a). Illicit psychoactive substance use, abuse and dependence in a population-based sample of Norwegian twins. *Psychological Medicine* **36**, 955–962.
- Kendler KS, Gardner CO, Prescott CA (2001). Panic syndromes in a population-based sample of male and female twins. *Psychological Medicine* **31**, 989–1000.
- Kendler KS, Gatz M, Gardner CO, Pedersen NL (2006b). A Swedish national twin study of lifetime major depression. *American Journal of Psychiatry* **163**, 109–114.
- Kendler KS, Gatz M, Gardner CO, Pedersen NL (2007). Clinical indices of familial depression in the Swedish Twin Registry. *Acta Psychiatrica Scandinavica* **115**, 214–220.
- Kendler KS, Karkowski LM, Neale MC, Prescott CA (2000). Illicit psychoactive substance use, heavy use, abuse, and dependence in a US population-based sample of male twins. *Archives of General Psychiatry* **57**, 261–269.
- Kendler KS, Neale MC, Kessler RC, Heath AC, Eaves LJ (1994). The clinical characteristics of major depression as indices of the familial risk to illness. *British Journal of Psychiatry* **165**, 66–72.
- Kendler KS, Pedersen N, Johnson L, Neale MC, Mathe AA (1993). A pilot twin study of affective illness, including hospital- and population-ascertained subsamples. *Archives of General Psychiatry* **50**, 699–706.
- Kendler KS, Pedersen NL, Neale MC, Mathe AA (1995). A pilot Swedish twin study of affective illness including hospital- and population-ascertained subsamples: results of model fitting. *Behavior Genetics* **25**, 217–232.
- Kendler KS, Prescott CA (1999). A population-based twin study of lifetime major depression in men and women. *Archives of General Psychiatry* **56**, 39–44.
- Kendler KS, Prescott CA, Myers J, Neale MC (2003). The structure of genetic and environmental risk factors for common psychiatric and substance use disorders in men and women. *Archives of General Psychiatry* **60**, 929–937.
- Kieseppa T, Partonen T, Haukka J, Kaprio J, Lonnqvist J (2004). High concordance of bipolar I disorder in a nationwide sample of twins. *American Journal of Psychiatry* **161**, 1814–1821.
- Klump KL, Miller KB, Keel PK, McGue M, Iacono WG (2001). Genetic and environmental influences on anorexia nervosa syndromes in a population-based twin sample. *Psychological Medicine* **31**, 737–740.
- Leckman JF, Caruso KA, Prusoff BA, Weissman MM, Merikangas KR, Pauls DL (1984a). Appetite disturbance and excessive guilt in major depression: use of family study data to define depressive subtypes. *Archives of General Psychiatry* **41**, 839–844.
- Leckman JF, Weissman MM, Prusoff BA, Caruso KA, Merikangas KR, Pauls DL, Kidd KK (1984b). Subtypes of depression: family study perspective. *Archives of General Psychiatry* **41**, 833–838.
- Levinson DF, Zubenko GS, Crowe RR, Depaulo JR, Scheftner WS, Weissman MM, Holmans P, Zubenko WN, Boutelle S, Murphy-Eberenz K, MacKinnon D, McInnis MG, Marta DH, Adams P, Sassoon S, Knowles JA, Thomas J, Chellis J (2003). Genetics of recurrent early-onset depression (GenRED): design and preliminary clinical characteristics of a repository sample for genetic linkage studies. *American Journal of Medical Genetics. Part B, Neuropsychiatric Genetics* **119B**, 118–130.
- Lyons MJ, Toomey R, Meyer JM, Green AI, Eisen SA, Goldberg J, True WR, Tsuang MT (1997). How do genes influence marijuana use? The role of subjective effects. *Addiction* **92**, 409–417.
- Mackintosh M-A, Gatz M, Wetherell JL, Pedersen NL (2006). A twin study of lifetime generalized anxiety disorder (GAD) in older adults: genetic and environmental influences shared by neuroticism and GAD. *Twin Research and Human Genetics* **9**, 30–37.
- McGuffin P, Katz R, Watkins S, Rutherford J (1996). A hospital-based twin register of the heritability of DSM-IV unipolar depression. *Archives of General Psychiatry* **53**, 129–136.
- McGuffin P, Rijdsdijk F, Andrew M, Sham P, Katz R, Cardno A (2003). The heritability of bipolar affective disorder and the genetic relationship to unipolar depression. *Archives of General Psychiatry* **60**, 497–502.
- McHugh PR (2005). Striving for coherence: psychiatry's efforts over classification. *Journal of the American Medical Association* **293**, 2526–2528.
- McHugh PR, Slavney PR (1998). *The Perspectives of Psychiatry*. Johns Hopkins University Press: Baltimore, MD.
- Meyer JM, Breitner JCS (1998). Multiple threshold model for the onset of Alzheimer's disease in the NAS-NRC Twin Panel. *American Journal of Medical Genetics* **81**, 92–97.
- Pedersen NL, Posner SF, Gatz M (2001). Multiple-threshold models for genetic influences on age of onset for Alzheimer's disease: findings in Swedish twins. *American Journal of Medical Genetics* **105**, 724–728.
- Raiha I, Kaprio J, Koskenvuo M, Rajala T, Sourander L (1996). Alzheimer's disease in Finnish twins. *Lancet* **347**, 573–578.
- Roth M (2001). Unitary or binary nature of classification of depressive illness and its implications for the scope of manic depressive disorder. *Journal of Affective Disorders* **64**, 1–18.
- Roth M, Gurney C, Garside RF, Kerr TA (1972). Studies in the classification of affective disorders: the relationship

- between anxiety states and depressive illnesses: I. *British Journal of Psychiatry* **121**, 147–161.
- Roy M-A, Neale MC, Pedersen NL, Mathe AA, Kendler KS** (1995). A twin study of generalized anxiety disorder and major depression. *Psychological Medicine* **25**, 1037–1049.
- Scherrer JF, True WR, Xian H, Lyons MJ, Eisen SA, Goldberg J, Lin N, Tsuang MT** (2000). Evidence for genetic influences common and specific to symptoms of generalized anxiety and panic. *Journal of Affective Disorders* **57**, 25–35.
- Schomerus G, Matschinger H, Angermeyer MC** (2006). Alcoholism: illness beliefs and resource allocation preferences of the public. *Drug and Alcohol Dependence* **82**, 204–210.
- Sheehan DV** (1983). *The Anxiety Disease*. Bantam Books: New York.
- Shorter E** (2009). *Before Prozac: The Troubled History of Mood Disorders in Psychiatry*. Oxford University Press: New York, NY.
- Sullivan PF, Kendler KS, Neale MC** (2003). Schizophrenia as a complex trait: evidence from a meta-analysis of twin studies. *Archives of General Psychiatry* **60**, 1187–1192.
- Sullivan PF, Neale MC, Kendler KS** (2000). Genetic epidemiology of major depression: review and meta-analysis. *American Journal of Psychiatry* **157**, 1552–1562.
- Thomasson HR, Li TK** (1993). How alcohol and aldehyde dehydrogenase genes modify alcohol drinking, alcohol flushing, and the risk for alcoholism. *Alcohol Health and Research World* **17**, 167–172.
- To SE, Vega CP** (2006). Alcoholism and pathways to recovery: new survey results on views and treatment options. *Medscape General Medicine* **8**, 2.
- Wade TD, Bulik CM, Neale M, Kendler KS** (2000). Anorexia nervosa and major depression: shared genetic and environmental risk factors. *American Journal of Psychiatry* **157**, 469–471.
- Wade TD, Tiggemann M, Bulik CM, Fairburn CG, Wray NR, Martin NG** (2008). Shared temperament risk factors for anorexia nervosa: a twin study. *Psychosomatic Medicine* **70**, 239–244.
- Weissman MM, Merikangas KR, Wickramaratne P, Kidd KK, Prusoff BA, Leckman JF, Pauls DL** (1986). Understanding the clinical heterogeneity of major depression using family data. *Archives of General Psychiatry* **43**, 430–434.
- Wirdefeldt K, Gatz M, Schalling M, Pedersen NL** (2004). No evidence for heritability of Parkinson disease in Swedish twins. *Neurology* **63**, 305–311.