

# Cable Project

Darice, Elise, Sarah

December 7, 2017

## 1 Perturbation

### 1.1 PDE for perturbation function

$$\frac{\partial \psi}{\partial t} = c \frac{\partial^2 \psi}{\partial \xi^2} + \frac{\partial \psi}{\partial \xi} - (1 - \delta(V(\xi) - \theta))\psi \quad (1)$$

### 1.2 Separation of Variables

We assume a separation of variables,  $\psi(\xi, t) = S(\xi)T(t)$ , which leads to

$$\frac{1}{T} \frac{\partial T}{\partial t} = \frac{1}{S} \frac{\partial^2 S}{\partial \xi^2} + \frac{c}{S} \frac{\partial S}{\partial \xi} - 1 + \delta(V(\xi) - \theta) = \lambda$$

The time-dependent equation is a homogeneous ODE:

$$\frac{1}{T} \frac{dT}{dt} = \lambda \implies \int \frac{dT}{T} = \int \lambda dt \implies T(t) = e^{\lambda t} \quad (2)$$

For the spatial dependence, we split the domain of the equation into three regions, depending on the value of  $\delta(V(\xi) - \theta)$ , with the boundary condition that  $\lim_{\xi \rightarrow \pm\infty} \psi = 0$ :

- 1)  $\xi < 0$  and  $\delta(V(\xi) - \theta) = 0$
- 2)  $\xi > 0$  and  $\delta(V(\xi) - \theta) = 0$
- 3)  $\xi = 0$  and  $\delta(V(\xi) - \theta) = \infty$

#### 1.2.1 Region 1 and 2

In Region 1 and 2, we solve the equation:

$$\frac{d^2 S}{d\xi^2} + c \frac{dS}{d\xi} - (\lambda + 1)S = 0 \quad (3)$$

The boundary conditions are in Region 1:  $\lim_{\xi \rightarrow -\infty} S_1 = 0$ ; and in Region 2:  $\lim_{\xi \rightarrow \infty} S_2 = 0$ . The solutions have the form:

$$S_1(\xi) = c_1 e^{\frac{-c + \sqrt{c^2 + 4(\lambda + 1)}}{2} \xi} S_2(\xi) = c_2 e^{\frac{-c - \sqrt{c^2 + 4(\lambda + 1)}}{2} \xi} \quad (4)$$

### 1.2.2 Region 3

At Region 3, we integrate the equation at a small interval around the point of discontinuity to examine the effect  $\delta(V(\xi) - \theta)$  has on the solution. After separation of variables, the inhomogeneous PDE 1 becomes

$$\frac{1}{T} \frac{\partial T}{\partial t} = \frac{1}{S} \frac{\partial^2 S}{\partial \xi^2} + \frac{c}{S} \frac{\partial S}{\partial \xi} - 1 + \delta(V(\xi) - \theta) = \lambda$$

The spatial equation is

$$\frac{d^2 S}{d\xi^2} + c \frac{dS}{d\xi} - [\lambda + 1 + \delta(V(\xi) - \theta)]S = 0 \quad (5)$$

Since there is a discontinuity due to the dirac delta function here at  $V(\xi) = \theta$ ,  $\xi = 0$ , we integrate a small interval  $(+\epsilon, -\epsilon)$  around the discontinuity. Then, we taking the limit  $\epsilon \rightarrow 0$  gives us information as to how the solutions in Region 1 and 2 (4) connect in Region 3.

$$\int_{-\epsilon}^{+\epsilon} \frac{d^2 S}{d\xi^2} d\xi + c \int_{-\epsilon}^{+\epsilon} \frac{dS}{d\xi} d\xi - \int_{-\epsilon}^{+\epsilon} (\lambda + 1)S d\xi - \int_{-\epsilon}^{+\epsilon} \delta(V(\xi) - \theta)S d\xi = 0$$

For the first two terms, the fundamental theorem of calculus gives us the first derivative evaluated at the endpoints and the function evaluated at the endpoints.

$$\frac{dS}{d\xi}(+\epsilon) - \frac{dS}{d\xi}(-\epsilon) + c[S(+\epsilon) - S(-\epsilon)] - \int_{-\epsilon}^{+\epsilon} (\lambda + 1)S d\xi - \int_{-\epsilon}^{+\epsilon} \delta(V(\xi) - \theta)S d\xi = 0$$

Now we take the limit:

$$\lim_{\xi \rightarrow 0} \left[ \frac{dS}{d\xi}(+\epsilon) - \frac{dS}{d\xi}(-\epsilon) + c[S(+\epsilon) - S(-\epsilon)] - \int_{-\epsilon}^{+\epsilon} (\lambda + 1)S d\xi - \int_{-\epsilon}^{+\epsilon} \delta(V(\xi) - \theta)S d\xi \right] = 0$$

Of the remaining two integrals, the first one goes to zero as the bounds shrink the zero since the integral is continuous, even if  $S$  is not. The second integral involving the dirac delta function evaluates to  $S(0)$  by the *sampling property*, since we are specifically looking at the small region centered around where  $V(\xi) = \theta$ . Therefore we get:

$$\lim_{\xi \rightarrow 0} \left[ \frac{dS}{d\xi}(+\epsilon) - \frac{dS}{d\xi}(-\epsilon) + c[S(+\epsilon) - S(-\epsilon)] \right] = S(0)$$

$S$  must be continuous if we are assuming that the derivative  $\frac{dS}{d\xi}$  exists so  $S(0^+) = S(0^-)$  and the constants in front of the homogeneous solutions (4) are equal,  $c_1 = c_2 = K$ , revealing a relationship between the value of the spatial term and the discontinuity between its derivative at  $\xi = 0$ .

$$S(0) = \lim_{\xi \rightarrow 0} \left[ K \left( \frac{-c - \sqrt{c^2 + 4(\lambda + 1)}}{2} e^{\frac{-c - \sqrt{c^2 + 4(\lambda + 1)}}{2}} - \frac{-c + \sqrt{c^2 + 4(\lambda + 1)}}{2} e^{\frac{-c + \sqrt{c^2 + 4(\lambda + 1)}}{2}} \right) \right]$$

Taking the limit, we find the following, where  $S(0) \in \mathbb{R}$  and  $K \in \mathbb{R}$  since this is a physical system and we assume the solutions are real-valued.

$$S(0) = K \left( \frac{-c - \sqrt{c^2 + 4(\lambda + 1)}}{2} - \frac{-c + \sqrt{c^2 + 4(\lambda + 1)}}{2} \right) \quad (6)$$

$S$  is the spatial part of some arbitrary perturbation function that satisfies  $\lim_{\xi \rightarrow \pm\infty} S(\xi) = 0$ , and its value at zero could be either non-zero, or it could be 0. Since the boundary conditions provided do not fix  $S(0)$  or  $K$ , we will examine both cases. In the latter case, if  $S(0) = 0$  then  $K = c_1 = 0$  and we find the trivial solution of  $S = 0$  which gives us  $\psi(\xi, t) = 0$ . In the following section, we explore the former case.

### 1.3 Eigenvalues

Now to examine the family of solutions that depend on  $\lambda$ , we start with the  $\lambda = 0$  case. Assuming we can rescale  $S(0)$  to unity, eq (6) simplifies to:

$$1 = K \left( \frac{-c - \sqrt{c^2 + 4}}{2} - \frac{-c + \sqrt{c^2 + 4}}{2} \right) = -K\sqrt{c^2 + 4}$$

or

$$K = -\frac{1}{\sqrt{c^2 + 4}}$$

Plugging this constant back into our solutions for Region 1 and 2 (4) and multiplying it by the time-dependent solution (2), we find a solution for  $\psi(\xi, t)$ :

$$\psi(\xi, t) = \begin{cases} \frac{-1}{\sqrt{c^2 + 4}} e^{\frac{-c + \sqrt{c^2 + 4}}{2}}, & \xi < 0 \\ \frac{-1}{\sqrt{c^2 + 4}} e^{\frac{-c - \sqrt{c^2 + 4}}{2}}, & \xi > 0 \end{cases} \quad (7)$$

Recalling our traveling wave solution  $V(\xi)$  (??), we note that  $\psi(\xi, t) = V'(\xi)$ . This solution corresponds to  $v(x, t) = V(\xi) + \epsilon V'(\xi)$ , a Taylor expansion linearization around the traveling wave solution. This perturbation is independent of time. We plot  $v(x, t)$  with various values of  $\epsilon$  to show its effect on  $V(\xi)$ .




Figure 1:  $v(x, t) = V(\xi) + \epsilon V'(\xi)$  with various  $\epsilon$  values

## References

- [1] Haberman, Richard. *Applied Partial Differential Equations with Fourier Series and Boundary Value Problems*. Pearson.
- [2] Tara LaForce. *Green's Function Course Notes*.  
<https://web.stanford.edu/class/energy281/GreensFunctions.pdf>
- [3] University of California, Davis. *Linearizing Equations About Rest Points*.  
<https://www.math.ucdavis.edu/~temple/MAT22C/LecturesMat22CW15/5-LinearizedEquationsAboutRestPoints-22C-W15.pdf>
- [4] USPAS. *Table of Fourier Transform Pairs*.  
<http://uspas.fnal.gov/materials/110DU/FourierTransformPairs.pdf>