
1914



We let γ be the equator containing 67 and in the meridian orthogonal to γ and going through the midpoint C between γ_1 1'12' and γ_1 '11'0.

We let γ_{11} and γ_{12} be the equators bisecting 11 and 12 and γ_1 and γ_0 be the equators containing 1'10' and 0'10'.

Note that the two pairs are antipodal and one is contained in γ_{11}, the other in γ_{12}, so

$$\{P_1, P_2\} \subset \gamma_{11} \cap \gamma_{12}$$

The symmetry with respect to γ maps γ_1 into γ_{12} and γ_0 into γ_{11}. So γ_{11} and γ_{12} intersect transversally and

$$\gamma_{11} \cap \gamma_{12} = \{P_1, P_2\}$$

and $s(10)$ is one of the P_i's, say P_1.
By the construction rules of derived-one net 4 is different from both P_1 and P_2.
So P_1 and 4 determine a unique equator going through them, which is the bisector ℓ of 10. This ℓ goes also through P_2.
The symmetry with respect to m maps

γ_1 into γ_{11} and γ_0 into γ_{12}, so again it maps 4 in either P_1 or P_2. In particular it maps ℓ into ℓ. So ℓ must be m.

The symmetry with respect to C maps γ_1 into γ_0 and γ_{12} into γ_{11}, so it maps $\{P_1, P_2\}$ into $\{P_1, P_2\}$. So either $\{P_1, P_2\}$ are the north and south pole, (a) or $P_1 = C$ and P_2 its antipodal, & vice versa. (b)

But $s(P_1) = 10$ and the s is the symmetry with respect to γ, either s maps P_1 into itself, (b) or P_1 into P_2 (a).

So either $4 = P_1$ or $4 = P_2$, which is a contradiction