My solutions to

Deep Learning: Foundations and Concepts

Dario Miro Konopatzki

11 Structured Distributions

11.3

$$
\begin{aligned}
p(a=0) &= \sum_{b,c} p(0,b,c) \\
&= p(0,0,0) + p(0,0,1) + p(0,1,0) + p(0,1,1) \\
&= 0.192 + 0.144 + 0.048 + 0.216 \\
&= 0.6 \\
p(b=0) &= \sum_{a,c} p(a,0,c) \\
&= p(0,0,0) + p(0,0,1) + p(1,0,0) + p(1,0,1) \\
&= 0.192 + 0.144 + 0.192 + 0.064 \\
&= 0.592 \\
p(c=0) &= \sum_{a,b} p(a,b,0) \\
&= p(0,0,0) + p(0,1,0) + p(1,0,0) + p(1,1,0) \\
&= 0.192 + 0.048 + 0.192 + 0.048 \\
&= 0.48 \\
p(c=1) &= 1 - p(c=0) \\
&= 0.52
\end{aligned}
$$

$$
\begin{aligned}
p(a=0,b=0) &= \sum_{c} p(0,0,c) \\
&= p(0,0,0) + p(0,0,1) \\
&= 0.192 + 0.144
\end{aligned}
$$

$$
\begin{aligned}
&= 0.336 \\
p(a=0, c=0) &= \sum_b p(0,b,0) \\
&= p(0,0,0) + p(0,1,0) \\
&= 0.192 + 0.048 \\
&= 0.24 \\
p(a=0, c=1) &= \sum_b p(0,b,1) \\
&= p(0,0,1) + p(0,1,1) \\
&= 0.144 + 0.216 \\
&= 0.36 \\
p(a=1, c=0) &= \sum_b p(1,b,0) \\
&= p(1,0,0) + p(1,1,0) \\
&= 0.192 + 0.048 \\
&= 0.24 \\
p(b=0, c=0) &= \sum_a p(a,0,0) \\
&= p(0,0,0) + p(1,0,0) \\
&= 0.192 + 0.192 \\
&= 0.384 \\
p(b=0, c=1) &= \sum_a p(a,0,1) \\
&= p(0,0,1) + p(1,0,1) \\
&= 0.144 + 0.064 \\
&= 0.208 \\
p(b=1, c=0) &= \sum_a p(a,1,0) \\
&= p(0,1,0) + p(1,1,0) \\
&= 0.048 + 0.048 \\
&= 0.096
\end{aligned}
$$

$$
p(a=0|c=0) = \frac{p(a=0, c=0)}{p(c=0)}
$$

$$
\begin{aligned}
&= \frac{0.24}{0.48} \\
&= 0.5 \\
p(a=1|c=0) &= 1 - p(a=0|c=0) \\
&= 1 - 0.5 \\
&= 0.5 \\
p(a=0|c=1) &= \frac{p(a=0, c=1)}{p(c=1)} \\
&= \frac{0.36}{0.52} \\
&= \frac{9}{13} \\
p(a=1|c=1) &= 1 - p(a=0|c=1) \\
&= 1 - \frac{9}{13} \\
&= \frac{4}{13} \\
p(b=0|c=0) &= \frac{p(b=0, c=0)}{p(c=0)} \\
&= \frac{0.384}{0.48} \\
&= 0.8 \\
p(b=1|c=0) &= 1 - p(b=0|c=0) \\
&= 1 - 0.8 \\
&= 0.2 \\
p(b=0|c=1) &= \frac{p(b=0, c=1)}{p(c=1)} \\
&= \frac{0.208}{0.52} \\
&= 0.4 \\
p(b=1|c=1) &= 1 - p(b=0|c=1) \\
&= 1 - 0.4 \\
&= 0.6
\end{aligned}
$$

$$
p(a=0, b=0|c=0) = \frac{p(0,0,0)}{p(c=0)}
$$

$$
\begin{aligned}
&= \frac{0.192}{0.48} \\
&= 0.4 \\
&= 0.5 \cdot 0.8 \\
&= p(a=0|c=0)p(b=0|c=0) \\
p(a=0,b=1|c=0) &= \frac{p(0,1,0)}{p(c=0)} \\
&= \frac{0.048}{0.48} \\
&= 0.1 \\
&= 0.5 \cdot 0.2 \\
&= p(a=0|c=0)p(b=1|c=0) \\
p(a=1,b=0|c=0) &= \frac{p(1,0,0)}{p(c=0)} \\
&= \frac{0.192}{0.48} \\
&= 0.4 \\
&= 0.5 \cdot 0.8 \\
&= p(a=1|c=0)p(b=0|c=0) \\
p(a=1,b=1|c=0) &= \frac{p(1,1,0)}{p(c=0)} \\
&= \frac{0.048}{0.48} \\
&= 0.1 \\
&= 0.5 \cdot 0.2 \\
&= p(a=1|c=0)p(b=1|c=0)
\end{aligned}
$$

$$
\begin{aligned}
p(a=0,b=0|c=1) &= \frac{p(0,0,1)}{p(c=1)} \\
&= \frac{0.144}{0.52} \\
&= \frac{18}{65} \\
&= \frac{9}{13} \cdot 0.4 \\
&= p(a=0|c=1)p(b=0|c=1)
\end{aligned}
$$

$$
\begin{aligned}
p(a=0,b=1|c=1) &= \frac{p(0,1,1)}{p(c=1)} \\
&= \frac{0.216}{0.52} \\
&= \frac{27}{65} \\
&= \frac{9}{13}\cdot 0.6 \\
&= p(a=0|c=1)p(b=1|c=1) \\
p(a=1,b=0|c=1) &= \frac{p(1,0,1)}{p(c=1)} \\
&= \frac{0.064}{0.52} \\
&= \frac{8}{65} \\
&= \frac{4}{13}\cdot 0.4 \\
&= p(a=1|c=1)p(b=0|c=1) \\
p(a=1,b=1|c=1) &= \frac{p(1,1,1)}{p(c=1)} \\
&= \frac{0.096}{0.52} \\
&= \frac{12}{65} \\
&= \frac{4}{13}\cdot 0.6 \\
&= p(a=1|c=1)p(b=1|c=1)
\end{aligned}
$$

11.4

$$
\begin{aligned}
p(a=1) &= 1-p(a=0) \\
&= 1-0.6 \\
&= 0.4
\end{aligned}
$$

$$
\begin{aligned}
p(c=0|a=0) &= \frac{p(a=0,c=0)}{p(a=0)} \\
&= \frac{0.24}{0.6}
\end{aligned}
$$

$$
\begin{aligned}
&= 0.4 \\
p(c=1|a=0) &= 1 - p(c=0|a=0) \\
&= 1 - 0.4 \\
&= 0.6 \\
p(c=0|a=1) &= \frac{p(a=1, c=0)}{p(a=1)} \\
&= \frac{0.24}{0.4} \\
&= 0.6 \\
p(c=1|a=1) &= 1 - p(c=0|a=1) \\
&= 1 - 0.6 \\
&= 0.4
\end{aligned}
$$

$$
\begin{aligned}
p(0,0,0) &= 0.192 \\
&= 0.6 \cdot 0.4 \cdot 0.8 \\
&= p(a=0)p(c=0|a=0)p(b=0|c=0) \\
p(0,0,1) &= 0.144 \\
&= 0.6 \cdot 0.6 \cdot 0.4 \\
&= p(a=0)p(c=1|a=0)p(b=0|c=1) \\
p(0,1,0) &= 0.048 \\
&= 0.6 \cdot 0.4 \cdot 0.2 \\
&= p(a=0)p(c=0|a=0)p(b=1|c=0) \\
p(0,1,1) &= 0.216 \\
&= 0.6 \cdot 0.6 \cdot 0.6 \\
&= p(a=0)p(c=1|a=0)p(b=1|c=1) \\
p(1,0,0) &= 0.192 \\
&= 0.4 \cdot 0.6 \cdot 0.8 \\
&= p(a=1)p(c=0|a=1)p(b=0|c=0) \\
p(1,0,1) &= 0.064 \\
&= 0.4 \cdot 0.4 \cdot 0.4 \\
&= p(a=1)p(c=1|a=1)p(b=0|c=1) \\
p(1,1,0) &= 0.048 \\
&= 0.4 \cdot 0.6 \cdot 0.2 \\
&= p(a=1)p(c=0|a=1)p(b=1|c=0)
\end{aligned}
$$

$$
\begin{aligned}
p(1,1,1) &= 0.096 \\
&= 0.4 \cdot 0.4 \cdot 0.6 \\
&= p(a=1)p(c=1|a=1)p(b=1|c=1)
\end{aligned}
$$



Directed graph corresponding to the factorization from 11.4.

11.11

$$
\begin{aligned}
p(a,b|d) &= \int p(a,b,c|d)dc \\
&= \int p(a|d)p(b,c|d)dc \qquad\qquad \text{by assumption } a \perp\!\!\!\perp b,c \mid d \\
&= p(a|d)\int p(b,c|d)dc \\
&= p(a|d)p(b|d)
\end{aligned}
$$

11.12

Let $A = \{x\}$, C Markov blanket of x, and B the remaining variables. Consider any path from x to any node in B. Such a path has to pass through some node $c \in C$.

1. If c parent of x and
   - path via child of c, then tail-to-tail at c.
   - path via parent of c, then head-to-tail at c.
2. If c child of x and
   - path via child of c, then head-to-tail at c.
   - path via parent d of c, then d co-parent of x (i.p. $d \in C$) and one continues by considering 3. below for d.
3. c co-parent of x and
   - path via child of c, then tail-to-tail at c.

- path via parent of c, then head-to-tail at c

By the above, and since $c \in C$, the path is blocked. Because it was chosen arbitrarily, all paths are blocked. Hence $\{x\}$ is d-separated from B by C.