

My solutions to  
Deep Learning: Foundations and Concepts

Dario Miro Konopatzki

## 11 Structured Distributions

### 11.3

$$\begin{aligned} p(a = 0) &= \sum_{b,c} p(0, b, c) \\ &= p(0, 0, 0) + p(0, 0, 1) + p(0, 1, 0) + p(0, 1, 1) \\ &= 0.192 + 0.144 + 0.048 + 0.216 \\ &= 0.6 \end{aligned}$$

$$\begin{aligned} p(b = 0) &= \sum_{a,c} p(a, 0, c) \\ &= p(0, 0, 0) + p(0, 0, 1) + p(1, 0, 0) + p(1, 0, 1) \\ &= 0.192 + 0.144 + 0.192 + 0.064 \\ &= 0.592 \end{aligned}$$

$$\begin{aligned} p(c = 0) &= \sum_{a,b} p(a, b, 0) \\ &= p(0, 0, 0) + p(0, 1, 0) + p(1, 0, 0) + p(1, 1, 0) \\ &= 0.192 + 0.048 + 0.192 + 0.048 \\ &= 0.48 \end{aligned}$$

$$\begin{aligned} p(c = 1) &= 1 - p(c = 0) \\ &= 0.52 \end{aligned}$$

$$\begin{aligned} p(a = 0, b = 0) &= \sum_c p(0, 0, c) \\ &= p(0, 0, 0) + p(0, 0, 1) \\ &= 0.192 + 0.144 \end{aligned}$$

$$\begin{aligned}
&= 0.336 \\
p(a = 0, c = 0) &= \sum_b p(0, b, 0) \\
&= p(0, 0, 0) + p(0, 1, 0) \\
&= 0.192 + 0.048 \\
&= 0.24 \\
p(a = 0, c = 1) &= \sum_b p(0, b, 1) \\
&= p(0, 0, 1) + p(0, 1, 1) \\
&= 0.144 + 0.216 \\
&= 0.36 \\
p(a = 1, c = 0) &= \sum_b p(1, b, 0) \\
&= p(1, 0, 0) + p(1, 1, 0) \\
&= 0.192 + 0.048 \\
&= 0.24 \\
p(b = 0, c = 0) &= \sum_a p(a, 0, 0) \\
&= p(0, 0, 0) + p(1, 0, 0) \\
&= 0.192 + 0.192 \\
&= 0.384 \\
p(b = 0, c = 1) &= \sum_a p(a, 0, 1) \\
&= p(0, 0, 1) + p(1, 0, 1) \\
&= 0.144 + 0.064 \\
&= 0.208 \\
p(b = 1, c = 0) &= \sum_a p(a, 1, 0) \\
&= p(0, 1, 0) + p(1, 1, 0) \\
&= 0.048 + 0.048 \\
&= 0.096 \\
p(a = 0|c = 0) &= \frac{p(a = 0, c = 0)}{p(c = 0)}
\end{aligned}$$

$$\begin{aligned}
&= \frac{0.24}{0.48} \\
&= 0.5 \\
p(a = 1|c = 0) &= 1 - p(a = 0|c = 0) \\
&= 1 - 0.5 \\
&= 0.5 \\
p(a = 0|c = 1) &= \frac{p(a = 0, c = 1)}{p(c = 1)} \\
&= \frac{0.36}{0.52} \\
&= \frac{9}{13} \\
p(a = 1|c = 1) &= 1 - p(a = 0|c = 1) \\
&= 1 - \frac{9}{13} \\
&= \frac{4}{13} \\
p(b = 0|c = 0) &= \frac{p(b = 0, c = 0)}{p(c = 0)} \\
&= \frac{0.384}{0.48} \\
&= 0.8 \\
p(b = 1|c = 0) &= 1 - p(b = 0|c = 0) \\
&= 1 - 0.8 \\
&= 0.2 \\
p(b = 0|c = 1) &= \frac{p(b = 0, c = 1)}{p(c = 1)} \\
&= \frac{0.208}{0.52} \\
&= 0.4 \\
p(b = 1|c = 1) &= 1 - p(b = 0|c = 1) \\
&= 1 - 0.4 \\
&= 0.6 \\
p(a = 0, b = 0|c = 0) &= \frac{p(0, 0, 0)}{p(c = 0)}
\end{aligned}$$

$$\begin{aligned}
&= \frac{0.192}{0.48} \\
&= 0.4 \\
&= 0.5 \cdot 0.8 \\
&= p(a = 0|c = 0)p(b = 0|c = 0) \\
p(a = 0, b = 1|c = 0) &= \frac{p(0, 1, 0)}{p(c = 0)} \\
&= \frac{0.048}{0.48} \\
&= 0.1 \\
&= 0.5 \cdot 0.2 \\
&= p(a = 0|c = 0)p(b = 1|c = 0) \\
p(a = 1, b = 0|c = 0) &= \frac{p(1, 0, 0)}{p(c = 0)} \\
&= \frac{0.192}{0.48} \\
&= 0.4 \\
&= 0.5 \cdot 0.8 \\
&= p(a = 1|c = 0)p(b = 0|c = 0) \\
p(a = 1, b = 1|c = 0) &= \frac{p(1, 1, 0)}{p(c = 0)} \\
&= \frac{0.048}{0.48} \\
&= 0.1 \\
&= 0.5 \cdot 0.2 \\
&= p(a = 1|c = 0)p(b = 1|c = 0) \\
p(a = 0, b = 0|c = 1) &= \frac{p(0, 0, 1)}{p(c = 1)} \\
&= \frac{0.144}{0.52} \\
&= \frac{18}{65} \\
&= \frac{9}{13} \cdot 0.4 \\
&= p(a = 0|c = 1)p(b = 0|c = 1)
\end{aligned}$$

$$\begin{aligned}
p(a = 0, b = 1|c = 1) &= \frac{p(0, 1, 1)}{p(c = 1)} \\
&= \frac{0.216}{0.52} \\
&= \frac{27}{65} \\
&= \frac{9}{13} \cdot 0.6 \\
&= p(a = 0|c = 1)p(b = 1|c = 1) \\
p(a = 1, b = 0|c = 1) &= \frac{p(1, 0, 1)}{p(c = 1)} \\
&= \frac{0.064}{0.52} \\
&= \frac{8}{65} \\
&= \frac{4}{13} \cdot 0.4 \\
&= p(a = 1|c = 1)p(b = 0|c = 1) \\
p(a = 1, b = 1|c = 1) &= \frac{p(1, 1, 1)}{p(c = 1)} \\
&= \frac{0.096}{0.52} \\
&= \frac{12}{65} \\
&= \frac{4}{13} \cdot 0.6 \\
&= p(a = 1|c = 1)p(b = 1|c = 1)
\end{aligned}$$

## 11.4


$$\begin{aligned}
p(a = 1) &= 1 - p(a = 0) \\
&= 1 - 0.6 \\
&= 0.4
\end{aligned}$$

$$\begin{aligned}
p(c = 0|a = 0) &= \frac{p(a = 0, c = 0)}{p(a = 0)} \\
&= \frac{0.24}{0.6}
\end{aligned}$$

$$\begin{aligned}
&= 0.4 \\
p(c = 1|a = 0) &= 1 - p(c = 0|a = 0) \\
&= 1 - 0.4 \\
&= 0.6 \\
p(c = 0|a = 1) &= \frac{p(a = 1, c = 0)}{p(a = 1)} \\
&= \frac{0.24}{0.4} \\
&= 0.6 \\
p(c = 1|a = 1) &= 1 - p(c = 0|a = 1) \\
&= 1 - 0.6 \\
&= 0.4
\end{aligned}$$

$$\begin{aligned}
p(0, 0, 0) &= 0.192 \\
&= 0.6 \cdot 0.4 \cdot 0.8 \\
&= p(a = 0)p(c = 0|a = 0)p(b = 0|c = 0) \\
p(0, 0, 1) &= 0.144 \\
&= 0.6 \cdot 0.6 \cdot 0.4 \\
&= p(a = 0)p(c = 1|a = 0)p(b = 0|c = 1) \\
p(0, 1, 0) &= 0.048 \\
&= 0.6 \cdot 0.4 \cdot 0.2 \\
&= p(a = 0)p(c = 0|a = 0)p(b = 1|c = 0) \\
p(0, 1, 1) &= 0.216 \\
&= 0.6 \cdot 0.6 \cdot 0.6 \\
&= p(a = 0)p(c = 1|a = 0)p(b = 1|c = 1) \\
p(1, 0, 0) &= 0.192 \\
&= 0.4 \cdot 0.6 \cdot 0.8 \\
&= p(a = 1)p(c = 0|a = 1)p(b = 0|c = 0) \\
p(1, 0, 1) &= 0.064 \\
&= 0.4 \cdot 0.4 \cdot 0.4 \\
&= p(a = 1)p(c = 1|a = 1)p(b = 0|c = 1) \\
p(1, 1, 0) &= 0.048 \\
&= 0.4 \cdot 0.6 \cdot 0.2 \\
&= p(a = 1)p(c = 0|a = 1)p(b = 1|c = 0)
\end{aligned}$$

$$\begin{aligned}
p(1, 1, 1) &= 0.096 \\
&= 0.4 \cdot 0.4 \cdot 0.6 \\
&= p(a = 1)p(c = 1|a = 1)p(b = 1|c = 1)
\end{aligned}$$



Directed graph corresponding to the factorization from 11.4.

### 11.11

$$\begin{aligned}
p(a, b|d) &= \int p(a, b, c|d)dc \\
&= \int p(a|d)p(b, c|d)dc && \text{by assumption } a \perp\!\!\!\perp b, c \mid d \\
&= p(a|d) \int p(b, c|d)dc \\
&= p(a|d)p(b|d)
\end{aligned}$$

### 11.12

Let  $A = \{x\}$ ,  $C$  Markov blanket of  $x$ , and  $B$  the remaining variables. Consider any path from  $x$  to any node in  $B$ . Such a path has to pass through some node  $c \in C$ .

1. If  $c$  parent of  $x$  and
  - path via child of  $c$ , then tail-to-tail at  $c$ .
  - path via parent of  $c$ , then head-to-tail at  $c$ .
2. If  $c$  child of  $x$  and
  - path via child of  $c$ , then head-to-tail at  $c$ .
  - path via parent  $d$  of  $c$ , then  $d$  co-parent of  $x$  (i.p.  $d \in C$ ) and one continues by considering 3. below for  $d$ .
3.  $c$  co-parent of  $x$  and
  - path via child of  $c$ , then tail-to-tail at  $c$ .

- path via parent of  $c$ , then head-to-tail at  $c$

By the above, and since  $c \in C$ , the path is blocked. Because it was chosen arbitrarily, all paths are blocked. Hence  $\{x\}$  is d-separated from  $B$  by  $C$ .

### 11.13

$$a \perp\!\!\!\perp b \mid \emptyset$$

$$\begin{aligned}
p(a, b) &= \sum_{\{c, d\}} p(a, b, c, d) \\
&= \sum_{\{c, d\}} p(a)p(b)p(c|a, b)p(d|c) \quad \text{graph structure} \\
&= p(a)p(b) \sum_c p(c|a, b) \sum_d p(d|c) \\
&= p(a)p(b) \sum_c (p(c|a, b) \cdot 1) \\
&= p(a)p(b)
\end{aligned}$$

$$a \not\perp\!\!\!\perp b \mid d$$

Consider the distribution and results from exercise 11.14 below. Then

$$\begin{aligned}
p(B = 0, F = 0 | D = 0) &= \frac{p(B = 0, F = 0, D = 0)}{p(D = 0)} \\
&= \frac{p(F = 0 | B = 0, D = 0)p(B = 0, D = 0)}{p(D = 0)} \\
&= \frac{\frac{41}{374} \cdot 0.0748}{0.352} \\
&= \frac{41}{1760} \\
&\neq \frac{289}{6400} \\
&= \frac{0.0748}{0.352} \cdot 0.2125 \\
&= \frac{p(B = 0, D = 0)}{p(D = 0)} p(F = 0 | D = 0) \\
&= p(B = 0 | D = 0) p(F = 0 | D = 0)
\end{aligned}$$



## 11.14

### Auxiliary Calculations

$$\begin{aligned}p(B = 0) &= 1 - p(B = 1) \\&= 1 - 0.9 \\&= 0.1\end{aligned}$$

$$\begin{aligned}p(F = 0) &= 1 - p(F = 1) \\&= 1 - 0.9 \\&= 0.1\end{aligned}$$

$$\begin{aligned}p(D = 0) &= \sum_G p(D = 0|G)p(G) \\&= p(D = 0|G = 0)p(G = 0) \\&\quad + p(D = 0|G = 1)p(G = 1) \\&= p(D = 0|G = 0)p(G = 0) \\&\quad + (1 - p(D = 1|G = 1))(1 - p(G = 0)) \\&= 0.9 \cdot 0.315 + (1 - 0.9) \cdot (1 - 0.315) \\&= 0.352\end{aligned}$$

$$\begin{aligned}p(G = 1|F = 0) &= 1 - p(G = 0|F = 0) \\&= 1 - 0.81 \\&= 0.19\end{aligned}$$

$$\begin{aligned}p(D = 0|G = 1) &= 1 - p(D = 1|G = 1) \\&= 1 - 0.9 \\&= 0.1\end{aligned}$$

$$\begin{aligned}p(G = 0|B = 0, F = 0) &= 1 - p(G = 1|B = 0, F = 0) \\&= 1 - 0.1 \\&= 0.9\end{aligned}$$

$$\begin{aligned}p(G = 0|B = 0, F = 1) &= 1 - p(G = 1|B = 0, F = 1) \\&= 1 - 0.2 \\&= 0.8\end{aligned}$$

$$\begin{aligned}
p(B = 0, D = 0) &= \sum_{F,G} p(B = 0, F, G, D = 0) \\
&= \sum_{F,G} p(B = 0) p(F) p(G|B = 0, F) p(D = 0|G) \\
&= p(B = 0) \sum_{F,G} p(F) p(G|B = 0, F) p(D = 0|G) \\
&= p(B = 0) \sum_F p(F) \underbrace{\sum_G p(G|B = 0, F) p(D = 0|G)}_{\star} \\
&= p(B = 0) (p(F = 0) \cdot \star_{[F=0]} + p(F = 1) \cdot \star_{[F=1]}) \\
&= 0.1 \cdot (0.1 \cdot 0.82 + 0.9 \cdot 0.74) \\
&= 0.0748
\end{aligned}$$

$$\begin{aligned}
\star_{[F=0]} &= p(G = 0|B = 0, F = 0) p(D = 0|G = 0) \\
&\quad + p(G = 1|B = 0, F = 0) p(D = 0|G = 1) \\
&= 0.9 \cdot .9 + 0.1 \cdot 0.1 \\
&= 0.82
\end{aligned}$$

$$\begin{aligned}
\star_{[F=1]} &= p(G = 0|B = 0, F = 1) p(D = 0|G = 0) \\
&\quad + p(G = 1|B = 0, F = 1) p(D = 0|G = 1) \\
&= 0.8 \cdot 0.9 + 0.2 \cdot 0.1 \\
&= 0.74
\end{aligned}$$

## Part 1

$$\begin{aligned}
p(F = 0|D = 0) &= \sum_G p(F = 0, G|D = 0) \\
&= \frac{1}{p(D = 0)} \sum_G p(F = 0, G, D = 0) \\
&= \frac{p(F = 0)}{p(D = 0)} \sum_G p(G|F = 0) p(D = 0|G) \\
&= \frac{p(F = 0)}{p(D = 0)} (p(G = 0|F = 0) p(D = 0|G = 0) \\
&\quad + p(G = 1|F = 0) p(D = 0|G = 1)) \\
&= \frac{0.1}{0.352} (0.81 \cdot 0.9 + 0.19 \cdot 0.1) \\
&= 0.2125
\end{aligned}$$

## Part 2

$$\begin{aligned}
p(F = 0|B = 0, D = 0) &= \sum_G p(F = 0, G|B = 0, D = 0) \\
&= \frac{1}{p(B = 0, D = 0)} \sum_G p(F = 0, G, B = 0, D = 0) \\
&= \frac{1}{p(B = 0, D = 0)} \sum_G p(B = 0)p(F = 0) \\
&\quad \cdot p(G|B = 0, F = 0)p(D = 0|G) \\
&= \frac{p(B = 0)p(F = 0)}{p(B = 0, D = 0)} \\
&\quad \cdot \sum_G p(G|B = 0, F = 0)p(D = 0|G) \\
&= \frac{p(B = 0)p(F = 0)}{p(B = 0, D = 0)} \\
&\quad \cdot (p(G = 0|B = 0, F = 0)p(D = 0|G = 0) \\
&\quad + p(G = 1|B = 0, F = 0)p(D = 0|G = 1)) \\
&= \frac{0.1 \cdot 0.1}{0.0748} (0.9 \cdot 0.9 + 0.1 \cdot 0.1) \\
&\approx 0.109 \\
&< 0.2125 \quad \text{Part 1} \\
&= p(F = 0|D = 0)
\end{aligned}$$

- Observing  $(D = 0)$  increases probability that  $(F = 0)$  compared to prior probability  $p(F = 0)$ .
- Observing  $(B = 0)$  explains away observation  $(D = 0)$ .