

Simulation and Analysis of 1D Wave Propagation under Various Physical Models

Dario Liotta



UNIVERSITÀ
DEGLI STUDI
DI PADOVA




Dipartimento
di Fisica
e Astronomia
Galileo Galilei

September 6th 2025


Course of **Quantum Information and Computing**
Academic Year 2024/2025

Numerical methods for differential equations

Ordinary Differential Equations




Partial Differential Equations




Numerical methods for differential equations

Ordinary Differential Equations



Partial Differential Equations



Introduction to the problem


Solving a **PDE** means to find a function u such that

$$\mathcal{L}u = f$$

where \mathcal{L} is a differential operator and f is a source term.

The equation holds in a domain Ω and is completed by prescribing **boundary conditions** on $\partial\Omega$.

In most physical
applications \mathcal{L} is a
second-order operator



Introduction to the problem


Solving a **PDE** means to find a function u such that

$$\mathcal{L}u = f$$

where \mathcal{L} is a differential operator and f is a source term.

The equation holds in a domain Ω and is completed by prescribing **boundary conditions** on $\partial\Omega$.

In most physical
applications \mathcal{L} is a
second-order operator



Weak formulation

Galerkin methods rely on a **weak formulation**

- Multiply by a **test function** v and integrate over the entire domain

$$-\int_{\Omega} (\Delta u) v d\Omega = \int_{\Omega} f v d\Omega$$

- Integrate by parts the left hand side

$$-\int_{\Omega} (\Delta u) v d\Omega = \int_{\Omega} \nabla u \cdot \nabla v d\Omega - \int_{\partial\Omega} \frac{\partial u}{\partial n} v ds$$

- Substitute and get the new expression

$$\int_{\Omega} \nabla u \cdot \nabla v d\Omega = \int_{\Omega} f v d\Omega + \int_{\partial\Omega} \frac{\partial u}{\partial n} v ds$$

Weak formulation

Galerkin methods rely on a **weak formulation**

- Multiply by a **test function** v and integrate over the entire domain

$$-\int_{\Omega} (\Delta u) v d\Omega = \int_{\Omega} f v d\Omega$$

- Integrate by parts the left hand side

$$-\int_{\Omega} (\Delta u) v d\Omega = \int_{\Omega} \nabla u \cdot \nabla v d\Omega - \int_{\partial\Omega} \frac{\partial u}{\partial n} v ds$$

- Substitute and get the new expression

$$\int_{\Omega} \nabla u \cdot \nabla v d\Omega = \int_{\Omega} f v d\Omega + \int_{\partial\Omega} \frac{\partial u}{\partial n} v ds$$

Weak formulation

Galerkin methods rely on a **weak formulation**

- Multiply by a **test function** v and integrate over the entire domain

$$-\int_{\Omega} (\Delta u) v d\Omega = \int_{\Omega} f v d\Omega$$

- Integrate by parts the left hand side

$$-\int_{\Omega} (\Delta u) v d\Omega = \int_{\Omega} \nabla u \cdot \nabla v d\Omega - \int_{\partial\Omega} \frac{\partial u}{\partial n} v ds$$

- Substitute and get the new expression

$$\int_{\Omega} \nabla u \cdot \nabla v d\Omega = \int_{\Omega} f v d\Omega + \int_{\partial\Omega} \frac{\partial u}{\partial n} v ds$$

Weak formulation

Galerkin methods rely on a **weak formulation**

- Multiply by a **test function** v and integrate over the entire domain

$$-\int_{\Omega} (\Delta u) v d\Omega = \int_{\Omega} f v d\Omega$$

- Integrate by parts the left hand side

$$-\int_{\Omega} (\Delta u) v d\Omega = \int_{\Omega} \nabla u \cdot \nabla v d\Omega - \int_{\partial\Omega} \frac{\partial u}{\partial n} v ds$$

- Substitute and get the new expression

$$\int_{\Omega} \nabla u \cdot \nabla v d\Omega = \int_{\Omega} f v d\Omega + \int_{\partial\Omega} \frac{\partial u}{\partial n} v ds$$

About the test function

The test function v is introduced to check whether the PDE is satisfied on average throughout the domain.

The problem becomes to find u such that

$$a(u, v) = F(v) \quad \forall v \in V$$

where

$$a(u, v) = \int_{\Omega} \nabla u \cdot \nabla v d\Omega \quad \text{is a bilinear form}$$

$$F(v) = \int_{\Omega} f v d\Omega + \int_{\partial\Omega} \frac{\partial u}{\partial n} v ds \quad \text{is a linear functional}$$

About the test function

The test function v is introduced to check whether the PDE is satisfied on average throughout the domain.

The problem becomes to find u such that

$$a(u, v) = F(v) \quad \forall v \in V$$

where

$$a(u, v) = \int_{\Omega} \nabla u \cdot \nabla v d\Omega \quad \text{is a bilinear form}$$

$$F(v) = \int_{\Omega} f v d\Omega + \int_{\partial\Omega} \frac{\partial u}{\partial n} v ds \quad \text{is a linear functional}$$

Benefits of the weak formulation

Strong formulation

$$u \in C^2(\Omega)$$

Holds pointwise in Ω

Derivatives exist classically

Weak formulation

$$u, v \in H^1(\Omega)^*$$

Holds on average on Ω

Derivatives exist in the
distributional sense

In short: weak formulation requires **less regularity**

* $H^1(\Omega)$ is a **Sobolev space** of functions with square-integrable first derivatives:

$$w \in H^1(\Omega) = \left\{ w \in L^2(\Omega) \mid \nabla w \in L^2(\Omega)^d \right\}$$

Benefits of the weak formulation

Strong formulation

$$u \in C^2(\Omega)$$

Holds pointwise in Ω

Derivatives exist classically

Weak formulation

$$u, v \in H^1(\Omega)^*$$

Holds on average on Ω

Derivatives exist in the
distributional sense

In short: weak formulation requires **less regularity**

* $H^1(\Omega)$ is a **Sobolev space** of functions with square-integrable first derivatives:

$$w \in H^1(\Omega) = \left\{ w \in L^2(\Omega) \mid \nabla w \in L^2(\Omega)^d \right\}$$

On boundary conditions

Another difference lies in the boundary condition prescription.



$v = 0$ on $\partial\Omega \Rightarrow$ cancels boundary term
(no information available on $\frac{\partial u}{\partial n}$)


v free on $\partial\Omega$

$u = g$ enforced on $\partial\Omega$ (final solution)

$\frac{\partial u}{\partial n} = h$ naturally enters weak form

On boundary conditions

Another difference lies in the boundary condition prescription.



$v = 0$ on $\partial\Omega \Rightarrow$ cancels boundary term
(no information available on $\frac{\partial u}{\partial n}$)



v free on $\partial\Omega$

$u = g$ enforced on $\partial\Omega$ (final solution)

$\frac{\partial u}{\partial n} = h$ naturally enters weak form

On boundary conditions

Another difference lies in the boundary condition prescription.



$\mathbf{v} = \mathbf{0}$ on $\partial\Omega \Rightarrow$ cancels boundary term
(no information available on $\frac{\partial u}{\partial n}$)

\mathbf{v} free on $\partial\Omega$

$\mathbf{u} = \mathbf{g}$ enforced on $\partial\Omega$ (final solution)

$\frac{\partial \mathbf{u}}{\partial n} = \mathbf{h}$ naturally enters weak form

Shape functions

Galerkin methods allow to find an approximate solution

$$u_h \in V_h \subset H^1(\Omega) \quad \text{where } V_h \text{ is a **finite-dimensional** space}$$

In this framework, the goal is to find u_h such that

$$a(u_h, v_h) = F(v_h) \quad \forall v_h \in V_h$$

A **basis of function** $\{\phi_i\}$ is chosen to express u_h and to use it as test:

$$u_h = \sum_{j=1}^N u_j \phi_j \implies a \left(\sum_{j=1}^N u_j \phi_j, \phi_i \right) = F(\phi_i) \quad \forall i = 1, \dots, N$$

Functions ϕ_i model the solution \longrightarrow **shape functions**

Shape functions

Galerkin methods allow to find an approximate solution

$$u_h \in V_h \subset H^1(\Omega) \quad \text{where } V_h \text{ is a **finite-dimensional** space}$$

In this framework, the goal is to find u_h such that

$$\boxed{a(u_h, v_h) = F(v_h) \quad \forall v_h \in V_h}$$

A **basis of function** $\{\phi_i\}$ is chosen to express u_h and to use it as test:

$$u_h = \sum_{j=1}^N u_j \phi_j \implies a \left(\sum_{j=1}^N u_j \phi_j, \phi_i \right) = F(\phi_i) \quad \forall i = 1, \dots, N$$

Functions ϕ_i model the solution \longrightarrow **shape functions**

Shape functions

Galerkin methods allow to find an approximate solution

$$u_h \in V_h \subset H^1(\Omega) \quad \text{where } V_h \text{ is a **finite-dimensional** space}$$

In this framework, the goal is to find u_h such that

$$\boxed{a(u_h, v_h) = F(v_h) \quad \forall v_h \in V_h}$$

A **basis of function** $\{\phi_i\}$ is chosen to express u_h and to use it as test:

$$u_h = \sum_{j=1}^N u_j \phi_j \implies a \left(\sum_{j=1}^N u_j \phi_j, \phi_i \right) = F(\phi_i) \quad \forall i = 1, \dots, N$$

Functions ϕ_i model the solution \longrightarrow **shape functions**

Mesh discretization

FEM approach consists in the subdivision of the domain in a so-called **mesh**

This choice brings several advantages:


- Good approximation of **complex geometries**
- Better capture of **local effects**
- Possibility of **adaptive refinement**
- Natural construction of a **global solution**

Mesh discretization


FEM approach consists in the subdivision of the domain in a so-called **mesh**

This choice brings several advantages:

- Good approximation of **complex geometries**
- Better capture of **local effects**
- Possibility of **adaptive refinement**
- Natural construction of a **global solution**



Elements



Title

In **F**inite **E**lement **M**ethod, V_h is generated by **local basis functions** with compact support. Usually, one discretizes the domain using a so-called mesh