

Multivariate Stochastic Calculus

Monday, February 13, 2012
4:56 PM

Sponsored by the National Grid Foundation **nationalgrid**
The power of action.

Def: d-dimensional Brownian Motion is a process

$W_t = (W_t^1, W_t^2, \dots, W_t^d)'$ with the following prop:

(i) each W_t^i is a Brownian Motion

(ii) if $i \neq j \Rightarrow W_t^i$ and W_t^j are independent

- associated with a d-dim Brownian Motion we have a filtration

$\{\mathbb{F}_t\}$ such that

(iii) (**Information accumulates**) : $\mathbb{F}_s \subseteq \mathbb{F}_t$ for all set

(iv) (**Adaptivity**) : W_t is \mathbb{F}_t measurable for all t

(v) (**Independence of future increments**) : for all $t \leq u$

$W_u - W_t$ is independent of \mathbb{F}_t

$$(i) \rightsquigarrow dW_t^i \cdot dW_t^i = dt$$

$$(ii) \rightsquigarrow dW_t^i \cdot dW_t^j = 0 \quad \text{for } i \neq j$$

(we need to justify this claim)

Proof : $0 = t_0 < t_1 < \dots < t_n = T$ partition of $[0, T]$

the sample cross-variation :

$$C_{\Pi} = \sum_{k=0}^{n-1}' [W_{t_{k+1}}^i - W_{t_k}^i] \cdot [W_{t_{k+1}}^j - W_{t_k}^j]$$

$$\cdot E C_{\Pi} = \sum_{k=0}^{n-1}' E[W_{t_{k+1}}^i - W_{t_k}^i] \cdot E[W_{t_{k+1}}^j - W_{t_k}^j] = 0$$

$$\cdot \text{Var } C_{\Pi} = E C_{\Pi}^2 = E \sum_{k=0}^{n-1}' (W_{t_{k+1}}^i - W_{t_k}^i)^2 (W_{t_{k+1}}^j - W_{t_k}^j)^2$$

$$= \sum_{k=0}^{n-1}' (t_{k+1} - t_k)^2 \leq \|\Pi\| \cdot T \xrightarrow{\text{as } \|\Pi\| \rightarrow 0} 0$$

$$\left. \begin{array}{l} E C_{\Pi} = 0 \\ \text{Var } C_{\Pi} \rightarrow 0 \end{array} \right\} \Rightarrow C_{\Pi} \xrightarrow[\text{as } \| \Pi \| \rightarrow 0]{} 0$$

Ito-Doeblin Formula for Multiple Processes

X_t, Y_t : Ito processes

$d=2$

$$W_t = \begin{pmatrix} W_t^1 \\ W_t^2 \end{pmatrix} \in \mathbb{R}^2$$

$$\left\{ \begin{array}{l} X_t = X_0 + \int_0^t \theta_u^1 du + \int_0^t \sigma_u^{11} dW_u^1 + \int_0^t \sigma_u^{12} dW_u^2 \\ Y_t = Y_0 + \int_0^t \theta_u^2 du + \int_0^t \sigma_u^{21} dW_u^1 + \int_0^t \sigma_u^{22} dW_u^2 \end{array} \right.$$

in vector form :

$$\begin{pmatrix} X_t \\ Y_t \end{pmatrix} = \begin{pmatrix} X_0 \\ Y_0 \end{pmatrix} + \int_0^t \theta_u du + \int_0^t \Gamma_u dW_u$$

$$\theta_t = \begin{pmatrix} \theta_t^1 \\ \theta_t^2 \end{pmatrix} \in \mathbb{R}^2 \quad \Gamma_t = \begin{pmatrix} \sigma_t^{11} & \sigma_t^{12} \\ \sigma_t^{21} & \sigma_t^{22} \end{pmatrix} \in \mathbb{R}^{2 \times 2}$$

$$d\langle X \rangle_t = dX_t \cdot dX_t = (\sigma_t^{11})^2 dt + (\sigma_t^{12})^2 dt$$

$$d\langle Y \rangle_t = dY_t \cdot dY_t = (\sigma_t^{21})^2 dt + (\sigma_t^{22})^2 dt$$

$$d\langle X, Y \rangle_t = dX_t \cdot dY_t = (\sigma_t^{11} \cdot \sigma_t^{21} + \sigma_t^{12} \cdot \sigma_t^{22}) dt$$

Two-dimensional Ito formula: $f(t, x, y)$ is \mathbb{C}^2 :

$f_t, f_x, f_y, f_{xx}, f_{xy}, f_{yy}$ are defined & continuous

$$\boxed{df(t, X_t, Y_t) = f_t(t, X_t, Y_t) dt + f_x(t, X_t, Y_t) dX_t + f_y(t, X_t, Y_t) dY_t + \frac{1}{2} f_{xx}(t, X_t, Y_t) d\langle X \rangle_t + f_{xy}(t, X_t, Y_t) d\langle X, Y \rangle_t + \frac{1}{2} f_{yy}(t, X_t, Y_t) d\langle Y \rangle_t}$$

example : $f(x,y) = xy \Rightarrow f_x = y, f_y = x$
 $f_{xx} = 0, f_{yy} = 0, f_{xy} = 1$

$$\boxed{d(X_t Y_t) = Y_t dX_t + X_t dY_t + d\langle XY \rangle_t}$$

↳ Stochastic version of the product rule

example : Correlated Stock Prices

- $\frac{dS_t^1}{S_t^1} = \alpha_1 dt + \sigma_1 dW_t^1 \quad \Leftrightarrow dS_t^1 = \alpha_1 S_t^1 dt + \sigma_1 S_t^1 dW_t^1$
- $\frac{dS_t^2}{S_t^2} = \alpha_2 dt + \sigma_2 \left[\rho dW_t^1 + \sqrt{1-\rho^2} dW_t^2 \right]$
 $= dW_t^3$

where W^1 and W^2 are independent B.M

$$\sigma_1 > 0, \sigma_2 > 0 \quad -1 \leq \rho \leq 1$$

- to analyze the second stock price, define :

$$\boxed{W_t^3 = \rho W_t^1 + \sqrt{1-\rho^2} W_t^2} \quad \text{martingale}$$

- is W^3 also a Brownian Motion also ?

we need to check if $d\langle W^3 \rangle_t = dt$

$$d\langle W^3 \rangle_t = dW_t^3 \cdot dW_t^3 = \rho^2 dt + (1-\rho^2) dt = dt$$

$\Rightarrow W_t^3$ is also a Brownian Motion.

So we can represent S^2 as :

$$\frac{dS_t^2}{S_t^2} = \alpha_2 dt + \sigma_2 dW_t^3 \quad (\text{GBM})$$

$$dW_t^1 W_t^3 = \underbrace{W_t^1 dW_t^3}_t + \underbrace{W_t^3 dW_t^1}_t + \underbrace{dW_t^1 \cdot dW_t^3}_{\rho dt}$$

$$W_t^1 W_t^3 = \int_0^t W_s^1 dW_s^3 + \int_0^t W_s^3 dW_s^1 + \rho t$$


$$\Rightarrow \mathbb{E} W_t^1 W_t^3 = \rho t$$

$$\text{corr}(W_t^1, W_t^3) = \frac{\mathbb{E}(W_t^1 \cdot W_t^3)}{t} = \rho$$

ρ = correlation between W^1 and W^3


[Correlated Wiener Processes \(Wolfram Demonstrations Project\)](#)

Correlated Wiener Processes




This Demonstration displays the paths of two correlated Wiener processes.

Correlated Wiener Processes



This Demonstration displays the paths of two correlated Wiener processes.

Correlated Wiener Processes



This Demonstration displays the paths of two correlated Wiener processes.