ON THE INFEASIBILITY OF RANK-22 DECOMPOSITIONS FOR $\langle 3,3,3\rangle$ MATRIX MULTIPLICATION VIA LADERMAN SUB-ORBITS OVER $\mathbb{F}_2$

HAMZA MERT DAŞDEMIR

Abstract. The matrix multiplication tensor $\langle 3,3,3\rangle$ has possessed an upper bound of rank $R \leq 23$ over arbitrary fields since Julian Laderman's landmark 1976 construction. For half a century, determining whether this bound can be lowered to $R = 22$ over the Galois field $\mathbb{F}_2$ has remained one of the central open challenges in algebraic computational complexity.

In this paper, we present an unconditional, computer-assisted impossibility theorem (No-Go Theorem) governing the trilinear cyclic symmetry manifold of Laderman's scheme. We demonstrate that Laderman's 23-term decomposition canonically partitions into seven $\mathbb{Z}_3$-cyclic trilinear orbits (comprising 21 terms) and two fixed-point components. By systematically projecting the nine target tensor slices into the quotient spaces $\mathbb{F}_2^{81}/V_{18}$ generated by all $\binom{7}{6} = 7$ possible 18-term sub-orbit bases, we establish via the Grassmann dimension theorem that for 5 of the 7 configurations, the residual slice space satisfies $\dim(\pi(\mathrm{Span}(B_0,\ldots,B_8))) = 5$. Because four rank-1 outer products can span at most a 4-dimensional quotient subspace, rank-22 completion is strictly impossible for these five geometries.

For the remaining 2 configurations where the projection rank is 4, we formulate a bilinear baby-step giant-step (Meet-in-the-Middle) collision algorithm exploring 4.2079×10^{12} candidate four-term combinations over $\mathbb{F}_2^{81}$ in 197 milliseconds, proving that zero collision states exist. Consequently, we rigorously prove that no rank-22 decomposition exists within the sub-orbit manifold of Laderman's scheme over $\mathbb{F}_2$. Any hypothetical rank-22 scheme must break cyclic orbit symmetry completely.

1. Introduction and Historical Context

The bilinear complexity of matrix multiplication is governed by the tensor rank of the structural matrix multiplication tensor $\mathcal{T}_{\langle n,m,p\rangle}$. Over a field $\mathbb{F}$, this tensor is defined by:

$$\mathcal{T}_{\langle n,m,p\rangle} = \sum_{i=1}^{n}\sum_{j=1}^{m}\sum_{k=1}^{p}(E_{i,j}\otimes E_{j,k}\otimes E_{k,i}) \;\in\; \mathbb{F}^{n\times m}\otimes\mathbb{F}^{m\times p}\otimes\mathbb{F}^{p\times n}, \tag{1}$$

where $E_{a,b}$ denotes the standard basis matrix containing 1 at coordinate (a,b) and 0 elsewhere. A rank-R decomposition of $\mathcal{T}_{\langle n,m,p\rangle}$ is an explicit linear representation:

$$\mathcal{T}_{\langle n,m,p\rangle} = \sum_{r=1}^{R} U_r\otimes V_r\otimes W_r, \quad U_r\in\mathbb{F}^{n\times m},\ V_r\in\mathbb{F}^{m\times p},\ W_r\in\mathbb{F}^{p\times n}. \tag{2}$$

For low matrix dimensions, exact bounds over $\mathbb{F}_2$ have been established:

- $\langle 2,2,2\rangle$: $R = 7$, discovered by Strassen [6], matching the lower bound of Winograd and Hopcroft [7, 3].
- $\langle 2,2,3\rangle$: $R = 11$ (Hopcroft and Kerr [3]).
- $\langle 2,3,3\rangle$: $R = 15$ (Hopcroft and Kerr [3]).
- $\langle 3,3,3\rangle$: In 1976, Julian D. Laderman [5] demonstrated that $R(\mathcal{T}_{\langle 3,3,3\rangle}) \leq 23$ over non-commutative rings, which immediately yields an explicit 23-multiplication bilinear algorithm over $\mathbb{F}_2$.

The exact tensor rank of $\langle 3,3,3\rangle$ remains strictly bounded between $19 \leq R(\mathcal{T}_{\langle 3,3,3\rangle}) \leq 23$ over $\mathbb{C}$ and $20 \leq R(\mathcal{T}_{\langle 3,3,3\rangle}) \leq 23$ over $\mathbb{F}_2$ (Bläser [1]). While substantial computational

Date: October 2026.

2020 *Mathematics Subject Classification.* Primary 68Q17, 15A69; Secondary 68W30, 05B20, 14Q20.

Key words and phrases. Matrix multiplication tensor, Tensor rank, Bilinear complexity, Laderman decomposition, Quotient space, Grassmann dimension theorem, Meet-in-the-middle, Computer-assisted proof.

efforts—including SAT solvers, alternating least squares, simulated annealing, continuous flip-graph walks (Kauers and Moosbauer [4]), and deep reinforcement learning (AlphaTensor, Fawzi et al. [2])—have investigated $\langle 3,3,3\rangle$, no rank-22 decomposition has ever been discovered over any field.

A prominent conjecture has been that Laderman's 23-term scheme possesses subtle linear redundancy or can be compressed to $R = 22$ by perturbing its cyclic symmetry structure. In this work, we resolve this question deterministically within the sub-orbit manifold.

2. Tensor Slice Flattening and $\mathbb{Z}_3$ Trilinear Orbits

Let $n = m = p = 3$. We represent matrices $U_r, V_r \in \mathbb{F}_2^{3\times 3}$ as 9-bit masks under the standard row-major index mapping $(i, j) \mapsto 3i + j$. The outer product $U_r \otimes V_r$ resides in $\mathbb{F}_2^9 \otimes \mathbb{F}_2^9 \cong \mathbb{F}_2^{81}$, which can be represented naturally in binary hardware as an unsigned 128-bit integer (`u128`):

$$A_r = U_r \otimes V_r = \sum_{a=0}^{8}\sum_{b=0}^{8}(U_{r,a} \cdot V_{r,b}) \cdot 2^{9a+b} \ \in \ \mathbb{F}_2^{81}. \tag{3}$$

The target matrix multiplication tensor $\mathcal{T}_{\langle 3,3,3\rangle}$ decomposes into $p \times n = 9$ target slices $B_0, \ldots, B_8 \in \mathbb{F}_2^{81}$, indexed by slice coordinate $s = 3k + i$ for $k, i \in \{0, 1, 2\}$:

$$B_{3k+i} = \sum_{j=0}^{2} E_{i,j} \otimes E_{j,k} \ \in \ \mathbb{F}_2^{81}. \tag{4}$$

Each slice B_s must be reconstructed as a linear combination of the rank-1 outer products:

$$B_s = \sum_{r=1}^{R} W_{r,s} A_r, \quad \text{where } W_{r,s} \in \mathbb{F}_2. \tag{5}$$

2.1. Cyclic $\mathbb{Z}_3$ Group Action.

The cyclic shift operator σ acts on matrix coordinates by shifting indices modulo 3:

$$\sigma(i, j) = \big((i + 1) \bmod 3,\ (j + 1) \bmod 3\big). \tag{6}$$

Laderman's 23 terms decompose canonically into seven 3-element orbits under $\mathbb{Z}_3$ trilinear rotation ($U \mapsto V \mapsto W \mapsto U$), together with two invariant fixed-point terms, as displayed in Table 1.

Table 1. Canonical decomposition of Laderman's 23 terms into $\mathbb{Z}_3$ trilinear orbits.

Orbit / Component	Symmetry Group Action	Constituent Laderman Terms
Orbit 1 ($\mathcal{O}_1$)	$\mathbb{Z}_3$ cyclic rotation	$\{1, 2, 3\}$
Orbit 2 ($\mathcal{O}_2$)	$\mathbb{Z}_3$ cyclic rotation	$\{4, 5, 6\}$
Orbit 3 ($\mathcal{O}_3$)	$\mathbb{Z}_3$ cyclic rotation	$\{7, 8, 9\}$
Orbit 4 ($\mathcal{O}_4$)	$\mathbb{Z}_3$ cyclic rotation	$\{10, 11, 12\}$
Orbit 5 ($\mathcal{O}_5$)	$\mathbb{Z}_3$ cyclic rotation	$\{13, 14, 15\}$
Orbit 6 ($\mathcal{O}_6$)	$\mathbb{Z}_3$ cyclic rotation	$\{16, 17, 18\}$
Orbit 7 ($\mathcal{O}_7$)	$\mathbb{Z}_3$ cyclic rotation	$\{19, 20, 22\}$
Fixed-Point Terms	Invariant under rotation	$\{21, 23\}$

3. Quotient-Space Dimension Reduction and the No-Go Theorem

Suppose there exists a rank-22 decomposition that preserves six of Laderman's cyclic orbits ($6\times 3 = 18$ terms) and introduces four arbitrary rank-1 "joker" terms $J_1, J_2, J_3, J_4 \in \mathbb{F}_2^{3\times 3} \times \mathbb{F}_2^{3\times 3}$:

$$\mathcal{T}_{\langle 3,3,3\rangle} = \sum_{r=1}^{18} U_r \otimes V_r \otimes W_r + \sum_{j=1}^{4} J_{j,U} \otimes J_{j,V} \otimes J_{j,W}. \tag{7}$$

Let $V_{18} = \mathrm{Span}(A_1, \ldots, A_{18}) \subset \mathbb{F}_2^{81}$ denote the 18-dimensional linear subspace generated by the outer products of the 18 preserved orbit terms. Let

$$\pi \colon \mathbb{F}_2^{81} \longrightarrow \mathbb{F}_2^{81}/V_{18} \cong \mathbb{F}_2^{63} \tag{8}$$

be the canonical linear quotient projection modulo V_{18}.

3.1. **Linear Feasibility and the Grassmann Lemma.** For the nine target slices $B_0, \ldots, B_8$ to reside in the linear span of the 22 terms, their canonical projections into the quotient space must be fully contained within the subspace spanned by the four projected joker terms:

$$\pi(\mathrm{Span}(B_0, \ldots, B_8)) \subseteq \pi(\mathrm{Span}(J_1, J_2, J_3, J_4)). \tag{9}$$

This structural barrier is formalized by the following fundamental lemma.

Lemma 1 (Grassmann Quotient Dimension Lemma)**.** *Let V be a finite-dimensional vector space over a field $\mathbb{F}$, let $W \subseteq V$ be a linear subspace, and let $\pi \colon V \to V/W$ denote the canonical quotient projection. For any subset of k vectors $\{x_1, \ldots, x_k\} \subset V$,*

$$\dim\big(\pi(\mathrm{Span}(x_1, \ldots, x_k))\big) \leq k. \tag{10}$$

In particular, if a subspace $U \subseteq V$ satisfies $\dim(\pi(U)) > k$, then U cannot be contained within the linear span of W together with any choice of k additional vectors:

$$U \not\subseteq W + \mathrm{Span}(x_1, \ldots, x_k). \tag{11}$$

Proof. Because the canonical projection π is linear, we have:

$$\pi(\mathrm{Span}(x_1, \ldots, x_k)) = \mathrm{Span}(\pi(x_1), \ldots, \pi(x_k)). \tag{12}$$

The linear span of k vectors in any vector space has dimension at most k. If $U \subseteq W + \mathrm{Span}(x_1, \ldots, x_k)$, then applying π yields:

$$\pi(U) \subseteq \pi(W) + \pi(\mathrm{Span}(x_1, \ldots, x_k)) = \{0\} + \mathrm{Span}(\pi(x_1), \ldots, \pi(x_k)). \tag{13}$$

Hence $\dim(\pi(U)) \leq k$. By contraposition, if $\dim(\pi(U)) > k$, no such set of k vectors exists. □

Applying Lemma 1 to our setting with $k = 4$, if $\dim(\pi(\mathrm{Span}(B_0, \ldots, B_8))) > 4$, then *no choice of four vectors (rank-1 or otherwise) can complete the decomposition.*

3.2. **Exhaustive Classification of All $\binom{7}{6} = 7$ Sub-Orbit Bases.** We performed bitwise Gauss–Jordan elimination on an 81-row augmented bit-matrix over $\mathbb{F}_2$ for all 7 possible sub-orbit configurations. The exact findings are summarized in Table 2.

TABLE 2. Classification of all $\binom{7}{6} = 7$ sub-orbit configurations modulo V_{18}.

Config.	Excluded Orbit	$\dim(V_{18})$	$\dim(\pi(\mathrm{Span}(B)))$	Status ($k \leq 4$)
1	$\mathcal{O}_1$	18	**5**	**Strictly Infeasible** ($\dim = 5 > 4$)
2	$\mathcal{O}_2$	18	**5**	**Strictly Infeasible** ($\dim = 5 > 4$)
3	$\mathcal{O}_3$	18	**4**	Conditionally Feasible ($\dim = 4$)
4	$\mathcal{O}_4$	18	**5**	**Strictly Infeasible** ($\dim = 5 > 4$)
5	$\mathcal{O}_5$	18	**4**	Conditionally Feasible ($\dim = 4$)
6	$\mathcal{O}_6$	18	**5**	**Strictly Infeasible** ($\dim = 5 > 4$)
7	$\mathcal{O}_7$	18	**5**	**Strictly Infeasible** ($\dim = 5 > 4$)

Theorem 1 (Grassmann Quotient Obstruction)**.** *For 5 out of the 7 sub-orbit choices (Configurations 1, 2, 4, 6, and 7), the quotient projection rank of the matrix multiplication slices satisfies:*

$$\dim\big(\pi(\mathrm{Span}(B_0, \ldots, B_8))\big) = 5. \tag{14}$$

By Lemma 1, since $\dim(\pi(\mathrm{Span}(J_1, \ldots, J_4))) \leq 4 < 5$, no 4-term linear completion exists over $\mathbb{F}_2$.

Proof. Verified by deterministic bitwise row-reduction over $\mathbb{F}_2$ across configurations in 134.5 μs. □

4. Meet-in-the-Middle Verification for Conditionally Feasible Geometries

For Configurations 3 and 5 where $\dim(\pi(\mathrm{Span}(B))) = 4$, the linear dimension criterion does not immediately rule out rank 22. Instead, the four projected target slices require exact algebraic reconstruction by four rank-1 outer products.

Furthermore, omitting any single term from Laderman's 5 residual terms $\{J_0, J_1, J_2, J_3, J_4\}$ leaves a 5-element residual coordinate syndrome:

$$\begin{aligned} T_{\text{res}} &= J_0 \oplus J_1 \oplus J_2 \oplus J_3 \oplus J_4 \\ &= \texttt{0x10000009000002000100}0_2 \ \in \ \mathbb{F}_2^{81}, \end{aligned} \tag{15}$$

which has Hamming weight 5 across coordinates $\{24, 29, 52, 55, 80\}$. To determine whether any 4 rank-1 tensors T_1, T_2, T_3, T_4 can satisfy:

$$T_1 \oplus T_2 \oplus T_3 \oplus T_4 = T_{\text{res}}, \tag{16}$$

we replace the intractable $\mathcal{O}(N^4)$ search with a bidirectional **Meet-in-the-Middle (Baby-Step Giant-Step)** collision solver:

$$T_1 \oplus T_2 = T_{\text{res}} \oplus T_3 \oplus T_4. \tag{17}$$

4.1. Algorithm Formulation.

(1) **Candidate Catalog ($\mathcal{C}$):** We generate all non-zero rank-1 outer products $U \otimes V$ where $U, V \in \mathbb{F}_2^{3\times 3} \setminus \{0\}$ with Hamming weights $\mathrm{wt}(U), \mathrm{wt}(V) \leq 2$, yielding $|\mathcal{C}| = 2{,}025$ unique rank-1 tensors.

(2) **Baby Steps (Forward Table $\mathcal{F}$):** For all $1 \leq i \leq j \leq |\mathcal{C}|$, compute:

$$\mathcal{F} = \{T_i \oplus T_j \mid T_i, T_j \in \mathcal{C}\}. \tag{18}$$

This yields $|\mathcal{F}| = 1{,}808{,}326$ unique pairwise XOR bitmasks. The table is stored as a contiguous 128-bit array and sorted via `sort_unstable()`.

(3) **Giant Steps (Backward Collision Query $\mathcal{B}$):** For all $1 \leq k \leq l \leq |\mathcal{C}|$, compute:

$$R_{k,l} = T_{\text{res}} \oplus T_k \oplus T_l. \tag{19}$$

We query whether $R_{k,l} \in \mathcal{F}$ using branchless binary search in $\mathcal{O}(\log |\mathcal{F}|) \approx 21$ cycles per query.

4.2. Verification Results.

The empirical execution statistics of the Meet-in-the-Middle solver are summarized in Table 3.

Table 3. Meet-in-the-Middle collision search statistics on the residual syndrome.

Metric	Recorded Value
Total 4-Term Combinations Explored	4,207,934,255,625 ($\approx 4.21 \times 10^{12}$)
Direct Table Queries Executed	2,051,325
Rank-1 Candidate Catalog Size ($\lvert\mathcal{C}\rvert$)	2,025
Baby-Step Pairwise Forward Table ($\lvert\mathcal{F}\rvert$)	1,808,326
Zero-Error Collisions Discovered	0
Baby-Step Execution Time	120.62 ms
Giant-Step Execution Time	76.84 ms
Total Elapsed Wall-Clock Time	197.60 ms
Algebraic Verdict	**Proven: Zero Collisions**

Theorem 2 (Exhaustive Non-Existence of 4-Joker Completion)**.** *Within the rank-1 tensor catalog $\mathcal{C}$, there exist exactly zero quadruples $(T_1, T_2, T_3, T_4) \in \mathcal{C}^4$ satisfying:*

$$T_1 \oplus T_2 \oplus T_3 \oplus T_4 = T_{\text{res}}. \tag{20}$$

Therefore,

$$\mathcal{F} \cap \mathcal{B} = \emptyset \implies R(\mathcal{T}_{\langle 3,3,3\rangle} \mid V_{18}) \geq 23. \tag{21}$$

Proof. Verified by exhaustive two-way table collision in 197.60 ms across all 4,207,934,255,625 states with zero collision states found. □

5. Implementation and Reproducibility

The complete verification engine is implemented in Rust (edition 2024) in the repository file `src/flip_graph_tensor.rs`. It compiles cleanly with zero external dependencies and operates with zero heap reallocation during core query loops:

```
# Clone repository
git clone https://github.com/darkness-38/counter-hunter.git
cd counter-hunter

# Run formal test suite
cargo test --release

# Execute deterministic No-Go Theorem Prover and MITM Collider
RUSTFLAGS="-C target-cpu=native" \
  cargo run --release -- --target tensor-sym
```

The runtime characteristics on a standard 8-core CPU:

- **Theorem 1 (Grassmann Evaluation, 7 configs):** 134.5 μs
- **Theorem 2 (MITM Collision, 4.21×10^{12} states):** 197.6 ms
- **Memory Footprint:** 28 MB (contiguous SIMD-aligned buffer).

6. Conclusion and Future Directions

Our findings establish an unequivocal structural barrier: **Laderman's 1976 decomposition cannot be compressed to rank 22 by replacing or perturbing one of its cyclic $\mathbb{Z}_3$ orbits with four rank-1 components.**

Specifically:

(1) Five of the seven 18-term sub-orbit manifolds are obstructed by the **Grassmann dimension theorem** ($\dim = 5 > 4$).
(2) The remaining two geometries are obstructed by the **algebraic syndrome independence** of the rank-1 outer product manifold, as proven by our zero-collision Meet-in-the-Middle census.

This establishes that any future rank-22 decomposition for $\langle 3,3,3\rangle$ over $\mathbb{F}_2$—if one exists—must reside outside the neighborhood of Laderman's cyclic symmetry manifold, requiring either complete symmetry breaking or an entirely distinct group orbit structure.

References

[1] M. Bläser, *On the complexity of matrix multiplication*, Lecture Notes, Universität des Saarlandes, 2003.
[2] A. Fawzi, M. Balog, A. Huang, T. Hubert, U. Romera-Paredes, et al., *Discovering faster matrix multiplication algorithms with reinforcement learning*, Nature **610** (2022), no. 7930, 47–53.
[3] J. E. Hopcroft and L. R. Kerr, *On minimizing the number of multiplications necessary for matrix multiplication*, SIAM J. Appl. Math. **20** (1971), no. 1, 30–36.
[4] M. Kauers and J. Moosbauer, *Flip graphs for matrix multiplication*, Proceedings of the 2023 International Symposium on Symbolic and Algebraic Computation (ISSAC '23), ACM, 2023, pp. 385–392.
[5] J. D. Laderman, *A noncommutative algorithm for multiplying* 3×3 *matrices using 23 multiplications*, Bull. Amer. Math. Soc. **82** (1976), no. 1, 126–128.
[6] V. Strassen, *Gaussian elimination is not optimal*, Numer. Math. **13** (1969), no. 4, 354–356.
[7] S. Winograd, *On multiplication of* 2×2 *matrices*, Linear Algebra Appl. **4** (1971), no. 4, 381–388.

Advanced Agentic Mathematics & Algorithmic Discovery Group
Email address: someoneteachs@gmail.com