

Harmonic maps of Riemannian manifolds

Tien NGUYEN MANH

Feb 20, 2018

Contents

| | | |
|----------|---|-----------|
| 1 | Harmonic maps | 1 |
| 1.1 | Variational approach: energy integral and tension field | 1 |
| 1.2 | Formulation using connection on vector bundle | 4 |
| 1.3 | The case of $E = f^*TM'$ | 6 |
| 1.3.1 | Energy functional and tension field | 6 |
| 1.3.2 | Fundamental form, some results in case of signed curvature | 7 |
| 1.4 | Example: Riemannian immersion | 8 |
| 1.4.1 | Second fundamental form. | 8 |
| 1.4.2 | The case of signed curvature. | 9 |
| 1.5 | Composition of maps | 9 |
| 2 | Nonlinear heat flow: Global equation and existence of harmonic maps. | 10 |
| 2.1 | Statement of the main results. | 10 |
| 2.2 | Strategy of the proof. | 11 |
| 2.3 | Global equation and Uniqueness of nonlinear heat equation. | 12 |
| 3 | A few energy estimates. | 14 |
| 3.1 | Estimate of density energies | 14 |
| 3.2 | Estimate of total energies | 16 |

1 Harmonic maps

1.1 Variational approach: energy integral and tension field

Notation. Let M, M', M'' be Riemannian manifolds of dimension n, n' and n'' respectively. We will use indices $i, j, k, \dots, \alpha, \beta, \gamma, \dots, a, b, c$ to denote local coordinates of M, M', M'' . Let $f : M \rightarrow M', f' : M' \rightarrow M''$ be a smooth maps, one denotes

$$f_i^\alpha = \frac{\partial f^\alpha}{\partial x^i}, \quad f_{ij}^\alpha = \frac{\partial^2 f^\alpha}{\partial x^i \partial x^j} - \Gamma_{ij}^k f_k^\alpha$$

so that $\nabla h = h_i dx^i$ and $\nabla(\nabla h) = h_{ij} dx^i \otimes dx^j$ and $-\Delta h = \text{Tr} \nabla(\nabla h) = g^{ij} h_{ij}$ for any smooth function h .

Definition 1. The *energy desity* of f at $p \in m$ is defined by

$$e(f)(p) = \frac{1}{2} \langle g, f^* g \rangle_p = \frac{1}{2} g^{ij} f_i^\alpha f_j^\beta g'_{\alpha\beta}$$

and the *energy functional* of f is

$$E(f) = \int_M e(f) dV = \frac{1}{2} \int_M g^{ij} f_i^\alpha f_j^\beta g'_{\alpha\beta} |\det(g_{ij})|^{\frac{1}{2}} dx^1 \wedge \dots \wedge dx^n$$

We recall that the inner product between 2 tensors of type (p, q) $S = S_{j_1 \dots j_q}^{i_1 \dots i_p}$, $T = T_{l_1 \dots l_q}^{k_1 \dots k_p}$ is $\prod_{m,n} g_{i_m k_m} g^{j_n l_n} S_{j_1 \dots j_q}^{i_1 \dots i_p} T_{l_1 \dots l_q}^{k_1 \dots k_p}$

Remark 2. The energy density is non-negative at every point. And $E(f) = 0$ if and only if $e(f) = 0$ at all points if and only if f is constant.

Definition 3. Let σ be a symmetric function of n variables and α be a symmetric $(0,2)$ tensor field, one can define the σ -*energy desity* of α at $P \in M$ to be $\sigma(\beta_1, \dots, \beta_n)(P)$ where β_i are eigenvalues of the linear operator $(g^{ik} \alpha_{ij})_{k,j}$. The σ -*energy* of α is $I_\sigma(\alpha) := \int_M \sigma(\alpha) dV$

Take $\alpha = f^* g'$, one calls $\sigma(\alpha)$ the σ -*energy density* of f and $I_\sigma(\alpha)$ the σ -*energy* of f .

Example 4. The energy functional $E(f)$ is $I_{\sigma_1}(f)$. $V(f) := I_{\sigma_{1/2}}(f)$ is called the *volume* of f .

Lemma 5 (variation of the energy). Let $f_t : M \rightarrow M'$ be a smooth family of smooth maps between Riemannian manifolds for $t \in (t_0, t_1)$. Then

$$\frac{d}{dt} E(f_t) = - \int_M \left(-\Delta f_t^\gamma + g^{ij} \Gamma_{\alpha\beta}^{\gamma} f_{t,i}^\alpha f_{t,j}^\beta \right) g'_{\gamma\nu} \frac{\partial f_t^\nu}{\partial t} dV, \quad \forall t \in (t_0, t_1)$$

Proof. One has

$$\begin{aligned} \frac{dE}{dt}(f_t) &= \frac{1}{2} \int \left[2g^{ij} f_i^\alpha \frac{\partial^2 f_t^\beta}{\partial x^j \partial t} g'_{\alpha\beta} + g^{ij} f_i^\alpha f_j^\beta \frac{\partial g'_{\alpha\beta}}{\partial y^\nu} \frac{df_t^\nu}{dt} \right] dV(g) \\ &= \frac{1}{2} \int \left[- \left(2g^{ij} f_i^\alpha g'_{\alpha\beta} \right)_j \frac{df_t^\beta}{dt} + g^{ij} f_i^\alpha f_j^\beta \frac{\partial g'_{\alpha\beta}}{\partial y^\nu} \frac{df_t^\nu}{dt} \right] dV(g) \end{aligned}$$

The first term is

$$\begin{aligned} - \left(2g^{ij} f_i^\alpha g'_{\alpha\beta} \right)_j \frac{df_t^\beta}{dt} &= -2g^{ij} f_{ij}^\alpha \frac{df_t^\beta}{dt} g'_{\alpha\beta} - 2g^{ij} f_i^\alpha \frac{df_t^\beta}{dt} \frac{\partial g'_{\alpha\beta}}{\partial y^\nu} f_j^\nu \\ &= 2\Delta f^\alpha g'_{\alpha\beta} \frac{df_t^\beta}{dt} - 2g^{ij} f_i^\alpha f_j^\beta \frac{\partial g'_{\alpha\beta}}{\partial y^\nu} \frac{df_t^\nu}{dt} \end{aligned}$$

It remains to check that

$$-2\frac{\partial g'_{\alpha\nu}}{\partial y^\beta} + \frac{\partial g'_{\alpha\beta}}{\partial y^\nu} = -2\Gamma_{\alpha\beta}^{\gamma'} g'_{\gamma\nu}$$

when we are allowed to permute α, β , which is routine. \square

Definition 6. 1. A **vector field along** $f : M \longrightarrow M'$ is a smooth application $v : M \longrightarrow TM'$ such that $\pi \circ v = f$ where $\pi : TM' \longrightarrow M'$ is the canonical projection. In other words, it is the association of each point $P \in M$ a tangent vector at $f(P)$

2. The **tension field** of f is the vector field along f defined by

$$\tau(f)^\gamma := -\Delta f^\gamma + g^{ij} \Gamma_{\alpha\beta}^{\gamma'} f_i^\alpha f_j^\beta$$

By the Lemma 5, $\tau(f)$ is the unique vector field along f such that $\frac{d}{dt} E(f_t) = -\int_M \langle \tau(f), \frac{df_t}{dt} \rangle$. In particular, if f_t is the variation of f along a vector field v along f , i.e. $f_t(P) = \exp_{f(P)}(tv(P))$ then $\frac{d}{dt} E(f_t) = -\langle \tau(f), v \rangle$.

3. $f : M \longrightarrow M'$ is called **harmonic** if $\tau(f) = 0$, or equivalently if f is a critical point of E .

In normal coordinates of M at P and M' at $f(P)$, the tension field of f is given by

$$\tau^\gamma(f)(P) = \sum_i \frac{\partial^2 f^\gamma}{\partial (x^i)^2}(P)$$

Remark 7. 1. If M' is flat, i.e. $R'_{\alpha\beta\gamma\delta} = 0$ then $\tau(f)^\gamma = -\Delta f^\gamma$ is linear in f . We refine the definition of harmonic function.

2. Since $\tau(f)$ depends locally on f , isometries and covering maps are harmonic.

Proposition 8 (Holomorphicity implies harmonicity). *Holomorphic maps between Kahler manifolds are harmonic.*

Proof. We recall that exponential function $\exp_P : T_P M \longrightarrow M'$ on a Kahler manifold M is holomorphic for any $P \in M$. In fact, let $v \in T_P M$ and $\delta v \in T_v(T_P M)$ be a tangent vector at v and denote abusively by J the complex structure of the complex vector space $T_P M$ and that of M , one needs to see that

$$D \exp_P(v) \cdot J \delta v = J(\exp_P(v)) D \exp_P(v) \cdot \delta v \quad (1)$$

In fact, let Y_1, Y_2 be Jacobi fields along $U(t) = \exp_P(tv)$ the geodesics of M starting at P in direction v with $Y_1(0) = Y_2(0) = 0, \dot{Y}_1(0) = \delta v, \dot{Y}_2(0) = J \delta v$ then the LHS of (1) is $Y_2(1)$, and the RHS is $J(U(1))Y_1(1)$. Then one can see that $Y_2(t) - J(U(t))Y_1(t) = 0$ for every $t \in [0, 1]$ since it is true at $t = 0$ and the derivative with respect to t vanishes as $\nabla_U J = 0$.

Therefore, at a point P of a Kahler manifold M , there exist holomorphic coordinates $z^j = x^j + iy^j$ of M in a neighborhood of P such that $\{x_j, y_j : j = \overline{1, n/2}\}$ are normal coordinates centered in P . Using such coordinates for $P \in M$ and $f(P) \in M'$, one has $\Delta f^\gamma = 0$ since f^γ is holomorphic and $\Gamma_{\alpha\beta}^{\gamma'}(P) = 0$ by normality, it follows that $\tau(f) = 0$ at every point $P \in M$. \square

1.2 Formulation using connection on vector bundle

Setup and notation. Let E be a metric vector bundle over a Riemannian manifold M , i.e. each fiber of E is equipped with an inner product that we denote by $(g'_{\alpha\beta})$. The metric of M is denoted by (g_{ij}) . Let n and m be the dimension of M of the fiber.

Covariant derivatives and exterior derivatives. We recall that a **covariant derivative** or a **connection** $\tilde{\nabla}$ of E is uniquely determined in local coordinates by an $m \times m$ matrix A of 1-forms, in other words, it is an 1-form on M with value in $\text{Hom}_M(E, E)$ which depends on the local frame of E (i.e. A is not a tensor with value in E). A is called the **connection form** of $\tilde{\nabla}$. Locally

$$\tilde{\nabla}_X(s^\alpha \tilde{e}_\alpha) = (\nabla_X s^\alpha) \tilde{e}_\alpha + A_\beta^\alpha(X) s^\beta \tilde{e}_\alpha.$$

When one prefers to work with forms rather than tensors with value in E , one uses an **exterior derivative**, a map $D : A^p(M, E) \longrightarrow A^{p+1}(M, E)$ which turns an p -form with value in E to an $p + 1$ -form with value in E . Locally

$$D(s^\alpha \tilde{e}_\alpha) = (ds^\alpha) \tilde{e}_\alpha + A_\beta^\alpha \wedge s^\beta \tilde{e}_\alpha.$$

and

$$D^2(s^\alpha \tilde{e}_\alpha) = (dA + A \wedge A) \wedge s.$$

One notes $\Theta := dA + A \wedge A$, which is an $m \times m$ matrix of 2-forms of M . Unlike A , Θ , seen as an 2-form with value in $\text{Hom}_M(E, E)$ does not depend on the local frame of E , i.e. Θ transforms as a $(0,2)$ tensor with value in E , called the **curvature form**.

The fibrewise metric structure of E and the metric tensor of M give rise to a pointwise inner product of (p, q) tensors of M with value in E , in particular a pointwise inner product $(s, s') \mapsto s \cdot s'$ from $A^p(M, E) \times A^p(M, E)$ to $C^\infty(M)$. Integrated over M , the pointwise inner product gives rise to a global inner product $\int_M \langle \cdot, \cdot \rangle$ of $A^p(M, E)$. One denotes by $\delta : A^{p+1}(M, E) \longrightarrow A^p(M, E)$ the adjoint operator of $D : A^p(M, E) \longrightarrow A^{p+1}(M, E)$ with respect to this inner product, i.e. $\int_M \langle Ds, s' \rangle_{A^{p+1}(M, E)} = \int_M \langle s, \delta s' \rangle_{A^p(M, E)}$ for all $s \in A^p(M, E), s' \in A^{p+1}(M, E)$.

Laplacian operator and harmonic forms. The **Hodge Laplacian** is defined as a endomorphism of $A^p(M, E)$ given by

$$\tilde{\Delta} = D\delta + \delta D$$

and a form $s \in A^p(M, E)$ is called **harmonic** if $\tilde{\Delta}s = 0$. Since the Laplacian operator represents the *Dirichlet integral*, i.e.

$$\int_M \langle Ds, Ds' \rangle + \int_M \langle \delta s, \delta s' \rangle = \int_M \langle \tilde{\Delta}s, s' \rangle,$$

one has $\tilde{\Delta}s = 0$ if and only if $Ds = \delta s = 0$.

Riemannian connected bundle. The metric vector bundle E over M is called a **Riemannian-connected bundle** if it is equipped with a connection $\tilde{\nabla}$ under which the metric g' of E is parallel, i.e. $\tilde{\nabla}g' = 0$, in other words, the matrix A in an orthonormal frame is anti-symmetric: $A + {}^tA = 0$. Unless explicitly indicated, we always suppose that our metric vector bundle E is Riemannian-connected and the metric g' is parallel to the connection being used.

Example 9. The case of our interest is when we have a smooth map $f : M \longrightarrow M'$ and $E = f^*TM'$ is a metric vector bundle over M under the metric g' induced from M' . Taking the connection $\tilde{\nabla}$ to be the Levi-Civita connection ∇' on M' , meaning

$$\tilde{\nabla}_X s = \nabla'_{f_*X} s,$$

for any vector field s along f , one can see that E is a Riemannian-connected bundle over M .

Lemma 10. Let E be a Riemannian-connected bundle and $s = s_i^\alpha dx^i \tilde{e}_\alpha \in A^1(M, E)$, one has

1. $\delta s = (\delta s)^\alpha \tilde{e}_\alpha \in A^0(M, E)$ where

$$(\delta s)^\alpha = -g^{ij} \left(\nabla_i s_j^\alpha + A_{\beta i}^\alpha s_j^\beta \right),$$

2. $\Delta s = (\Delta s)_i dx^i$ where $(\Delta s)_i$ is an $m \times m$ matrix given by

$$(\Delta s)_i = -\tilde{\nabla}^k \tilde{\nabla}_k s_i + {}^t \left(\Theta_i^h - \text{Ric}_i^h \right) s_h$$

where:

- the indices i, h, k correspond to local coordinates of M ,
 - Θ_i^h is the curvature form of $\tilde{\nabla}$ with its indices raised by the metric g of M ,
 - $\text{Ric}_i^h = \text{Ric}_i^h I_m$ is the Ricci curvature tensor of (M, g) with indices raised by the metric g , multiplied by the identity $m \times m$ matrix,
 - $\tilde{\nabla}^k = g^{hk} \tilde{\nabla}_h$.
3. With $s \cdot s'$ denoting the pointwise inner product of $A^1(M, E)$ and $\langle \cdot, \cdot \rangle_E$ denoting the metric g' of E , one has

$$-\frac{1}{2} \Delta(s \cdot s) = -s \cdot \Delta s + \langle \tilde{\nabla}_i s_k, \tilde{\nabla}^i s^k \rangle_E - \left\langle {}^t \left(\Theta_i^h - \text{Ric}_i^h \right) s_h, s^i \right\rangle_E \quad (2)$$

where the superscript i, h are raised by the metric g .

Proof. Computational in nature. □

1.3 The case of $E = f^*TM'$

1.3.1 Energy functional and tension field

Our interest will be the case of Example 9 where $E = f^*TM'$ for a smooth map $f : M \longrightarrow M'$ of Riemannian manifolds is a Riemannian-connected bundle over M with the connection $\tilde{\nabla}$ given by the Levi-Civita connection of M' .

In this section, the tangent map $Tf : TM \longrightarrow TM'$ can be interpreted as a form f_* in $A^1(M, E)$. The energy functional can be rewritten as

$$E(f) = \frac{1}{2} \int_M f_i^\alpha f_j^\beta g'^{ij} g'_{\alpha\beta} dV = \frac{1}{2} \langle f_*, f_* \rangle_{A^1(M, E)}.$$

Proposition 11. *Let $f : M \longrightarrow M'$ and $E = f^*TM'$ be the Riemannian-connected bundle over M . Then:*

1. $A_\alpha^\beta = \Gamma_{\gamma\alpha}^{\prime\beta} f_i^\gamma dx^i$ where $\Gamma_{\gamma\alpha}^{\prime\beta}$ are Christoffel symbols of (M', g') .
2. $Df_* = 0$ where f_* is considered as an element of $A^1(M, E)$. Hence $\tilde{\Delta}f_* = D\delta f_*$.
3. The tension field of f is $\tau(f) = -\delta f_*$.

Proof. 1. One has

$$A_{\alpha i}^\beta \tilde{e}_\beta = \tilde{\nabla}_i \tilde{e}_\alpha = \tilde{\nabla}_{f_i^\gamma \tilde{e}_\gamma} \tilde{e}_\alpha = f_i^\gamma \Gamma_{\gamma\alpha}^{\prime\beta} \tilde{e}_\beta,$$

Therefore $A_\alpha^\beta = f_i^\gamma \Gamma_{\gamma\alpha}^{\prime\beta} dx^i$.

2. By direct computation:

$$Df_* = \left(\frac{\partial^2 f^\alpha}{\partial x^i \partial x^j} + \Gamma_{\gamma\beta}^{\prime\alpha} f_i^\gamma f_j^\beta \right) dx^j \wedge dx^i \otimes \tilde{e}_\alpha = 0$$

since it is the product of a symmetric quantity in (i, j) and an anti-symmetric one.

3. Using the first part of Lemma 10 for $s = f_* = f_i^\alpha dx^i \otimes \tilde{e}_\alpha$, one has $\delta f_* = -g^{ij} \left(\nabla_i \nabla_j f^\gamma + \Gamma_{\alpha\beta}^{\prime\gamma} f_i^\alpha f_j^\beta \right) \tilde{e}_\gamma = -\tau(f)$

□

It follows immediately that

Corollary 12. *$f : M \longrightarrow M'$ is a harmonic map of compact Riemannian manifolds if and only if f_* is harmonic as form in $A^1(M, f^*TM')$.*

1.3.2 Fundamental form, some results in case of signed curvature

Definition 13. The *fundamental form* of a map $f : M \longrightarrow M'$ of Riemannian manifolds is the $(0,2)$ symmetric tensor on M with value in $E = f^*TM'$ defined by

$$\beta(f) := \tilde{\nabla} f_* = \left(f_{ij}^\gamma + \Gamma_{\alpha\beta}^{\prime\gamma} f_i^\alpha f_j^\beta \right) dx^i \otimes dx^j \otimes \tilde{e}_\gamma.$$

The function f is called **totally geodesic** if $\beta(f) = 0$ identically on M .

Remark 14. 1. The tension field $\tau(f) = g^{ij}\beta(f)_{ij}$ is the trace of the fundamental form.

2. If f is totally geodesic then it is harmonic.

When $s = f_*$, Lemma 10 and Remark ?? become Lemma 15, with no more than direct computation. The appearance of Riemann curvature tensor R' of (M', g') is due to the formula

$$R'^\rho_{\sigma\mu\nu} = \partial_\mu \Gamma'^\rho_{\nu\sigma} - \partial_\nu \Gamma'^\rho_{\mu\sigma} + \Gamma'^\rho_{\mu\lambda} \Gamma'^\lambda_{\nu\sigma} - \Gamma'^\rho_{\nu\lambda} \Gamma'^\lambda_{\mu\sigma}.$$

Lemma 15. 1. $Q(f_*)$ is given by

$$Q(f_*) = R'_{\alpha\beta\gamma\delta} f_i^\alpha f_j^\beta f_k^\gamma f_l^\delta g^{ik} g^{jl} - \text{Ric}^{ij} f_i^\alpha f_j^\beta g'_{\alpha\beta}$$

and

$$Q(f_*)^{ij}_{\alpha\beta} = R'_{\alpha\beta\gamma\delta} f_k^\gamma f_l^\delta g^{ik} g^{jl} - \text{Ric}^{ij} g'_{\alpha\beta}.$$

2. If f is harmonic then

$$-\Delta e(f) = |\beta(f)|^2 - R'_{\alpha\beta\gamma\delta} f_i^\alpha f_j^\beta f_k^\gamma f_l^\delta g^{ik} g^{jl} + \text{Ric}^{ij} f_i^\alpha f_j^\beta g'_{\alpha\beta}$$

where $|\beta(f)|$ is the pointwise norm of $\beta(f)$.

The previous computation of $Q(f_*)$ in term of Riemannian curvature of M' and Ricci curvature of M give the following result in case the curvature of M and M' are of definite sign.

Notation. Given a Riemannian manifold M , we will use the following notation:

1. $\text{Ric} \geq 0$ (resp. $\text{Ric} > 0$) if the Ricci curvature is positive semi-definite (resp. positive definite) as symmetric bilinear form.
2. $\text{Riem} \leq 0$ (resp. $\text{Riem} < 0$) if all sectional curvatures are negative (resp. strictly negative), i.e. $R_{ijhk} u^i v^j u^h v^k \leq 0$ (resp. $R_{ijhk} u^i v^j u^h v^k < 0$) for non-colinear vectors u, v .

Corollary 16. Let $f : M \longrightarrow M'$ be a map of Riemannian manifolds.

1. If f is harmonic and $Q(f_*) \leq 0$ then f is totally geodesic and $e(f)$ is constant.

2. If $\text{Ric}(M) \geq 0$ and $\text{Riem}(M') \leq 0$ then f is harmonic if and only if f is totally geodesic.

Proof. All the statements are consequence of 2) of Lemma 15 and the fact that $\int_M \Delta e(f) dV = 0$, noticing that

- $\text{Ric}^{ij} f_i^\alpha f_j^\beta g'_{\alpha\beta}$ is $\text{Ric} \otimes g'$ applied doubly to $f_i^\alpha dx^i \otimes \tilde{e}_\alpha$.
- $R'_{\alpha\beta\gamma\delta} f_i^\alpha f_j^\beta f_k^\gamma f_l^\delta g^{ik} g^{jl}$ is $(f^* R')_{ijkl} g^{ik} g^{jl}$. In a normal coordinate at P where $g^{ik} = \delta_{ik}$, $g^{jl} = \delta_{jl}$, it is the sum of sectional curvatures of tangent planes formed by $f_* e_i, f_* e_j$, and therefore negative.

□

1.4 Example: Riemannian immersion

Let $f : M \rightarrow M'$ be a Riemannian immersion, i.e. Tf is injective and $f^* g' = g$. We will see that the fundamental form $\beta(f)$ that we defined earlier is the same as usual definition in courses of Riemannian geometry.

1.4.1 Second fundamental form.

One defines the symmetric (0,2)-tensor Π of $f^* TM'$ as the unique normal vector of M such that

$$\langle \Pi_{ij}, \xi_\sigma \rangle := -\langle \tilde{\nabla}_i \xi_\sigma, f_* e_j \rangle$$

for every vector field ξ_σ of M' orthogonal to M .

Lemma 17 (Second fundamental form). *If f is a Riemannian immersion then $\beta(f)_{ij} = -\Pi_{ij}$ and they are orthogonal to M . In particular, if f is totally geodesic then it maps geodesics of M to geodesics of M'*

Proof. One has

$$\begin{aligned} \langle \tilde{\nabla}_i \xi_\sigma, f_* e_j \rangle &= \langle \xi_\sigma, \tilde{\nabla}_i (f_* e_j) \rangle = \langle \xi_\sigma, \tilde{\nabla}_i (f_l^\gamma dx^l \otimes \tilde{e}_\gamma) e_j + f_* \nabla_i e_j \rangle \\ &= \langle \xi_\sigma, (f_{il}^\gamma dx^l \tilde{e}_\gamma + f_l^\gamma dx^l \tilde{\nabla}_i \tilde{e}_\gamma) e_j \rangle \\ &= \langle \xi_\sigma, f_{ij}^\gamma \tilde{e}_\gamma + f_j^\gamma A_{\gamma i}^\alpha \tilde{e}_\alpha \rangle = \left\langle \xi_\sigma, \left(f_{ij}^\gamma + \Gamma_{\alpha\beta}^{\gamma} f_i^\alpha f_j^\beta \right) \tilde{e}_\gamma \right\rangle \\ &= \langle \xi_\sigma, \tilde{\nabla}_i (f_*) . e_j \rangle = \langle \xi_\sigma, \beta(f)_{ij} \rangle \end{aligned}$$

where we used $\xi_\sigma \perp f_* e_j$ in the first line and $\xi_\sigma \perp f_*([e_i, e_j])$ in the second line. Hence $\Pi_{ij} \equiv -\beta(f)_{ij}$ modulo an element in TM . It remains to see that $\beta(f)_{ij} \perp M$ in order to conclude $\Pi = -\beta(f)$. By definition, one has $\beta(f)_{ij} = \tilde{\nabla}_i (f_*) . e_j$ and

$$\begin{aligned} \langle \beta(f)_{ij}, f_* e_k \rangle &= \langle \tilde{\nabla}_i (f_*) . e_j, f_* e_k \rangle = \tilde{\nabla}_i \langle f_* e_j, f_* e_k \rangle - \langle \nabla_i e_j, e_k \rangle - \langle f_* e_j, \tilde{\nabla}_i (f_* e_k) \rangle \\ &= \nabla_i \langle e_j, e_k \rangle - \langle \nabla_i e_j, e_k \rangle - \langle \beta(f)_{ik}, f_* e_j \rangle - \langle e_j, \nabla_i e_k \rangle \\ &= -\langle \beta(f)_{ik}, f_* e_j \rangle \end{aligned}$$

Then using the symmetric of $\beta(f)_{ij}$, one has $\langle \beta(f)_{ij}, f_* e_k \rangle = 0$.

Finally, if $\beta(f) = 0$ and X is a geodesic vector field of M , one needs to prove that $f_* X$ is a geodesic vector field of M' . In fact

$$\tilde{\nabla}_X(f_* X) = (\tilde{\nabla}_X f_*)X + f_* \nabla_X X = \beta(f)(X, X) = 0.$$

Hence $f_* X$ is a geodesic field of M' . \square

Example 18. The inclusion $x \mapsto (x, y_0)$ of a Riemannian manifold M to the Riemannian product $M \times N$ is totally geodesic.

Definition 19. Given an orthonormal frame $(\xi_\sigma)_{1 \leq \sigma \leq n'-n}$, the *mean normal curvature field* of M in M' at $P \in M$ is defined as

$$\xi(P) := \sum_{\sigma=1}^{n'-n} g^{ij} \langle \Pi_{ij}, \xi_\sigma \rangle \xi_\sigma = - \sum_{\sigma=1}^{n'-n} \langle \tau(f), \xi_\sigma \rangle \xi_\sigma.$$

The immersion f is said to be *minimal* if ξ vanishes identically on M .

Remark 20. 1. Since $(\xi_\sigma)_{1 \leq \sigma \leq n'-n}$ is an orthonormal frame, one also has

$$\xi(P) = -g^{ij} \langle \tilde{\nabla}_i \xi_\sigma, f_* e_j \rangle \xi_\sigma(P) = - \sum_{\sigma=1}^{n'-n} \operatorname{div}(\xi_\sigma(P)) \xi_\sigma(P)$$

2. The mean normal curvature field is the tension field of f , i.e. $\xi = -\tau(f)$. Minimal immersions are exactly harmonic immersion.

1.4.2 The case of signed curvature.

If $f : M \rightarrow M'$ is a Riemannian immersion then the Ricci term of Lemma 15 is actually the scalar curvature of M , one has

Proposition 21. Let $f : M \rightarrow M'$ be a Riemannian immersion. Suppose that $\operatorname{Riem}(M') \leq 0$ and $r = g^{ij} \operatorname{Ric}_{ij} < 0$ at one point of M . If f is harmonic then it is constant.

1.5 Composition of maps

The following results come from direct computation of the second fundamental form and tension field of composition of maps between Riemannian manifolds. Again, we use indices i, j, k, \dots for M , $\alpha, \beta, \gamma, \dots$ for M' and a, b, c, \dots for M'' .

Proposition 22. Let $f : M \rightarrow M'$ and $f' : M' \rightarrow M''$ be smooth maps of Riemannian manifolds, then

$$\beta(f' \circ f)_{ij}^a = \beta(f)_{ij}^\gamma f_\gamma'^a + \beta(f')_{\alpha\beta}^a f_i^\alpha f_j^\beta \quad (3)$$

and

$$\tau(f' \circ f)^a = \tau(f)^\gamma f_\gamma'^a + g^{ij} \beta(f')_{\alpha\beta}^a f_i^\alpha f_j^\beta \quad (4)$$

Therefore,

| | | |
|------------------------------|------------------------------|--|
| <i>If f' is</i> | <i>and f is</i> | <i>then $f' \circ f$ is</i> |
| <i>totally geodesic</i> | <i>totally geodesic</i> | <i>totally geodesic</i> |
| <i>totally geodesic</i> | <i>harmonic</i> | <i>harmonic</i> |

and the inverse of a totally geodesic map is totally geodesic.

Remark 23. 1. The Proposition says that SFF of composition follows chain rule: the SFF of $f' \circ f$ is a 2-tensor on M valued in TM'' is obtained by pushing forward the SFF of f and pulling back the SFF of f' and then adding the two. In Cartan formalism:

$$(f' \circ f)_{\alpha ij} = (f'_{\alpha k}(f) f_{ki})_j = f'_{\alpha kh} f_{hj} f_{ki} + f'_{\alpha k} f_{kij} = f^* \beta(f') + f'_* \beta(f)$$

2. It is not true in general that the composition of harmonic maps are harmonic. For example, if one composes the harmonic maps $\mathbb{R} \rightarrow \mathbb{R}^2 : x \mapsto (x, 2x)$ and $\mathbb{R}^2 \rightarrow \mathbb{R} : (x, y) \mapsto x^2 - y^2$, the result is $\mathbb{R} \rightarrow \mathbb{R} : x \mapsto -3x^2$, which is not harmonic.

Proposition 24 (composition with immersion). *If $f' : M' \rightarrow M''$ is a Riemannian immersion and $f : M \rightarrow M'$ then*

1. *Energy functionals: $E(f) = E(f' \circ f)$.*
2. *Tension fields: $\tau(f)$ is the projection of $\tau(f' \circ f)$ to M' .*

Proof. 1. One has $e(f) = \frac{1}{2} \langle g, f^* g' \rangle = \frac{1}{2} \langle g, (f' \circ f)^* g'' \rangle = e(f' \circ f)$.

2. One has $\tau(f' \circ f)^a = \tau(f)^a + g^{ij} \beta(f')^a_{\alpha\beta} f_i^\alpha f_j^\beta$ by (4). The conclusion follows since the second term is normal to M' .

□

The following immediate corollary of Proposition 24 is a generalization of the fact that a curve on a surface M' of \mathbb{R}^3 is geodesic if and only if its curvature vector is orthogonal to M' .

Corollary 25. *If $f' : M' \rightarrow M''$ is a Riemannian immersion, then a map $f : M \rightarrow M'$ is harmonic if and only if $\tau(f' \circ f) \perp M'$.*

2 Nonlinear heat flow: Global equation and existence of harmonic maps.

2.1 Statement of the main results.

We want to prove in the next part existence of harmonic map between manifolds M and M' by deforming any map $f : M \rightarrow M'$ using the τ -flow, meaning solving the PDE:

$$\begin{cases} \frac{df_t}{dt} = \tau(f_t), & t \in [\alpha, \omega] \\ f_\alpha = f, \end{cases} \quad (5)$$

The equation makes sense because both $\frac{df_t}{dt}$ and $\tau(f_t)$ are vector fields along f_t . Since this is the gradient-descent equation for E , the energy of f_t decreases and we hope, under conditions, to obtain convergence of $\{f_t\}$ to a critical point f_∞ of E , this will prove that any homotopy class of $C^\infty(M, M')$ has at least a harmonic map.

It is proved by Eells and Sampson [?] that

Theorem 26 (Eells-Sampson). *Let M and M' be compact Riemannian manifolds with $\text{Riem}(M') \leq 0$ then there exists a harmonic map $f : M \rightarrow M'$ in each homotopy class.*

Several boundary conditions, of Dirichlet, Neumann or mixed type, are also taken into account by Hamilton [?], as an example, we will state the Dirichlet problem:

Theorem 27 (Hamilton). *Let M and M' be compact Riemannian manifolds possibly with boundary. Suppose that M' has $\text{Riem}(M') \leq 0$ and $\partial M'$ is convex, then any relative homotopy class of $C^\infty(M, M')$ has a harmonic element.*

About the terminology, **relative homotopy class** means that we only deform f among maps with the same value on ∂M . The **convexity of $\partial M'$** means that the geodesic at any point in $\partial M'$ with initial tangent vector parallel to the boundary does not enter the interior of M' in short time. This condition can be expressed using the Christoffel symbols of M' at the point in question: If M' is coordinated by y^1, \dots, y^n with $M' = \{y^n \geq 0\}$, then the convexity is translated as $\Gamma_{\alpha\beta}^n \geq 0$ as a symmetric form ($1 \leq \alpha, \beta \leq n-1$). This can be seen by the geometric interpretation of the second fundamental form of the embedding $s : \partial M' \hookrightarrow M'$, which is $\text{II}(s) = -\Gamma_{\alpha\beta}^n$.

It is easy to see that the convexity of $\partial M'$ is a necessary condition, as harmonic maps from \mathbb{R} are geodesics: Suppose the condition does not hold at $x \in \partial M'$, meaning that upto time t the geodesic flow of M' initially pointing into M' remains in the interior. The geodesic of $\partial M'$ of length less than t with the same initial tangent therefore cannot be deformed into a geodesic of M' in relative homotopy class.

2.2 Strategy of the proof.

In order to have a global frame, we will embed M' into an Euclidean space V , but we will not use the Euclidean metric of V . In fact, let T be a tubular neighborhood of M' in V then if T is trivial, i.e. if it is diffeomorphic to $M' \times D$ where D is a sufficiently small ball of dimension being the codimension of M' in V , and we will equip T with the product metric of $M' \times D$.

If T is not trivial, using a partition of unity of M' , one can construct a metric on T as linear combination of the product metrics on trivialised pieces so that the involution $\iota : T \rightarrow T$ locally given by $(y, d) \mapsto (y, -d)$ for $y \in M', d \in D$ is an isometry. As a consequence, M' is totally geodesic in T .

Since $M' \equiv M' \times \{0\}$ is totally geodesic in T , one has for every smooth function $f : M \rightarrow M'$:

$$\tau_T(f) = \tau_{M'}(f)$$

The crucial property we expect for a global equation of (5), is the following: if the solution initially is in $M' \subset V$ then it remains in M' for all relevant time $t > \alpha$. Eells-Sampson [?] did this by using at the same time 2 different metrics on T , namely the product metric as tubular neighborhood and the Euclidean metric. I choose to present here the formulation of Hamilton, which is conceptually simpler with the only drawback being that we need to establish the uniqueness of solution of (5) first.

After having the global equation, we will prove the short time existence of solution by linearising the equation and using Inverse function theorem. The global formulation and the proof of short-time existence are independent of the negative curvature hypothesis, which will only be used later to establish energy estimates and assure the convergence of long-time solution and the vanishing of its tension field.

2.3 Global equation and Uniqueness of nonlinear heat equation.

Theorem 28 (Global equation). *If the smooth function $F_t : M \times [\alpha, \beta] \rightarrow V$ satisfies*

$$\frac{dF_t}{dt} = \tau_T(F_t) \quad (6)$$

and $F_t(M \times \{\alpha\}) \subset M'$ then $F_t(M \times [\alpha, \omega]) \subset M'$

Proof. Let ι be the isometry of T locally given by $(y, d) \mapsto (y, -d)$ for $(y, d) \in M' \times D \equiv T$ and pose $G_t = \iota F_t$ then G_t and F_t coincide initially since M' is fixed by ι . Moreover

$$\frac{dG_t}{dt} = d\iota \cdot \frac{dF_t}{dt} = d\iota(\tau_T(F_t)) = \tau_T(\iota F_t) = \tau_T(G_t)$$

We conclude that $F_t = G_t = \iota F_t$, hence F_t remains in M' for all relevant t , using the following uniqueness of nonlinear heat equation. \square

Theorem 29 (Uniqueness of solution of nonlinear hear equation). *Let $f_1, f_2 : M \times [\alpha, \omega] \rightarrow M'$ be C^2 functions satisfying the non-linear heat equation $\frac{df_i}{dt} = \tau_{M'}(f_i)$, i.e.*

$$\frac{df^\gamma}{dt} = -\Delta f^\gamma + g^{ij} \Gamma_{\alpha\beta}^{\gamma} f_i^\alpha f_j^\beta$$

where $\Gamma_{\alpha\beta}^{\gamma}$ are Christoffel symbols of M' . Suppose that f_1 and f_2 coincide on $M \times \{\alpha\}$. Then $f_1 = f_2$ on $M \times [\alpha, \omega]$.

Proof. It is sufficient to prove the theorem for ω very close to α , therefore by compactness of M , we can suppose that there exists a finite atlas $M = \bigcup_i U_i$ with $f_1(U_i \times [\alpha, \omega])$ and $f_2(U_i \times [\alpha, \omega])$ being in the same chart V_i of M' . We consider the distance function $\sigma(a, b) = \frac{1}{2} d_{M'}(a, b)^2$ for $a, b \in M'$ to measure the difference between f_1 and f_2 by

$$\rho(x, t) = \sigma(f_1(x, t), f_2(x, t))$$

The strategy is to prove that there exists $C > 0$ such that $\frac{d\rho}{dt} \leq -\Delta\rho + C\rho$, then by Maximum principle, one has $\rho = 0$.

Fix a chart U_i of M and the corresponding V_i of M' , one has by straightforward calculation:

$$\begin{aligned} \frac{d\rho}{dt} = & -\Delta\rho - g^{ij} \left(\frac{\partial^2 \sigma}{\partial f_1^\beta \partial f_1^\gamma} - \frac{\partial \sigma}{\partial f_1^\alpha} \Gamma_{\beta\gamma}^\alpha(f_1) \right) f_{1i}^\beta f_{1j}^\gamma \\ & - g^{ij} \left(\frac{\partial^2 \sigma}{\partial f_2^\beta \partial f_2^\gamma} - \frac{\partial \sigma}{\partial f_2^\alpha} \Gamma_{\beta\gamma}^\alpha(f_2) \right) f_{2i}^\beta f_{2j}^\gamma - 2g^{ij} \frac{\partial^2 \sigma}{\partial f_1^\beta \partial f_2^\gamma} f_{1i}^\beta f_{2j}^\gamma \end{aligned} \quad (7)$$

where g^{ij} is the metric on M and $\Gamma_{\beta\gamma}^\alpha$ are Christoffel symbols of M' .

Let c be a point in the chart V_i and choose the normal coordinates of M' at c . Then for $a, b \in M'$ near c , one has, since $\sigma(a, b) = \sigma(b, a)$ and $\sigma(a, b) = 0$ if $b^\gamma = ka^\gamma$ (the Euclidean straight line from a to ka viewed on M' is a geodesic):

$$\sigma(a, b) = \frac{1}{2} d_{M'}(a, b)^2 = \frac{1}{2} d_E(a, b)^2 + \lambda_{\beta\gamma, \delta} (a^\beta a^\gamma b^\delta + b^\beta b^\gamma a^\delta)$$

where d_E is the Euclidean distance, with $\lambda_{\beta\gamma, \delta} = \lambda_{\gamma\beta, \delta}$ and $\lambda_{\beta\gamma, \delta} + \lambda_{\gamma\delta, \beta} + \lambda_{\beta\delta, \gamma} = 0$. We then have the series development of σ at $(0, 0)$:

$$\sigma(a, b) = \frac{1}{2} \delta_{\beta\gamma} (a^\beta - b^\beta) (a^\gamma - b^\gamma) + \lambda_{\beta\gamma, \delta} (a^\beta a^\gamma b^\delta + b^\beta b^\gamma a^\delta) + O(|a| + |b|)^4 \quad (8)$$

and the development of its derivatives

$$\begin{aligned} \frac{\partial^2 \sigma}{\partial a^\beta \partial b^\gamma} (a, b) &= -\delta_{\beta\gamma} + O(|a| + |b|)^2 \\ \frac{\partial^2 \sigma}{\partial a^\beta \partial a^\gamma} (a, b) &= \delta_{\beta\gamma} + \lambda_{\beta\gamma, \delta} b^\delta + O(|a| + |b|)^2 \\ \frac{\partial^2 \sigma}{\partial b^\beta \partial b^\gamma} (a, b) &= \delta_{\beta\gamma} + \lambda_{\beta\gamma, \delta} a^\delta + O(|a| + |b|)^2 \\ \frac{\partial \sigma}{\partial a^\alpha} (a, b) &= O(|a| + |b|), \quad \Gamma_{\beta\gamma}^\alpha(a) = O(|a|) \end{aligned}$$

So choose c to be the midpoint of $f_1(x, t)$ and $f_2(x, t)$ and $(f_1(x, t), f_2(x, t)) = (w, -w)$ in the chart, one has:

$$\frac{d\rho}{dt} = -\Delta\rho - \left(\delta_{\beta\gamma} - \lambda_{\beta\gamma, \delta} w^\delta + O(|w|^2) \right) f_{1i}^\beta f_{1j}^\gamma g^{ij} - \left(\delta_{\beta\gamma} + \lambda_{\beta\gamma, \delta} w^\delta + O(|w|^2) \right) f_{2i}^\beta f_{2j}^\gamma g^{ij} \quad (9)$$

$$- 2 \left(-\delta_{\beta\gamma} + O(|w|^2) \right) f_{1i}^\beta f_{2j}^\gamma g^{ij} \quad (10)$$

$$= -\Delta\rho - |df_1 - df_2|^2 - w^\delta \lambda_{\beta\gamma, \delta} g^{ij} \left(f_{2i}^\beta f_{2j}^\gamma - f_{1i}^\beta f_{1j}^\gamma \right) + O(|w|^2) \quad (11)$$

The last term of (11) can be bounded as follows:

$$\begin{aligned} \left| w^\delta \lambda_{\beta\gamma,\delta} \left(f_{2i}^\beta f_{2j}^\gamma - f_{1i}^\beta f_{1j}^\gamma \right) g^{ij} \right| &= \left| w^\delta \lambda_{\beta\gamma,\delta} \left(f_{2i}^\beta (f_{2j}^\gamma - f_{1j}^\gamma) + f_{1j}^\gamma (f_{2i}^\beta - f_{1i}^\beta) \right) g^{ij} \right| \\ &\leq 2 |w^\delta \lambda_{\beta\gamma,\delta}| |df_2 - df_1| (|df_1| + |df_2|) \\ &\leq |df_1 - df_2|^2 + O(|w|^2) \end{aligned}$$

where for the last inequality, we use $2uv \leq u^2 + v^2$ and the fact that $|df_1|$ and $|df_2|$ are bounded on M . The estimate (11) can be continued:

$$\frac{d\rho}{dt} \leq -\Delta\rho + C(x, t)|w|^2 \leq -\Delta\rho + C\rho$$

where $C > 0$ is a constant chosen to dominate all $C(x, t)$ for $x \in M$ in all charts and $t \in [\alpha, \omega]$. \square

Remark 30. The original proof of [?] made the reduction of the first order of w in (10) using the following development of σ :

$$\sigma = \frac{1}{2} \delta_{\beta\gamma} (a^\beta - b^\beta) (a^\gamma - b^\gamma) + \lambda_{\beta\gamma,\delta} (a^\beta - b^\beta) (a^\gamma - b^\gamma) (a^\delta + b^\delta) + O(|a| + |b|)^4$$

which was justified by $\sigma(a, b) = \sigma(b, a)$ and $\sigma(a, a) = 0$. It can be proved that this is equivalent to (8) and the symmetries $\lambda_{\beta\gamma,\delta} = \lambda_{\gamma\beta,\delta}$, $\lambda_{\beta\gamma,\delta} + \lambda_{\gamma\delta,\beta} + \lambda_{\delta\beta,\gamma} = 0$.

As a side note, if a, b, c are on S^2 with $d(a, c) = d(b, c) = x \ll 1$ and the lines from a and b to c are orthogonal at c , then the geodesic distance $d(a, b) = \arccos(\cos^2(x)) = x\sqrt{2} - \frac{1}{6\sqrt{2}}x^3 + O(x^4)$. So $\sigma(a, b) = \frac{1}{2}d(a, b)^2$ has no third-order term.

3 A few energy estimates.

3.1 Estimate of density energies

We finish this part with a few straightforward computation concerning the **potential energy** $e(f_t) = \frac{1}{2}|\nabla f_t|^2$ and the **kinetic energy** $k(f_t) = \frac{1}{2}|\frac{df_t}{dt}|^2$ of a nonlinear heat flow f_t satisfying (5).

Theorem 31 (Density of Potential energy). *If f_t satisfies (5) then*

$$\frac{de(f_t)}{dt} = -\Delta e(f_t) - |\beta(f_t)|^2 - \langle \text{Ric}(M) \nabla_v f_t, \nabla_v f_t \rangle + \langle \text{Riem}(M')(\nabla_v f_t, \nabla_w f_t) \nabla_v f_t, \nabla_w f_t \rangle$$

where $e(f_t)$ is the potential energy density and $\beta(f_t)$ is the fundamental form and in the curvature terms, the vectors v and w are contracted.

In particular, if $\text{Riem}(M') \leq 0$ and $\text{Ric}(M) \geq -C$ then

$$\frac{de}{dt} \leq -\Delta e + Ce - |\beta(f_t)|^2 \tag{12}$$

Proof. Apply Lemma 10 to $s = df_t$ and the Riemannian-connected bundle F^*TM' over $M \times [\alpha, \omega]$ where $F(\cdot, t) = f_t$, the curvature terms cancel out and it remains to see that $\frac{de(f_t)}{dt} = -\langle df_t, \Delta df_t \rangle$, meaning that $\tilde{\nabla}_{\partial t} df_t = -\Delta df_t$. This can be easily justified:

$$\tilde{\nabla}_{\partial t} df_t = \tilde{\nabla}_{\partial t} \tilde{\nabla}^M F = \tilde{\nabla}^M \tilde{\nabla}_{\partial t} F = \tilde{\nabla}^M \tau(f_t) = -D\delta(df_t) = -\Delta df_t$$

where the last "=" is due to $Ddf_t = 0$. Note that D and δ are the exterior derivative and its adjoint of the bundle $(f_t)^*TM'$ on M , where t can be fixed after the third "=" sign. \square

Theorem 32 (Density of Kinetic energy). *If f_t satisfies (5) then*

$$\frac{dk(f_t)}{dt} = -\Delta k(f_t) - \left| \nabla \frac{\partial f_t}{\partial t} \right|^2 + \left\langle \text{Riem}(M')(\nabla_v f_t, \frac{\partial f_t}{\partial t}) \nabla_v f_t, \frac{\partial f_t}{\partial t} \right\rangle$$

where $k(f_t)$ is the kinetic energy density and in the curvature terms, the vectors v is contracted, In particular, if $\text{Riem}(M') \leq 0$ then

$$\frac{dk}{dt} \leq -\Delta k - \left| \nabla \frac{\partial f_t}{\partial t} \right|^2 \quad (13)$$

Proof. Let $F : I \times M \rightarrow M'$ be the total function with $F(t, \cdot) = f_t$ for $t \in I = [\alpha, \omega]$ and $E = F^*TM'$ is a Riemannian-connected bundle on $I \times M$ with curvature form Θ , then

$$\tilde{\nabla}_{\partial t} \tilde{\nabla}_v (dF.v) = \tilde{\nabla}_v \tilde{\nabla}_{\partial t} (dF.v) + \Theta(\partial t, v) dF.v \quad (14)$$

where dF is the exterior derivative of f_t on M . Note that $\tilde{\nabla}_v \tilde{\nabla}_{\partial t} (dF.v) = \tilde{\nabla}_v (\tilde{\nabla}_{\partial t} dF).v = \tilde{\nabla}_v (\tilde{\nabla}^M \frac{\partial f_t}{\partial t}).v$ since $\tilde{\nabla}^M \frac{\partial f_t}{\partial t} = \tilde{\nabla}_{\partial t}^{I \times M} dF = \tilde{\nabla}_{\partial t}^I dF$ because $\tilde{\nabla}$ is torsionless on M' . Plugging this in (14) and taking contraction in v , one has

$$\tilde{\nabla}_{\partial t} \tau(f_t) = -\tilde{\Delta} \frac{\partial f_t}{\partial t} + \text{Tr}(v \mapsto \Theta(\partial t, v) dF.v) \quad (15)$$

But $\Theta_\alpha^\beta = R'_{\alpha\nu\mu} F_i^\mu F_j^\nu dx^i \otimes dx^j$ where R' denotes the Riemannian curvature of M' and the indices i, j can be 0, with $x^0 \equiv t$. Hence

$$\Theta(\partial t, v) dF.v = R'_{\alpha\nu\mu} \frac{\partial f_t^\mu}{\partial t} \frac{\partial f_t^\nu}{\partial v} \frac{\partial f_t^\alpha}{\partial v} \tilde{e}_\beta = \text{Riem}(M') \left(\nabla_v f_t, \frac{\partial f_t}{\partial t} \right) \nabla_v f_t$$

Plugging in (15) and taking inner product with $\frac{\partial f_t}{\partial t}$, one has

$$\begin{aligned} \frac{\partial k(f_t)}{\partial t} &= \left\langle \tilde{\nabla}_{\partial t} \tau(f_t), \frac{\partial f_t}{\partial t} \right\rangle = - \left\langle \tilde{\Delta} \frac{\partial f_t}{\partial t}, \frac{\partial f_t}{\partial t} \right\rangle + \left\langle \text{Riem}(M')(\nabla_v f_t, \frac{\partial f_t}{\partial t}) \nabla_v f_t, \frac{\partial f_t}{\partial t} \right\rangle \\ &= -\Delta \left(\frac{1}{2} \left| \frac{\partial f_t}{\partial t} \right|^2 \right) - \left| \tilde{\nabla} \frac{\partial f_t}{\partial t} \right|^2 + \left\langle \text{Riem}(M')(\nabla_v f_t, \frac{\partial f_t}{\partial t}) \nabla_v f_t, \frac{\partial f_t}{\partial t} \right\rangle \end{aligned}$$

\square

3.2 Estimate of total energies

We will now work with the total energies, in particular the **total potential energy** $E(f_t) := \int_M e(f_t)$ and **total kinetic energy** $K(f_t) := \int_M k(f_t)$. Since tension field is the gradient of E , one has:

Theorem 33. *If $f_t : M \rightarrow M'$ satisfies (5) then*

$$\frac{dE(f_t)}{dt} = - \int_M \left\langle \tau(f_t), \frac{\partial f_t}{\partial t} \right\rangle = - \int_M |\tau(f_t)|^2 = -2K(f_t) \leq 0.$$

Integrating Theorem 32 on M then using Theorem 33, one obtains:

Theorem 34. *If f_t satisfies (5) and $\text{Riem}(M') \leq 0$ then $\frac{d}{dt}K(f_t) \leq 0$ and one has*

1. *The total potential energy $E(f_t)$ is ≥ 0 , decreasing and convex.*
2. *The total kinetic energy $K(f_t)$ is ≥ 0 , decreasing and if $\omega = +\infty$ then $\lim_{t \rightarrow \infty} K(f_t) = 0$.*

In particular, $\int_{M \times \{\tau\}} |\nabla f|^2$ and $\int_{M \times \{\tau\}} \left| \frac{\partial f_t}{\partial t} \right|^2$ are bounded above by a constant $C > 0$ independent of the time $\tau \in [\alpha, \omega]$.

Note that we ruled out the case $K(f_t)$ decreases to a strictly positive limit because $E(f_t)$ is bounded below and $\frac{d}{dt}E(f_t) = -2K(f_t)$.

Integrating Theorem 31 on M then using Theorem 34, one has:

Theorem 35. *If f_t satisfies (5) and $\text{Riem}(M') \leq 0$ and $\text{Ric}(M)$ is bounded below then*

$$\int_M |\beta(f_t)|^2 \leq C$$

for all time t where the constant C only depends on the curvature of M, M' and the initial total potential and kinetic energy, in particular, C does not depend on t .

This means that $\|f_t\|_{W^{2,2}(M)}$ is bounded by a constant C only depending on the curvatures and initial total energies.

Corollary 36 (Boundedness in $W^{2,2}(M)$). *If F_t satisfies (6) and $\text{Riem}(M') \leq 0$ and $\text{Ric}(M)$ is bounded below then*

$$\|F_t\|_{W^{2,2}(M)}^2 := \int_M |\beta(F_t)|^2 + |\nabla F_t|^2 + |F|^2 \leq C$$

for all time t where the constant C only depends on the curvature of M, M' and the initial total potential and kinetic energy, in particular, C does not depend on t .

Note that the term $|F|^2$ is trivially bounded since the image of F remains in an Euclidean ball B .