

Examiner: Xiangfeng Yang (Tel: 013 28 57 88).**Things allowed (Hjälpmmedel):** a calculator.**Scores rating (Betygsgränser):** 8-11 points giving rate 3; 11.5-14.5 points giving rate 4; 15-18 points giving rate 5.
Write down all necessary steps in solutions in order to receive as many points as possible.

1 (3 points)

A system contains 3 independent components A , B and C . The probability that component A fails is $P(A) = 0.8$, the probability that component B fails is $P(B) = 0.5$, and the probability that component C fails is $P(C) = 0.3$.

- (1.1) (1p) Find the probability that all 3 components fail, namely, $P(A \cap B \cap C)$.
- (1.2) (1p) Find the probability that exactly 1 component fails.
- (1.3) (1p) Find the probability that at least 1 component fails.

Solution. (1.1) It is from independence of A , B and C that

$$P(A \cap B \cap C) = P(A) \cdot P(B) \cdot P(C) = 0.8 \cdot 0.5 \cdot 0.3 = 0.12.$$

(1.2)

$$\begin{aligned} P(\text{exactly 1 component fails}) &= P(\text{only A fails}) + P(\text{only B fails}) + P(\text{only C fails}) \\ &= P(A \cap B' \cap C') + P(A' \cap B \cap C') + P(A' \cap B' \cap C) \\ &= 0.8 \cdot 0.5 \cdot 0.7 + 0.2 \cdot 0.5 \cdot 0.7 + 0.2 \cdot 0.5 \cdot 0.3 = 0.28 + 0.07 + 0.03 = 0.38. \end{aligned}$$

(1.3)

$$P(\text{at least 1 component fails}) = P(A \cup B \cup C) = 1 - P(A' \cap B' \cap C') = 1 - 0.2 \cdot 0.5 \cdot 0.7 = 1 - 0.07 = 0.93.$$

□

2 (3 points)

Let (X, Y) be a two dimensional continuous random variable with the following joint probability density function

$$f(x, y) = \begin{cases} 8xy, & \text{if } 0 < x < 1 \text{ and } 0 < y < x, \\ 0, & \text{otherwise.} \end{cases}$$

- (2.1) (1p) Find the probability $P(X > 0.5 \cap Y < 0.5)$. (Also written as $P(X > 0.5, Y < 0.5)$.)
- (2.2) (2p) Are X and Y independent? Why?

Solution. (2.1) Draw integral region,

$$P(X > 0.5 \cap Y < 0.5) = \int_{0.5}^1 \left(\int_0^{0.5} 8xy dy \right) dx = \int_{0.5}^1 x dx = 0.375.$$

(2.2) Draw integral region, and marginal pmfs can be found as follows

$$f_X(x) = \int_0^x 8xy dy = 4x^3, \quad f_Y(y) = \int_y^1 8xy dx = 4y(1-y^2).$$

It is clear that $f(x, y) \neq f_X(x) \cdot f_Y(y)$, therefore X and Y are NOT independent.

□

3 (3 points)

Let X be the distribution of numbers of car accidents in one day on a certain highway, and assume that X has the following probability mass function (pmf) $p(x)$:

X	0	1	2
$p(x)$	0.75	0.15	0.10

(3.1) (1p) Find the mean $E(X)$ and the variance $V(X)$ of X .

(3.2) (2p) Let X_1, X_2, \dots, X_{36} denote numbers of car accidents in 36 days on the highway. Assume that X_1, X_2, \dots, X_{36} are independent and each has the same distribution as X . Find the probability that there are at most 20 car accidents in 36 days, namely, $P(X_1 + X_2 + \dots + X_{36} \leq 20)$.

Solution. (3.1)

$$\mu = E(X) = 0 \times 0.75 + 1 \times 0.15 + 2 \times 0.10 = 0.35,$$

$$\sigma^2 = V(X) = E(X^2) - \mu^2 = 0^2 \times 0.75 + 1^2 \times 0.15 + 2^2 \times 0.10 - 0.35^2 = 0.55 - 0.1225 = 0.4275.$$

(3.2)

$$\begin{aligned} P(X_1 + X_2 + \dots + X_{36} \leq 20) \\ &= P\left(\frac{X_1 + X_2 + \dots + X_{36}}{36} < 20/36\right) = P(\bar{X} < 20/36) \\ &= P\left(\frac{\bar{X} - \mu}{\sigma/\sqrt{n}} < \frac{20/36 - 0.35}{\sqrt{0.4275}/\sqrt{36}}\right) = P(N(0, 1) < 1.89) = \Phi(1.89) = 0.9706. \end{aligned}$$

□

4 (3 points)

Suppose that a population X has a probability density function (pdf) $f(x) = e^{-(x-\theta)}$ for $x \geq \theta$ (otherwise it is 0), where θ is an unknown parameter. A sample $\{x_1, x_2, \dots, x_n\}$ is taken from this population.

(4.1) (1p) Find a point estimate $\hat{\theta}_{MM}$ of θ based on the method of moments.

(4.2) (2p) Find a point estimate $\hat{\theta}_{ML}$ of θ based on the maximum-likelihood method.

Solution. (4.1) We first compute the mean $E(X)$ of X as follows

$$\begin{aligned} E(X) &= \int_{\theta}^{+\infty} x \cdot e^{-(x-\theta)} dx = e^{\theta} \int_{\theta}^{+\infty} x \cdot e^{-x} dx = -e^{\theta} \int_{\theta}^{+\infty} x \cdot (e^{-x})' dx \\ &= (\text{integration by parts}) = -e^{\theta} \left([xe^{-x}]_{\theta}^{+\infty} - \int_{\theta}^{+\infty} x' \cdot e^{-x} dx \right) \\ &= -e^{\theta} \left(-\theta e^{-\theta} - \int_{\theta}^{+\infty} \cdot e^{-x} dx \right) = -e^{\theta} (-\theta e^{-\theta} - e^{-\theta}) = \theta + 1. \end{aligned}$$

Therefore, the equation $E(X) = \bar{x}$ give

$$\theta + 1 = \bar{x} \implies \hat{\theta}_{MM} = \bar{x} - 1.$$

(4.2) The likelihood function is

$$L(\theta) = f(x_1) \cdot f(x_2) \cdot \dots \cdot f(x_n) = e^{-(x_1-\theta)} \cdot e^{-(x_2-\theta)} \cdot \dots \cdot e^{-(x_n-\theta)} = e^{n\theta - (x_1+x_2+\dots+x_n)}.$$

The logarithmic likelihood function is

$$\ln L(\theta) = n\theta - (x_1 + x_2 + \dots + x_n).$$

The derivative $\ln' L(\theta) = n > 0$ is strictly positive, implying that the function $\ln L(\theta)$ is increasing in θ . The non-trivial domain of $L(\theta)$ is: $\theta \leq \min\{x_1, x_2, \dots, x_n\}$ (Why? because for each $f(x_i)$ in $L(\theta)$ the domain is $\theta \leq x_i$, and combining all these n functions gives such a domain). This implies that

$$\hat{\theta}_{ML} = \min\{x_1, x_2, \dots, x_n\}.$$

□

5 (3 points)

Volvo company claims that 2025-model XC60 plug-in hybrid cars have 60 km electric range (which is the distance using an electric motor powered by battery). Let X be the distribution of electric ranges of all 2025-model XC60 plug-in hybrid cars, and assume $X \sim N(\mu, \sigma^2)$. A sample $\{x_1, x_2, \dots, x_{16}\}$ taken from X has a sample mean 56 km and a sample standard deviation 6 km.

(5.1) (1p) Construct a 95% two-sided confidence interval I_μ of μ .

(5.2) (2p) Does the above sample provide any evidence that $\mu < 60$? Answer this using an appropriate 99% one-sided confidence interval I_μ of μ . (Hint: one-sided!!! 99%!!!)

Solution. (5.1) This is Case 1.3:

$$I_\mu = \bar{x} \mp t_{\alpha/2}(n-1) \frac{s}{\sqrt{n}} = 56 \mp t_{0.025}(15) \frac{6}{\sqrt{16}} = 56 \mp 2.13 \frac{6}{\sqrt{16}} = 56 \mp 3.195 = (52.805, 59.195).$$

(5.2) One-sided CI is (Case 1.3)

$$I_\mu = (-\infty, \bar{x} + t_\alpha(n-1) \frac{s}{\sqrt{n}}) = (-\infty, 56 + t_{0.01}(15) \frac{6}{\sqrt{16}}) = (-\infty, 56 + 2.60 \cdot 1.5) = (-\infty, 59.9).$$

Since $59.9 < 60$, the sample DOES provide evidence that $\mu < 60$. □

6 (3 points)

It was reported by many teachers that students in ⑤ program at LiU do not usually attend lectures. The committee of ⑤ program started to investigate the attendance rate p . To this end, the committee randomly went to a lecture with 200 registered students from ⑤ program, and only 56 attended the lecture (that is, sample attendance rate is $\hat{p} = 56/200 = 28\%$). Based on this observation, can the committee conclude that $p < 30\%$ with confidence 97.5%? Answer this by hypothesis testing (HT).

(6.1) (1p) Write down H_0 and H_a .

(6.2) (1p) Is H_0 rejected? Why?

(6.3) (1p) Find the power $h(25\%)$ of the test in (6.2) when the real attendance rate is $p = 25\%$.

Solution. (6.1)

$$\begin{cases} H_0 : p = 30\% \\ H_a : p < 30\% \end{cases}$$

(6.2) Since $n\hat{p} \geq 10$ and $n(1 - \hat{p}) \geq 10$, it follows that

$$TS = \frac{\hat{p} - p_0}{\sqrt{p_0(1 - p_0)/n}} = \frac{28\% - 30\%}{\sqrt{30\%(1 - 30\%)/200}} = -0.617, \quad C = (-\infty, -z_\alpha) = (-\infty, -1.96).$$

It is clear that $TS \notin C$, so H_0 is NOT rejected, which suggests that there is no evidence showing that $p < 30\%$.
(6.3)

$$\begin{aligned} h(25\%) &= P(H_0 \text{ is rejected when } p = 25\%) = P\left(\frac{\hat{P} - p_0}{\sqrt{p_0(1 - p_0)/n}} < -1.96 \text{ when } p = 25\%\right) \\ &= P(\hat{P} < 0.2365 \text{ when } p = 25\%) = P\left(\frac{\hat{P} - p}{\sqrt{p(1 - p)/n}} < \frac{0.2365 - p}{\sqrt{p(1 - p)/n}} \text{ when } p = 25\%\right) \\ &= P(N(0, 1) < \frac{0.2365 - 0.25}{\sqrt{0.25(1 - 0.25)/200}}) = P(N(0, 1) < -0.44) = \Phi(-0.44) = 1 - \Phi(0.44) = 1 - 0.6700 = 0.3300. \end{aligned}$$

□

1. Basic probability

(1.1) Conditional probability $P(A|B) = \frac{P(A \cap B)}{P(B)}$.

(1.2) Total probability $P(B) = \sum_{i=1}^k P(B|A_i)P(A_i)$ where $\{A_i\}$ are disjoint and $\cup_{i=1}^k A_i = S$.

(1.3) Bayes' Theorem $P(A_j|B) = \frac{P(B|A_j)P(A_j)}{\sum_{i=1}^k P(B|A_i)P(A_i)}$ where $\{A_i\}$ are in (1.2).

2. Random variables (r.v.s)

(2.1) Discrete r.v. X has a pmf $p(x) = P(X = x)$ satisfying $p(x) \geq 0$ and $\sum p(x_i) = 1$,

X	x_1	x_2	\cdots	x_n	\cdots
$p(x)$	$p(x_1)$	$p(x_2)$	\cdots	$p(x_n)$	\cdots

Expectation (or *Expected value* or *mean*) $\mu_X = E(X) = \sum x_i p(x_i)$;
 Variance $\sigma_X^2 = V(X) = E(X - \mu_X)^2 = E(X^2) - \mu_X^2 = \sum x_i^2 p(x_i) - (\sum x_i p(x_i))^2$.

(2.2) Continuous r.v. X has a pdf $f(x)$ satisfying $f(x) \geq 0$ and $\int_{-\infty}^{\infty} f(x)dx = 1$,

$$P(a < X < b) = \int_a^b f(x)dx.$$

Expectation (or *Expected value* or *mean*) $\mu_X = E(X) = \int_{-\infty}^{\infty} xf(x)dx$;
 Variance $\sigma_X^2 = V(X) = E(X - \mu_X)^2 = E(X^2) - \mu_X^2 = \int_{-\infty}^{\infty} x^2 f(x)dx - (\int_{-\infty}^{\infty} xf(x)dx)^2$.

(2.3) Cumulative distribution function (cdf) of a r.v. X is $F(x) = P(X \leq x)$.

(2.4) X and Y are r.v.s, a , b and c are scalars, then

$$E(aX + bY + c) = aE(X) + bE(Y) + c,$$

$$V(aX + bY + c) = a^2V(X) + b^2V(Y) + 2ab \text{cov}(X, Y),$$

$$E(g(X, Y)) = \begin{cases} \sum_{i,j} g(x_i, y_j) \cdot p(x_i, y_j), & \text{for discrete } (X, Y), \\ \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} g(x, y) \cdot f(x, y)dxdy, & \text{for continuous } (X, Y). \end{cases}$$

(2.5) • Discrete r.v. (X, Y) has a joint pmf $p(x, y)$ satisfying $p(x, y) \geq 0$ and $\sum_{x_i} \sum_{y_i} p(x_i, y_i) = 1$.

The marginal pmf of X is $p_X(x) = \sum_y p(x, y)$;
 The marginal pmf of Y is $p_Y(y) = \sum_x p(x, y)$;

X and Y are *independent* if $p(x, y) = p_X(x) \cdot p_Y(y)$.

The marginal pdf of X is $f_X(x) = \int_{-\infty}^{\infty} f(x, y)dy$;

The marginal pdf of Y is $f_Y(y) = \int_{-\infty}^{\infty} f(x, y)dx$;
 X and Y are *independent* if $f(x, y) = f_X(x) \cdot f_Y(y)$.

3. Several special r.v.s

(3.1) $X \sim Bin(n, p)$ has a pmf $p(x) = P(X = x) = \binom{n}{x} \cdot p^x \cdot (1-p)^{n-x}$, $x = 0, 1, 2, \dots, n$.

$$E(X) = n \cdot p, \quad V(X) = n \cdot p \cdot (1-p).$$

(3.2) $X \sim Po(\lambda)$ has a pmf $p(x) = P(X = x) = \frac{e^{-\lambda} \lambda^x}{x!}$, $x = 0, 1, 2, \dots$.

$$E(X) = \lambda, \quad V(X) = \lambda.$$

(3.3) $X \sim Hypergeometric$ has a pmf $p(x) = P(X = x) = \frac{\binom{M}{x} \binom{N-M}{n-x}}{\binom{N}{n}}$.

(3.4) $X \sim Exp(\lambda)$ has a pdf

$$f(x) = \begin{cases} \lambda e^{-\lambda x}, & x \geq 0, \\ 0, & \text{otherwise.} \end{cases}$$

(3.5) $X \sim N(\mu, \sigma^2)$ has a pdf

$$f(x) = \frac{1}{\sigma \sqrt{2\pi}} e^{-\frac{(x-\mu)^2}{2\sigma^2}}, -\infty < x < \infty.$$

$E(X) = \mu, \quad V(X) = \sigma^2$

(3.6) $X \sim U(a, b)$ has a pdf

$$f(x) = \begin{cases} \frac{1}{b-a}, & a < x < b, \\ 0, & \text{otherwise.} \end{cases}$$

$$E(X) = \frac{a+b}{2}, \quad V(X) = \frac{(b-a)^2}{12}.$$

4. Central Limit Theorem (CLT)

Suppose that a population has mean = μ and variance = σ^2 . A random sample $\{X_1, X_2, \dots, X_n\}$ from this population is given. Then for large $n \geq 30$,

$$\frac{\bar{X} - \mu}{\sigma/\sqrt{n}} \sim N(0, 1).$$

- If the population is normal, then (1) holds for any n .
- Note that $\mu = E(\bar{X})$ and $(\sigma/\sqrt{n})^2 = V(\bar{X})$.

5. Several notations in statistics

(5.1) Sample mean: $\bar{X} = \frac{X_1 + X_2 + \dots + X_n}{n} = \frac{\sum X_i}{n}$; $\bar{x} = \frac{x_1 + x_2 + \dots + x_n}{n} = \frac{\sum x_i}{n}$.

(5.2) Sample variance:

$$S^2 = \frac{\sum (X_i - \bar{X})^2}{n-1} = \frac{1}{n-1} \left(\sum X_i^2 - \frac{(\sum X_i)^2}{n} \right); \quad s^2 = \frac{\sum (x_i - \bar{x})^2}{n-1} = \frac{1}{n-1} \left(\sum x_i^2 - \frac{(\sum x_i)^2}{n} \right).$$

• Capital letters \bar{X} and S^2 refer to the objects based on random sample (therefore they are in general r.v.s), while small letters \bar{x} and s^2 are the objects based on observations (so they are scalars).

(5.3) A point estimator of θ obtained by Maximum Likelihood method is denoted as $\hat{\theta}_{ML}$.

6. Confidence Interval (CI)

In this course, three types of confidence intervals are studied depending on the unknown population parameter(s): CI-1 (confidence intervals for population mean(s)), CI-2 (confidence intervals for population variance(s)), and CI-3 (confidence intervals for population proportion(s)).

CI-1: $(1 - \alpha)$ CI of a population mean μ

case 1.1 (any n) If population $X \sim N(\mu, \sigma^2)$ and σ^2 is known, then $\frac{\bar{X} - \mu}{\sigma/\sqrt{n}} \sim N(0, 1)$ and

$$I_\mu = \left(\bar{x} - z_{\alpha/2} \cdot \frac{\sigma}{\sqrt{n}}, \bar{x} + z_{\alpha/2} \cdot \frac{\sigma}{\sqrt{n}} \right) := \bar{x} \mp z_{\alpha/2} \cdot \frac{\sigma}{\sqrt{n}}.$$

case 1.2 ($n \geq 30$) For any population X , it holds that $\frac{\bar{X} - \mu}{\sigma/\sqrt{n}} \sim N(0, 1)$ and

$$I_\mu = \bar{x} \mp z_{\alpha/2} \cdot \frac{\sigma}{\sqrt{n}} \text{ or } I_\mu = \bar{x} \mp z_{\alpha/2} \cdot \frac{\hat{\sigma}}{\sqrt{n}}.$$

case 1.3 (any n) If population $X \sim N(\mu, \sigma^2)$ and σ^2 is unknown, then $\frac{\bar{X} - \mu}{S/\sqrt{n}} \sim T(n-1)$ and

$$I_\mu = \bar{x} \mp t_{\alpha/2}(n-1) \cdot \frac{s}{\sqrt{n}}.$$

CI-1': $(1 - \alpha)$ CI of the difference of two population means $\mu_X - \mu_Y$

case 1.1' (any n_1, n_2) If independent populations $X \sim N(\mu_X, \sigma_X^2)$, $Y \sim N(\mu_Y, \sigma_Y^2)$, and σ_X^2, σ_Y^2 are known, then $\frac{(\bar{X} - \bar{Y}) - (\mu_X - \mu_Y)}{\sqrt{\frac{\sigma_X^2}{n_1} + \frac{\sigma_Y^2}{n_2}}} \sim N(0, 1)$, and $I_{\mu_X - \mu_Y} = (\bar{x} - \bar{y}) \mp z_{\alpha/2} \cdot \sqrt{\frac{\sigma_X^2}{n_1} + \frac{\sigma_Y^2}{n_2}}$.

case 1.2' ($n_1, n_2 \geq 30$) For any independent populations X and Y , it holds that



$$\frac{(\bar{X} - \bar{Y}) - (\mu_X - \mu_Y)}{\sqrt{\frac{\sigma_X^2}{n_1} + \frac{\sigma_Y^2}{n_2}}} \sim N(0, 1) \text{ and}$$

$$I_{\mu_X - \mu_Y} = (\bar{x} - \bar{y}) \mp z_{\alpha/2} \cdot \sqrt{\frac{\sigma_X^2}{n_1} + \frac{\sigma_Y^2}{n_2}} \text{ or } I_{\mu_X - \mu_Y} = (\bar{x} - \bar{y}) \mp z_{\alpha/2} \cdot \sqrt{\frac{\hat{\sigma}_X^2}{n_1} + \frac{\hat{\sigma}_Y^2}{n_2}}.$$

case 1.3' (any n_1, n_2) If independent populations $X \sim N(\mu_X, \sigma_X^2)$, $Y \sim N(\mu_Y, \sigma_Y^2)$, where σ_X^2, σ_Y^2 are unknown but $\sigma_X^2 = \sigma_Y^2$, then

$$\frac{(\bar{X} - \bar{Y}) - (\mu_X - \mu_Y)}{\sqrt{\frac{1}{n_1} + \frac{1}{n_2}}} \sim T(n_1 + n_2 - 2), \text{ where } S^2 = \frac{(n_1 - 1)S_X^2 + (n_2 - 1)S_Y^2}{n_1 + n_2 - 2}, \text{ and}$$

$$I_{\mu_X - \mu_Y} = (\bar{x} - \bar{y}) \mp t_{\alpha/2}(n_1 + n_2 - 2) \cdot s \sqrt{\frac{1}{n_1} + \frac{1}{n_2}}.$$



CI-2: $(1 - \alpha)$ CI of population variance(s) σ^2


- If a population $X \sim N(\mu, \sigma^2)$ and σ^2 is unknown, then $\frac{(n-1)S^2}{\sigma^2} \sim \chi^2(n-1)$, and

$$I_{\sigma^2} = \left(\frac{(n-1)s^2}{\chi_{\alpha/2}^2(n-1)}, \frac{(n-1)s^2}{\chi_{1-\alpha/2}^2(n-1)} \right).$$

- If two independent populations $X \sim N(\mu_X, \sigma^2)$ and $Y \sim N(\mu_Y, \sigma^2)$, and σ^2 is unknown, then $\frac{(n_1+n_2-2)S^2}{\sigma^2} \sim \chi^2(n_1 + n_2 - 2)$, and

$$I_{\sigma^2} = \left(\frac{(n_1 + n_2 - 2)s^2}{\chi_{\alpha/2}^2(n_1 + n_2 - 2)}, \frac{(n_1 + n_2 - 2)s^2}{\chi_{1-\alpha/2}^2(n_1 + n_2 - 2)} \right),$$

where $S^2 = \frac{(n_1-1)S_X^2 + (n_2-1)S_Y^2}{n_1 + n_2 - 2}$.



CI-3: $(1 - \alpha)$ CI of population proportion(s)

- If a (large) population has an unknown proportion p , then $\frac{\hat{p} - p}{\sqrt{p(1-p)/n}} \sim N(0, 1)$ if $n\hat{p} \geq 10$ and $n(1-\hat{p}) \geq 10$ with $\hat{p} = x/n$, and $I_p = \hat{p} \mp z_{\alpha/2} \cdot \sqrt{\frac{\hat{p}(1-\hat{p})}{n}}$.

- If two independent (large) populations have unknown proportions p_1 and p_2 , then

$$\frac{(\hat{p}_1 - \hat{p}_2) - (p_1 - p_2)}{\sqrt{\frac{p_1(1-p_1)}{n_1} + \frac{p_2(1-p_2)}{n_2}}} \sim N(0, 1)$$

if $n_i\hat{p}_i \geq 10$ and $n_i(1 - \hat{p}_i) \geq 10$ for $i = 1, 2$, and $I_{p_1 - p_2} = (\hat{p}_1 - \hat{p}_2) \mp z_{\alpha/2} \cdot \sqrt{\frac{\hat{p}_1(1-\hat{p}_1)}{n_1} + \frac{\hat{p}_2(1-\hat{p}_2)}{n_2}}$.

7. Hypothesis Test (HT)

	H_0 is true	H_0 is false and $\theta = \theta_1$
reject H_0	(type I error or significance level) α	(power) $h(\theta_1) = 1 - h(\theta_1)$
don't reject H_0	1 - α	(type II error) $\beta(\theta_1) = 1 - h(\theta_1)$

reject $H_0 \Leftrightarrow TS \in C \Leftrightarrow p\text{-value} < \alpha$

χ^2 tests for populations (non-parametric tests)

Suppose that for a random sample of a population X , the n elements of it are classified into k disjoint groups $A_i, 1 \leq i \leq k$. For each group $A_i, 1 \leq i \leq k$, suppose that there are $N_i, 1 \leq i \leq k$ elements inside. Let $p_i = P(A_i)$ assuming a given distribution of X . Note that $p_1 + p_2 + \dots + p_k = 1$ and $N_1 + N_2 + \dots + N_k = n$. One wants to test the hypotheses

$$H_0 : P(A_i) = p_i, \quad 1 \leq i \leq k, \quad H_a : P(A_i) \neq p_i \text{ for some } 1 \leq i \leq k.$$

If n is large in the sense that $np_i \geq 5$ for all $1 \leq i \leq k$, then the test statistic is

$$\sum_{i=1}^k \frac{(N_i - np_i)^2}{np_i} \approx \chi^2(k-1).$$

Therefore the observation of the test statistic is

$$TS = \sum_{i=1}^k \frac{(n_i - np_i)^2}{np_i}, \text{ where } n_i \text{ is the observation of } N_i, 1 \leq i \leq k.$$

For the critical region C , one can take (note that if H_0 is true, then TS should be close to zero)

$$C = (\chi_\alpha^2(k-1), \infty).$$

The conclusion would be $TS \in C \iff H_0$ is rejected.

8. Linear and logistic regression

(Multiple) linear regression: $Y = \beta_0 + \beta_1 x_1 + \dots + \beta_k x_k + \varepsilon, \varepsilon \sim N(0, \sigma^2)$.

• Y : response variable (which is normal r.v.), $\{x_1, \dots, x_k\}$: predictors (which are scalars).

• sample: $\{(x_{11}, \dots, x_{1k}; y_1), (x_{21}, \dots, x_{2k}; y_2), \dots, (x_{n1}, \dots, x_{nk}; y_n)\}$.

• how to estimate $\beta_j \approx \hat{\beta}_j$: least square method, that is, to minimize $\sum_{i=1}^n (\hat{y}_i - y_i)^2$, where the estimated (multiple) linear regression line \hat{y} is

$$\hat{y} = \beta_0 + \hat{\beta}_1 x_1 + \dots + \hat{\beta}_k x_k.$$

• $\frac{\hat{\beta}_j - \beta_j}{s.e(\hat{\beta}_j)} \sim T(n-k-1)$, this helps determine whether or not the real $\beta_j = 0$?

• $\sigma^2 \approx \frac{SSE}{n-k-1}$, this gives an estimation of the size of the error.

• $R^2 = \frac{SSE}{SS_T}$ this gives how well the model is (if $R^2 \approx 1$, then the model fits the sample very well).

• How to test $\beta_1 = \dots = \beta_k = 0$? Use the random variable $\frac{SS_R/k}{SSE/(n-k-1)} \sim F(k, n-k-1)$.

Logistic regression: Let Y can only take 0 or 1 with $P(Y=1) = p$ and $P(Y=0) = 1-p$,

$$E(Y) = p(x_1, \dots, x_k) = \frac{e^{\beta_0 + \beta_1 x_1 + \dots + \beta_k x_k}}{1 + e^{\beta_0 + \beta_1 x_1 + \dots + \beta_k x_k}}.$$

• Y : response variable (which is Bernoulli r.v. $P(Y=1) = p$ and $P(Y=0) = 1-p$, so $E(Y) = p$), $\{x_1, \dots, x_k\}$: predictors (which are scalars).

• sample: $\{(x_{11}, \dots, x_{1k}; y_1), (x_{21}, \dots, x_{2k}; y_2), \dots, (x_{n1}, \dots, x_{nk}; y_n)\}$.

• how to estimate $\beta_j \approx \hat{\beta}_j$: maximal likelihood method (maximize $\prod_{i=1}^n p(x_{i1}, \dots, x_{ik})^{y_i} (1 - p(x_{i1}, \dots, x_{ik}))^{1-y_i}$).

• $\frac{\hat{\beta}_j - \beta_j}{s.e(\hat{\beta}_j)} \approx N(0, 1)$ for large $n \geq 30$, this helps determine whether or not the real $\beta_j = 0$?

• Classification of a new object $Y(x_1, \dots, x_k)$ as 1 or 0 according

$$Y(x_1, \dots, x_k) = \begin{cases} 1, & \text{if } \hat{p}(x_1, \dots, x_k) \geq 0.5, \\ 0, & \text{if } \hat{p}(x_1, \dots, x_k) < 0.5, \end{cases}$$

where the estimated logit function $\hat{p}(x_1, \dots, x_k)$ is

$$\hat{p}(x_1, \dots, x_k) = \frac{e^{\hat{\beta}_0 + \hat{\beta}_1 x_1 + \dots + \hat{\beta}_k x_k}}{1 + e^{\hat{\beta}_0 + \hat{\beta}_1 x_1 + \dots + \hat{\beta}_k x_k}}.$$

9. Tables

(9.1) Table for $N(0,1)$ standard normal random variable $\Phi(x) = P(N(0,1) \leq x)$, $x \geq 0$.

There is an important relation $\Phi(-x) = 1 - \Phi(x)$, $x \geq 0$.

x	0	1	2	3	4	5	6	7	8	9
0.0	0.5000	0.5040	0.5080	0.5120	0.5160	0.5199	0.5239	0.5279	0.5319	0.5359
0.1	0.5398	0.5438	0.5478	0.5517	0.5557	0.5596	0.5636	0.5675	0.5714	0.5753
0.2	0.5793	0.5832	0.5871	0.5910	0.5948	0.5987	0.6026	0.6064	0.6103	0.6141
0.3	0.6179	0.6217	0.6255	0.6293	0.6331	0.6368	0.6406	0.6443	0.6480	0.6517
0.4	0.6554	0.6591	0.6628	0.6664	0.6700	0.6736	0.6772	0.6808	0.6844	0.6879
0.5	0.6915	0.6950	0.6985	0.7019	0.7054	0.7088	0.7123	0.7157	0.7190	0.7224
0.6	0.7257	0.7291	0.7324	0.7357	0.7389	0.7422	0.7454	0.7486	0.7517	0.7549
0.7	0.7580	0.7611	0.7642	0.7673	0.7704	0.7734	0.7764	0.7794	0.7823	0.7852
0.8	0.7881	0.7910	0.7939	0.7957	0.8023	0.8051	0.8078	0.8106	0.8133	0.8160
0.9	0.8159	0.8186	0.8212	0.8238	0.8264	0.8289	0.8315	0.8340	0.8365	0.8389
1.0	0.8413	0.8438	0.8461	0.8485	0.8508	0.8531	0.8554	0.8577	0.8599	0.8621
1.1	0.8643	0.8665	0.8686	0.8708	0.8729	0.8749	0.8770	0.8790	0.8810	0.8830
1.2	0.8849	0.8869	0.8888	0.8907	0.8925	0.8944	0.8962	0.8980	0.8997	0.9015
1.3	0.9032	0.9049	0.9066	0.9082	0.9099	0.9115	0.9131	0.9147	0.9162	0.9177
1.4	0.9192	0.9222	0.9256	0.9251	0.9265	0.9279	0.9292	0.9306	0.9319	0.9330
1.5	0.9332	0.9345	0.9357	0.9370	0.9382	0.9394	0.9406	0.9418	0.9429	0.9441
1.6	0.9452	0.9463	0.9474	0.9484	0.9495	0.9505	0.9515	0.9525	0.9535	0.9545
1.7	0.9554	0.9564	0.9573	0.9582	0.9591	0.9599	0.9608	0.9616	0.9625	0.9633
1.8	0.9641	0.9649	0.9656	0.9664	0.9671	0.9678	0.9686	0.9693	0.9699	0.9706
1.9	0.9713	0.9719	0.9726	0.9732	0.9738	0.9744	0.9750	0.9756	0.9761	0.9767
2.0	0.9772	0.9778	0.9783	0.9788	0.9793	0.9798	0.9803	0.9808	0.9812	0.9817
2.1	0.9821	0.9826	0.9830	0.9834	0.9838	0.9842	0.9846	0.9850	0.9854	0.9857
2.2	0.9861	0.9864	0.9868	0.9871	0.9875	0.9878	0.9881	0.9884	0.9887	0.9890
2.3	0.9893	0.9896	0.9898	0.9901	0.9904	0.9906	0.9909	0.9911	0.9913	0.9916
2.4	0.9918	0.9920	0.9922	0.9925	0.9927	0.9929	0.9931	0.9932	0.9934	0.9936
2.5	0.9938	0.9940	0.9941	0.9943	0.9945	0.9946	0.9948	0.9949	0.9951	0.9952
2.6	0.9955	0.9956	0.9957	0.9959	0.9960	0.9961	0.9962	0.9963	0.9964	0.9965
2.7	0.9965	0.9966	0.9967	0.9968	0.9969	0.9970	0.9971	0.9972	0.9973	0.9974
2.8	0.9974	0.9975	0.9977	0.9978	0.9979	0.9979	0.9980	0.9981	0.9981	0.9981
2.9	0.9981	0.9982	0.9983	0.9984	0.9984	0.9985	0.9986	0.9986	0.9986	0.9986
3.0	0.9987	0.9987	0.9987	0.9988	0.9988	0.9989	0.9989	0.9990	0.9990	0.9990
3.1	0.9990	0.9991	0.9991	0.9991	0.9992	0.9992	0.9993	0.9993	0.9993	0.9993
3.2	0.9993	0.9993	0.9994	0.9994	0.9994	0.9994	0.9995	0.9995	0.9995	0.9995
3.3	0.9995	0.9995	0.9995	0.9996	0.9996	0.9996	0.9996	0.9996	0.9997	0.9997
3.4	0.9997	0.9997	0.9997	0.9997	0.9997	0.9997	0.9997	0.9997	0.9998	0.9998
3.5	0.9998	0.9998	0.9998	0.9998	0.9998	0.9998	0.9998	0.9998	0.9998	0.9998
3.6	0.9998	0.9998	0.9999	0.9999	0.9999	0.9999	0.9999	0.9999	0.9999	0.9999
3.7	0.9999	0.9999	0.9999	0.9999	0.9999	0.9999	0.9999	0.9999	0.9999	0.9999
3.8	0.9999	0.9999	0.9999	0.9999	0.9999	0.9999	0.9999	0.9999	0.9999	0.9999
3.9	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000
4.0	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000

(9.2) Table for $T(f)$ random variable $F(x) = P(T(f) \leq x)$, where f is a parameter called 'degrees of freedom'.

f	0.75	0.90	0.95	0.975	0.99	0.995	0.9975	0.9995	$F(x)$
1	1.00	3.08	6.31	12.71	31.82	63.66	127.32	636.62	
2	0.82	1.89	2.92	4.30	6.96	9.92	14.09	31.60	
3	0.76	1.64	2.35	3.18	4.54	5.84	7.45	12.92	
4	0.74	1.53	2.13	2.78	3.75	4.60	5.60	8.61	
5	0.73	1.48	2.02	2.57	3.36	4.03	4.77	6.87	
6	0.72	1.44	1.94	2.45	3.14	3.71	4.32	5.96	
7	0.71	1.41	1.89	2.36	3.00	3.50	4.03	5.41	
8	0.71	1.40	1.86	2.31	2.90	3.36	3.83	5.04	
9	0.70	1.38	1.83	2.26	2.82	3.25	3.69	4.78	
10	0.70	1.37	1.81	2.23	2.76	3.17	3.58	4.59	
11	0.70	1.36	1.80	2.20	2.72	3.11	3.50	4.44	
12	0.70	1.36	1.78	2.18	2.68	3.05	3.43	4.32	
13	0.69	1.35	1.77	2.16	2.65	3.01	3.37	4.22	
14	0.69	1.35	1.76	2.14	2.62	2.98	3.33	4.14	
15	0.69	1.34	1.75	2.13	2.60	2.95	3.29	4.07	
16	0.69	1.34	1.75	2.12	2.58	2.92	3.25	4.01	
17	0.69	1.33	1.74	2.11	2.57	2.90	3.22	3.97	
18	0.69	1.33	1.73	2.10	2.55	2.88	3.20	3.92	
19	0.69	1.33	1.73	2.09	2.54	2.86	3.17	3.88	
20	0.69	1.33	1.72	2.09	2.53	2.85	3.15	3.85	
21	0.69	1.32	1.72	2.08	2.52	2.83	3.14	3.82	
22	0.69	1.32	1.72	2.07	2.51	2.82	3.12	3.79	
23	0.69	1.32	1.71	2.07	2.50	2.81	3.10	3.77	
24	0.68	1.32	1.71	2.06	2.49	2.80	3.09	3.75	
25	0.68	1.32	1.71	2.06	2.49	2.79	3.08	3.73	
26	0.68	1.31	1.71	2.06	2.48	2.78	3.07	3.71	
27	0.68	1.31	1.70	2.05	2.47	2.77	3.06	3.69	
28	0.68	1.31	1.70	2.05	2.47	2.76	3.05	3.67	
29	0.68	1.31	1.70	2.05	2.46	2.76	3.04	3.66	
30	0.68	1.31	1.70	2.04	2.46	2.75	3.03	3.65	
40	0.68	1.30	1.68	2.02	2.42	2.70	2.97	3.55	
50	0.68	1.30	1.68	2.01	2.40	2.68	2.94	3.50	
60	0.68	1.30	1.67	2.00	2.39	2.66	2.91	3.46	
100	0.68	1.29	1.66	1.98	2.36	2.63	2.87	3.39	
∞	0.67	1.28	1.65	1.96	2.33	2.58	2.81	3.29	

(9.3) Table for $\chi^2(f)$ random variable $F(x) = P(\chi^2(f) \leq x)$, where f is a parameter.

f	$F(x)$										
	0.0005	0.001	0.005	0.01	0.025	0.05	0.10	0.20	0.30	0.40	0.50
1	0.00	0.00	0.00	0.00	0.00	0.02	0.06	0.15	0.27	0.45	1
2	0.00	0.00	0.01	0.02	0.05	0.10	0.21	0.45	0.71	1.39	2
3	0.02	0.02	0.07	0.11	0.22	0.35	0.58	1.01	1.42	2.37	3
4	0.06	0.09	0.21	0.30	0.48	0.71	1.06	1.65	2.19	2.75	4
5	0.16	0.21	0.41	0.55	0.83	1.15	1.61	2.34	3.00	3.66	5
6	0.30	0.38	0.68	0.87	1.24	1.64	2.20	3.07	3.83	4.57	6
7	0.48	0.60	0.99	1.24	1.69	2.17	2.83	3.82	4.67	5.49	7
8	0.71	0.86	1.34	1.65	2.18	2.73	3.49	4.59	5.53	6.42	8
9	0.97	1.15	1.73	2.09	2.70	3.33	4.17	5.38	6.39	7.36	9
10	1.26	1.48	2.16	2.56	3.25	3.94	4.87	6.18	7.27	8.30	10
11	1.59	1.83	2.60	3.05	3.82	4.57	5.58	6.99	8.15	9.24	11
12	1.93	2.21	3.07	3.57	4.40	5.23	6.30	7.81	9.03	10.18	12
13	2.31	2.62	3.57	4.11	5.01	5.89	7.04	8.63	9.93	11.13	13
14	2.70	3.04	4.07	4.66	5.63	6.57	7.79	9.47	10.82	12.08	14
15	3.11	3.48	4.60	5.23	6.26	7.26	8.55	10.31	11.72	13.03	15
16	3.54	3.94	5.14	5.81	6.91	7.96	9.31	11.15	12.62	13.98	16
17	3.98	4.42	5.70	6.41	7.56	8.67	10.09	12.00	13.53	14.94	17
18	4.44	4.90	6.26	7.01	8.23	9.39	10.86	12.86	14.44	15.89	18
19	4.91	5.41	6.84	7.63	8.91	10.12	11.65	13.72	15.35	16.85	19
20	5.40	5.92	7.43	8.26	9.59	10.85	12.44	14.58	16.27	17.81	20
21	5.90	6.45	8.03	8.90	10.28	11.59	13.24	15.44	17.18	18.77	21
22	6.40	6.98	8.64	9.54	10.98	12.34	14.04	16.31	18.10	19.73	22
23	6.92	7.53	9.26	10.20	11.69	13.09	14.85	17.19	19.02	20.69	23
24	7.45	8.08	9.89	10.86	12.40	13.85	15.66	18.06	19.94	21.65	24
25	7.99	8.65	10.52	11.52	13.12	14.61	16.47	18.94	20.87	22.62	25
26	8.54	9.22	11.16	12.20	13.84	15.38	17.29	19.82	21.79	23.58	26
27	9.09	9.80	11.81	12.88	14.57	16.15	18.11	20.70	22.72	24.54	27
28	9.66	10.39	12.46	13.56	15.31	16.93	18.94	21.59	23.65	25.51	28
29	10.23	10.99	13.12	14.26	16.05	17.71	19.77	22.48	24.58	26.48	29
30	10.80	11.59	13.79	14.95	16.79	18.49	20.60	23.36	25.51	27.44	30
40	16.91	17.92	20.71	22.16	24.43	26.51	29.05	32.34	34.87	37.13	40
50	23.46	24.67	27.99	29.71	32.36	34.76	37.69	41.45	44.31	46.86	50
60	30.34	31.74	35.53	37.48	40.48	43.19	46.46	50.64	53.81	56.62	60
100	59.90	61.92	67.33	70.06	74.22	77.93	82.36	87.95	92.13	95.81	100

Table for $\chi^2(f)$ random variable $F(x) = P(\chi^2(f) \leq x)$, where f is a parameter.

f	$F(x)$									
	0.60	0.70	0.80	0.90	0.95	0.975	0.99	0.995	0.999	0.9995
1	0.71	1.07	1.64	2.71	3.84	5.02	6.63	7.88	10.83	12.12
2	1.83	2.41	3.22	4.61	5.99	7.38	9.21	10.60	13.82	15.20
3	2.95	3.66	4.64	6.25	7.81	9.35	11.34	12.84	16.27	17.73
4	4.04	4.88	5.99	7.78	9.49	11.14	13.28	14.86	18.47	20.00
5	5.13	6.06	7.29	9.24	11.07	12.83	15.09	16.75	20.52	22.11
6	6.21	7.23	8.56	10.64	12.59	14.45	16.81	18.55	22.46	24.10
7	7.28	8.38	9.80	12.02	14.07	16.01	18.48	20.28	24.32	26.02
8	8.35	9.52	11.03	13.36	15.51	17.53	20.09	21.95	26.12	27.87
9	9.41	10.66	12.24	14.68	16.92	19.02	21.67	23.59	27.88	29.67
10	10.47	11.78	13.44	15.99	18.31	20.48	23.21	25.19	29.59	31.42
11	11.53	12.90	14.63	17.28	19.68	21.92	24.72	26.76	31.14	33.14
12	12.58	14.01	15.81	18.55	21.03	23.34	26.22	28.30	32.91	34.82
13	13.64	15.12	16.98	19.81	22.36	24.74	27.69	29.82	34.53	36.48
14	14.69	16.22	18.15	21.06	23.68	26.12	29.14	31.32	36.12	38.11
15	15.73	17.32	19.31	22.31	25.00	27.49	30.58	32.80	37.70	39.79
16	16.78	18.42	20.47	23.54	26.30	28.85	32.00	34.27	39.25	41.31
17	17.82	19.51	21.61	24.77	27.59	30.19	33.41	35.72	40.79	42.88
18	18.87	20.60	22.76	25.99	28.87	31.53	34.81	37.16	42.31	44.43
19	19.91	21.69	23.90	27.20	30.14	32.85	36.19	38.58	43.82	45.97
20	20.95	22.77	25.04	28.41	31.41	34.17	37.57	40.00	45.31	47.50
21	21.99	23.86	26.17	29.62	32.67	35.48	38.93	41.40	46.80	49.01
22	23.03	24.94	27.30	30.81	33.92	36.78	40.29	42.80	48.27	50.51
23	24.07	26.02	28.43	32.01	35.17	38.08	41.64	44.18	49.73	52.00
24	25.11	27.10	29.55	33.20	36.42	39.36	42.98	45.56	51.18	53.48
25	26.14	28.17	30.68	34.38	37.65	40.65	44.31	46.93	52.62	54.95
26	27.18	29.25	31.79	35.56	38.89	41.92	45.64	48.29	54.05	56.41
27	28.21	30.32	32.91	36.74	40.11	43.19	46.96	49.64	55.48	57.86
28	29.25	31.39	34.03	37.92	41.34	44.46	48.28	50.99	56.89	59.30
29	30.28	32.46	35.14	39.09	42.56	45.72	49.59	52.34	58.30	60.73
30	31.32	33.53	36.25	40.26	43.77	46.98	50.89	53.67	59.70	62.16
40	41.62	44.16	47.27	51.81	55.76	59.34	63.69	66.77	73.40	76.09
50	51.89	54.72	58.16	63.17	67.50	71.42	76.15	79.49	86.66	89.56
60	62.13	65.23	68.97	74.40	79.08	83.30	88.38	91.95	99.61	102.69
100	102.95	106.91	111.67	118.50	124.34	129.56	135.81	140.17	149.45	153.17

(9.4) Table for Binomial random variable $P(Bin(n, p) \leq k)$ if $p \leq 0.5$.
 If $p > 0.5$, then $P(Bin(n, p) \leq k) = P(Bin(n, 1-p) \geq n-k)$.

n	k	0.05	0.10	0.15	0.20	0.25	0.30	0.35	0.40	0.45	0.50	p
2	0	0.9025	0.8100	0.7225	0.6400	0.5625	0.4900	0.4225	0.3600	0.3025	0.2500	
3	0	0.9575	0.9900	0.9775	0.9600	0.9375	0.9100	0.8775	0.8400	0.7975	0.7500	
4	0	0.8574	0.7290	0.6141	0.5120	0.4219	0.3430	0.2746	0.2160	0.1664	0.1250	
5	0	0.9928	0.9720	0.9392	0.8960	0.8438	0.7840	0.7183	0.6480	0.5747	0.5000	
6	0	0.9999	0.9990	0.9966	0.9920	0.9844	0.9730	0.9571	0.9360	0.9089	0.8750	
7	0	0.8145	0.6561	0.5220	0.4096	0.3164	0.2401	0.1785	0.1296	0.0915	0.0625	
8	0	0.9860	0.9477	0.8905	0.8192	0.7383	0.6517	0.5630	0.4752	0.3910	0.3125	
9	0	0.9995	0.9963	0.9880	0.9728	0.9492	0.9163	0.8735	0.8208	0.7585	0.6875	
10	0	0.7738	0.5905	0.4437	0.3277	0.2373	0.1681	0.1160	0.0778	0.0503	0.0313	
11	0	0.9774	0.9185	0.8352	0.7373	0.6328	0.5282	0.4284	0.3370	0.2562	0.1875	
12	0	0.9988	0.9914	0.9734	0.9421	0.8965	0.8369	0.7648	0.6826	0.5931	0.5000	
13	0	0.9999	0.9995	0.9978	0.9933	0.9844	0.9692	0.9460	0.9130	0.8688	0.8125	
14	0	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	
15	0	0.7351	0.5314	0.3771	0.2621	0.1780	0.1176	0.0754	0.0467	0.0277	0.0156	
16	0	0.9672	0.8857	0.7765	0.6554	0.5339	0.4202	0.3191	0.2333	0.1636	0.1094	
17	0	0.9978	0.9842	0.9527	0.9011	0.8306	0.7443	0.6471	0.5443	0.4415	0.3438	
18	0	0.9999	0.9987	0.9941	0.9830	0.9624	0.9295	0.8826	0.8208	0.7447	0.6563	
19	0	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	
20	0	0.6983	0.4783	0.3206	0.2097	0.1335	0.0824	0.0490	0.0280	0.0152	0.0078	
21	0	0.9556	0.8503	0.7166	0.5767	0.4449	0.3294	0.2338	0.1586	0.1024	0.0625	
22	0	0.9962	0.9743	0.9262	0.8520	0.7564	0.6471	0.5323	0.4199	0.3124	0.2266	
23	0	0.9998	0.9973	0.9879	0.9667	0.9294	0.8740	0.8002	0.7102	0.6083	0.5000	
24	0	1.0000	0.9998	0.9988	0.9953	0.9871	0.9712	0.9444	0.9037	0.8471	0.7734	
25	0	1.0000	1.0000	0.9999	0.9996	0.9987	0.9962	0.9910	0.9812	0.9643	0.9375	
26	0	1.0000	1.0000	1.0000	0.9999	0.9998	0.9994	0.9984	0.9963	0.9922	0.9900	
27	0	0.6634	0.4305	0.2725	0.1678	0.1001	0.0576	0.0319	0.0168	0.0084	0.0039	
28	0	1.0000	0.9428	0.8131	0.6572	0.5033	0.3671	0.2553	0.1691	0.1064	0.0632	
29	0	2.0000	1.0000	0.9996	0.9950	0.9786	0.8862	0.8059	0.7064	0.5941	0.4770	
30	0	3.0000	1.0000	0.9996	0.9971	0.9896	0.9727	0.9420	0.8939	0.8263	0.7396	
31	0	4.0000	1.0000	0.9998	0.9988	0.9987	0.9747	0.9502	0.9115	0.8555	0.7733	
32	0	5.0000	1.0000	0.9998	0.9998	0.9987	0.9964	0.9915	0.9819	0.9643	0.9000	
33	0	6.0000	1.0000	0.9999	0.9999	0.9987	0.9964	0.9915	0.9819	0.9643	0.9000	
34	0	7.0000	1.0000	1.0000	1.0000	0.9999	0.9998	0.9993	0.9983	0.9961	0.9900	
35	0	8.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	
36	0	9.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	
37	0	10.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	
38	0	11.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	
39	0	12.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	

Table for Binomial random variable $P(Bin(n, p) \leq k)$ if $p \leq 0.5$.
 If $p > 0.5$, then $P(Bin(n, p) \leq k) = P(Bin(n, 1-p) \geq n-k)$.

n	k	0.05	0.10	0.15	0.20	0.25	0.30	0.35	0.40	0.45	0.50	p
11	0	0.5688	0.3138	0.1673	0.0859	0.0422	0.0198	0.0088	0.0036	0.0014	0.0005	
12	0	0.5404	0.2824	0.1422	0.0687	0.0317	0.0138	0.0057	0.0022	0.0008	0.0002	
13	0	0.5040	0.2242	0.1087	0.0544	0.0274	0.0124	0.0055	0.0023	0.0009	0.0003	
14	0	0.4816	0.1659	0.0843	0.0443	0.0274	0.0158	0.0085	0.0042	0.0016	0.0003	
15	0	0.4540	0.1122	0.0567	0.0317	0.0187	0.0095	0.0047	0.0023	0.0009	0.0003	
16	0	0.4204	0.0784	0.0419	0.0242	0.0137	0.0070	0.0034	0.0016	0.0006	0.0002	
17	0	0.3864	0.0530	0.0287	0.0153	0.0080	0.0042	0.0021	0.0009	0.0004	0.0001	
18	0	0.3530	0.0375	0.0197	0.0103	0.0053	0.0027	0.0013	0.0006	0.0003	0.0001	
19	0	0.3226	0.0267	0.0137	0.0070	0.0035	0.0017	0.0008	0.0004	0.0002	0.0001	
20	0	0.2923	0.0198	0.0103	0.0053	0.0027	0.0013	0.0006	0.0003	0.0002	0.0001	
21	0	0.2620	0.0146	0.0070	0.0035	0.0017	0.0008	0.0004	0.0002	0.0001	0.0001	
22	0	0.2326	0.0103	0.0053	0.0027	0.0013	0.0006	0.0003	0.0002	0.0001	0.0001	
23	0	0.2025	0.0070	0.0035	0.0017	0.0008	0.0004	0.0002	0.0001	0.0001	0.0001	
24	0	0.1720	0.0046	0.0020	0.0010	0.0005	0.0002	0.0001	0.0001	0.0001	0.0001	
25	0	0.1420	0.0023	0.0010	0.0005	0.0002	0.0001	0.0001	0.0001	0.0001	0.0001	
26	0	0.1120	0.0017	0.0007	0.0003	0.0001	0.0001	0.0001	0.0001	0.0001	0.0001	
27	0	0.0820	0.0013	0.0005	0.0002	0.0001	0.0001	0.0001	0.0001	0.0001	0.0001	
28	0	0.0520	0.0009	0.0004	0.0002	0.0001	0.0001	0.0001	0.0001	0.0001	0.0001	
29	0	0.0220	0.0005	0.0002	0.0001	0.0001	0.0001	0.0001	0.0001	0.0001	0.0001	
30	0	0.0020	0.0002	0.0001	0.0001	0.0001	0.0001	0.0001	0.0001	0.0001	0.0001	
31	0	0.0009	0.0001	0.0001	0.0001	0.0001	0.0001	0.0001	0.0001	0.0001	0.0001	
32	0	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	
33	0	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	
34	0	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	
35	0	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	
36	0	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	
37	0	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	
38	0	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	
39	0	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	

Table for Binomial random variable $P(Bin(n, p) \leq k)$ if $p \leq 0.5$.
If $p > 0.5$, then $P(Bin(n, p) \leq k) = P(Bin(n, 1-p) \geq n-k)$.

n	k	0.05	0.10	0.15	0.20	0.25	p	0.30	0.35	0.40	0.45	0.50
14	0	0.4877	0.2288	0.1028	0.0440	0.0178	0.0068	0.0024	0.0008	0.0002	0.0001	
1	0.8470	0.5284	0.3567	0.1979	0.1010	0.0475	0.0205	0.0081	0.0009			
2	0.9699	0.8416	0.6479	0.4481	0.2811	0.1608	0.0839	0.0398	0.0170	0.0065		
3	0.9558	0.9559	0.8535	0.6982	0.5213	0.3552	0.2205	0.1243	0.0632	0.0287	0.0123	
4	0.9996	0.9908	0.9533	0.8702	0.7415	0.5842	0.4227	0.2793	0.1672	0.0898	0.0464	
5	1.0000	0.9985	0.9885	0.9561	0.8883	0.7805	0.6405	0.4859	0.3373	0.2120	0.1260	0.0596
6	1.0000	0.9998	0.9978	0.9884	0.9617	0.9067	0.8164	0.6925	0.5461	0.3953	0.2348	0.0245
7	1.0000	1.0000	0.9997	0.9976	0.9897	0.9685	0.9247	0.8499	0.7414	0.6047	0.4478	0.2902
8	1.0000	1.0000	1.0000	0.9996	0.9978	0.9917	0.9757	0.8811	0.7830	0.6405	0.4743	0.3145
9	1.0000	1.0000	1.0000	1.0000	0.9997	0.9983	0.9940	0.9825	0.9574	0.9102	0.8011	0.6626
10	1.0000	1.0000	1.0000	1.0000	1.0000	0.9998	0.9989	0.9961	0.9886	0.9713	0.9177	0.6855
11	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	0.9999	0.9978	0.9935	0.9777	0.8338
12	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	0.9894	0.9283
13	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	0.9755
14	0	0.4633	0.2059	0.0874	0.0352	0.0134	0.0047	0.0016	0.0005	0.0001	0.0000	
1	0.8290	0.5490	0.3186	0.1671	0.0802	0.0353	0.0142	0.0052	0.0017	0.0005		
2	0.9638	0.8159	0.6042	0.3980	0.2361	0.1268	0.0617	0.0271	0.0107	0.0037		
3	0.9445	0.9444	0.8227	0.6482	0.4613	0.2969	0.1727	0.0905	0.0424	0.0176		
4	0.9994	0.9873	0.9383	0.8358	0.6865	0.5155	0.3519	0.2173	0.1204	0.0592		
5	0.9999	0.9978	0.9832	0.9389	0.8516	0.7216	0.5643	0.4032	0.2608	0.1509		
6	1.0000	0.9997	0.9964	0.9819	0.9434	0.8689	0.7548	0.6098	0.4522	0.3036		
7	1.0000	1.0000	0.9994	0.9958	0.9827	0.9500	0.8868	0.7869	0.6535	0.5000		
8	1.0000	1.0000	0.9999	0.9992	0.9958	0.9848	0.9578	0.9050	0.8182	0.6964		
9	1.0000	1.0000	1.0000	0.9990	0.9992	0.9963	0.9876	0.9662	0.9231	0.8491		
10	1.0000	1.0000	1.0000	0.9999	0.9993	0.9972	0.9907	0.9745	0.9408	0.8743		
11	1.0000	1.0000	1.0000	1.0000	0.9999	0.9995	0.9981	0.9937	0.9824	0.8809		
12	1.0000	1.0000	1.0000	1.0000	1.0000	0.9999	0.9997	0.9989	0.9963	0.9790		
13	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	0.9999	0.9995	0.9998		
14	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000		
15	0	0.4401	0.1853	0.0743	0.0281	0.0100	0.0033	0.0010	0.0003	0.0001	0.0000	
1	0.8108	0.5147	0.2839	0.1407	0.0635	0.0261	0.0098	0.0033	0.0010	0.0003		
2	0.9571	0.7892	0.5614	0.3518	0.1971	0.0994	0.0451	0.0183	0.0066	0.0021		
3	0.9360	0.9316	0.7899	0.5981	0.4050	0.2459	0.1339	0.0651	0.0281	0.0106		
4	0.9991	0.9830	0.9209	0.7982	0.6302	0.4499	0.2892	0.1666	0.0853	0.0384		
5	0.9999	0.9967	0.9765	0.9183	0.8103	0.6598	0.4900	0.3288	0.1976	0.1051		
6	1.0000	0.9995	0.9944	0.9733	0.9204	0.8247	0.6881	0.5272	0.3660	0.2272		
7	1.0000	0.9999	0.9930	0.9729	0.9256	0.8406	0.7161	0.5629	0.4018	0.2409		
8	1.0000	1.0000	0.9998	0.9985	0.9795	0.9743	0.9329	0.8579	0.7441	0.5982		
9	1.0000	1.0000	1.0000	0.9998	0.9924	0.9771	0.9417	0.8759	0.7728	0.6454		
10	1.0000	1.0000	1.0000	0.9997	0.9984	0.9938	0.9809	0.9514	0.8949	0.7439		
11	1.0000	1.0000	1.0000	1.0000	0.9997	0.9987	0.9951	0.9851	0.9616	0.8180		
12	1.0000	1.0000	1.0000	1.0000	1.0000	0.9998	0.9991	0.9965	0.9894	0.8988		
13	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000		
14	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000		
15	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000		

Table for Binomial random variable $P(Bin(n, p) \leq k)$ if $p \leq 0.5$.
If $p > 0.5$, then $P(Bin(n, p) \leq k) = P(Bin(n, 1-p) \geq n-k)$.

n	k	0.05	0.10	0.15	0.20	0.25	p	0.30	0.35	0.40	0.45	0.50
17	0	0.4181	0.1668	0.0631	0.0225	0.0075	0.0023	0.0007	0.0002	0.0000	0.0000	
1	0.7922	0.4818	0.2525	0.1182	0.0501	0.0193	0.0067	0.0021	0.0006	0.0001		
2	0.9497	0.7618	0.5198	0.3096	0.1637	0.0774	0.0327	0.0123	0.0041	0.0012		
3	0.9912	0.9174	0.7556	0.5489	0.3530	0.2019	0.1028	0.0464	0.0184	0.0064		
4	0.9988	0.9779	0.9013	0.7582	0.5739	0.3887	0.2348	0.1260	0.0596	0.0245		
5	0.9999	0.9953	0.9681	0.8943	0.7653	0.5968	0.4197	0.2639	0.1471	0.0717		
6	1.0000	0.9992	0.9623	0.8929	0.7752	0.6188	0.4478	0.2902	0.1662			
7	1.0000	0.9999	0.9883	0.9598	0.8787	0.7872	0.6405	0.4743	0.3145			
8	1.0000	1.0000	0.9997	0.9974	0.9876	0.9597	0.9006	0.8011	0.6626	0.5000		
9	1.0000	1.0000	1.0000	0.9999	0.9999	0.9970	0.9894	0.9699	0.9283	0.7597		
10	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000		
11	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000		
12	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000		
13	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000		
14	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000		
15	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000		
16	0	0.3774	0.1351	0.0456	0.0144	0.0042	0.0011	0.0003	0.0001	0.0000		
1	0.7547	0.4203	0.1985	0.0829	0.0310	0.0104	0.0031	0.0008	0.0002	0.0000		
2	0.9335	0.7054	0.4413	0.2369	0.1113	0.0462	0.0170	0.0055	0.0015	0.0004		
3	0.9836	0.8850	0.6841	0.4551	0.2631	0.1332	0.0591	0.0230	0.0077	0.0022		
4	0.9980	0.9648	0.8556	0.6733	0.4654	0.2822	0.1500	0.0696	0.0280	0.0096		
5	0.9998	0.9914	0.9463	0.8369	0.6678	0.4739	0.2968	0.1629	0.0777	0.0318		
6	1.0000	0.9983	0.9837	0.9324	0.8251	0.6655	0.4812	0.3081	0.1727	0.0835		
7	1.0000	0.9997	0.9959	0.9767	0.9225	0.8180	0.6656	0.4878	0.3169	0.1796		
8	1.0000	1.0000	0.9992	0.9933	0.9713	0.9161	0.8145	0.6675	0.4940	0.3238		
9	1.0000	1.0000	0.9999	0.9974	0.9817	0.9077	0.8159	0.6710	0.4915	0.3238		
10	1.0000	1.0000	1.0000	0.9997	0.9977	0.9895	0.9653	0.9115	0.8159	0.6762		
11	1.0000	1.0000	1.0000	0.9995	0.9972	0.9886	0.9648	0.9129	0.8204	0.6822		
12	1.0000	1.0000	1.0000	0.9999	0.9994	0.9969	0.9884	0.9658	0.9165	0.7914		
13	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000		
14	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000		
15	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000		
16	0	0.3774	0.1351	0.0456	0.0144	0.0042	0.0011	0.0003	0.0001	0.0000		

(9.5) Table for Poisson random variable $P(Po(\mu) \leq k)$.

k	μ									
k	0.1	0.2	0.3	0.4	0.5	0.6	0.7	0.8	0.9	1.0
0	0.9048	0.8187	0.7408	0.6703	0.6065	0.5488	0.4966	0.4493	0.4066	0.3679
1	0.9953	0.9825	0.9631	0.9384	0.9098	0.8781	0.8442	0.8088	0.7725	0.7358
2	0.9998	0.9989	0.9964	0.9921	0.9856	0.9769	0.9659	0.9526	0.9371	0.9197
3	1.0000	0.9999	0.9997	0.9992	0.9982	0.9966	0.9942	0.9909	0.9865	0.9810
4	1.0000	1.0000	0.9999	0.9998	0.9996	0.9992	0.9986	0.9977	0.9963	0.9956
5	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	0.9999	0.9998	0.9994
6	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000
7	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000
k	1.1	1.2	1.3	1.4	1.5	1.6	1.7	1.8	1.9	2.0
0	0.3329	0.3012	0.2725	0.2466	0.2231	0.2019	0.1827	0.1653	0.1496	0.1353
1	0.6890	0.6626	0.6268	0.5918	0.5578	0.5249	0.4932	0.4628	0.4337	0.4060
2	0.904	0.8795	0.8571	0.8335	0.8088	0.7834	0.7572	0.7306	0.7037	0.6767
3	0.9743	0.9662	0.9569	0.9463	0.9344	0.9212	0.9068	0.8913	0.8747	0.8571
4	0.9946	0.9923	0.9893	0.9857	0.9814	0.9763	0.9704	0.9636	0.9559	0.9473
5	0.9990	0.9985	0.9978	0.9968	0.9955	0.9940	0.9920	0.9896	0.9834	0.9775
6	0.9999	0.9997	0.9996	0.9994	0.9991	0.9987	0.9981	0.9974	0.9966	0.9955
7	1.0000	1.0000	0.9999	0.9999	0.9998	0.9997	0.9996	0.9994	0.9992	0.9989
8	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000
9	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000
k	3.2	3.4	3.6	3.8	4.0	4.2	4.4	4.6	4.8	5.0
0	0.0408	0.0334	0.0273	0.0224	0.0183	0.0150	0.0123	0.0101	0.0082	0.0067
1	0.1712	0.1468	0.1257	0.1074	0.0916	0.0780	0.0663	0.0563	0.0477	0.0404
2	0.3799	0.3397	0.3027	0.2689	0.2381	0.2102	0.1851	0.1626	0.1425	0.1247
3	0.6025	0.5584	0.5152	0.4735	0.4335	0.3954	0.3594	0.3257	0.2942	0.2650
4	0.7806	0.7442	0.7064	0.6678	0.6288	0.5898	0.5512	0.5132	0.4763	0.4405
5	0.8946	0.8705	0.8441	0.8156	0.7851	0.7531	0.7199	0.6858	0.6510	0.6160
6	0.9554	0.9421	0.9267	0.9091	0.8893	0.8675	0.8436	0.8180	0.7908	0.7622
7	0.9832	0.9769	0.9692	0.9559	0.9489	0.9361	0.9214	0.9049	0.8867	0.8666
8	0.9943	0.9917	0.9883	0.9840	0.9786	0.9721	0.9642	0.9549	0.9442	0.9319
9	0.9995	0.9992	0.9987	0.9981	0.9972	0.9959	0.9943	0.9922	0.9896	0.9863
k	5.2	5.4	5.6	5.8	6.0	6.5	7.0	7.5	8.0	8.5
0	0.0055	0.0045	0.0037	0.0030	0.0025	0.0015	0.0009	0.0006	0.0003	0.0002
1	0.0342	0.0289	0.0244	0.0206	0.0174	0.0113	0.0073	0.0047	0.0030	0.0019
2	0.1088	0.0948	0.0824	0.0715	0.0620	0.0430	0.0296	0.0203	0.0138	0.0093
3	0.2381	0.2133	0.1906	0.1700	0.1512	0.1118	0.0818	0.0591	0.0424	0.0301
4	0.4061	0.3733	0.3422	0.3127	0.2851	0.2237	0.1730	0.1321	0.0966	0.0744
5	0.5809	0.5461	0.5119	0.4783	0.4457	0.3690	0.3007	0.2414	0.1912	0.1496
6	0.7324	0.7017	0.6703	0.6384	0.6063	0.5265	0.4497	0.3782	0.3134	0.2562
7	0.8449	0.8149	0.7970	0.7710	0.7440	0.6728	0.5987	0.5246	0.4530	0.3856
8	0.9181	0.9027	0.8857	0.8672	0.8472	0.7916	0.7291	0.6620	0.5925	0.5231
9	0.9603	0.9512	0.9409	0.9292	0.9161	0.8774	0.8305	0.7764	0.7166	0.6530
10	0.9823	0.9775	0.9718	0.9651	0.9574	0.9332	0.9015	0.8622	0.8159	0.7634
11	0.9904	0.9875	0.9841	0.9799	0.9761	0.9467	0.9208	0.8881	0.8487	0.8047
12	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000
k	2.1	2.2	2.3	2.4	2.5	2.6	2.7	2.8	2.9	3.0
0	0.1225	0.1108	0.1003	0.0907	0.0821	0.0743	0.0672	0.0608	0.0550	0.0498
1	0.3796	0.3546	0.3309	0.3084	0.2873	0.2674	0.2487	0.2311	0.2146	0.1991
2	0.6496	0.6227	0.5960	0.5597	0.5348	0.5184	0.4936	0.4695	0.4460	0.4232
3	0.8386	0.8194	0.7787	0.7576	0.7360	0.7141	0.6919	0.6696	0.6474	0.6141
4	0.9379	0.9275	0.9162	0.9041	0.8912	0.8774	0.8477	0.8318	0.8153	0.7894
5	0.9796	0.9751	0.9700	0.9643	0.9580	0.9510	0.9433	0.9349	0.9258	0.9161
6	0.9941	0.9925	0.9884	0.9828	0.9794	0.9756	0.9713	0.9665	0.9623	0.9573
7	0.9985	0.9974	0.9967	0.9958	0.9947	0.9934	0.9919	0.9901	0.9881	0.9851
8	0.9997	0.9995	0.9994	0.9989	0.9985	0.9981	0.9976	0.9969	0.9962	0.9952
9	0.9999	0.9999	0.9999	0.9998	0.9997	0.9996	0.9995	0.9994	0.9993	0.9992
10	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000
11	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000
12	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000

Table for Poisson random variable $P(Po(\mu) \leq k)$.

k	9.0	9.5	10.0	11.0	12.0	13.0	14.0	15.0	16.0	17.0	μ
0	0.0001	0.0001	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000
1	0.0012	0.0008	0.0005	0.0002	0.0001	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000
2	0.0062	0.0042	0.0028	0.0012	0.0005	0.0002	0.0001	0.0000	0.0000	0.0000	0.0000
3	0.0212	0.0149	0.0103	0.0049	0.0023	0.0011	0.0005	0.0002	0.0001	0.0000	0.0000
4	0.0550	0.0403	0.0293	0.0151	0.0076	0.0037	0.0018	0.0009	0.0004	0.0002	0.0000
5	0.1157	0.0885	0.0671	0.0375	0.0203	0.0107	0.0055	0.0028	0.0014	0.0007	0.0000
6	0.2068	0.1649	0.1301	0.0786	0.0458	0.0259	0.0142	0.0076	0.0040	0.0021	0.0000
7	0.3239	0.2687	0.2202	0.1432	0.0895	0.0540	0.0316	0.0180	0.0100	0.0054	0.0000
8	0.4557	0.3918	0.3328	0.2320	0.1550	0.0998	0.0621	0.0374	0.0220	0.0126	0.0000
9	0.5874	0.5218	0.4579	0.3405	0.2424	0.1658	0.1094	0.0699	0.0433	0.0261	0.0000
10	0.7060	0.6453	0.5830	0.4599	0.3472	0.2517	0.1757	0.1185	0.0774	0.0491	0.0000
11	0.8030	0.7520	0.6968	0.5793	0.4616	0.3532	0.2600	0.1848	0.1270	0.0847	0.0000
12	0.8758	0.8364	0.7916	0.6887	0.5760	0.4631	0.3585	0.2676	0.1931	0.1350	0.0000
13	0.9261	0.8981	0.8645	0.7813	0.6815	0.5730	0.4644	0.3632	0.2745	0.2009	0.0000
14	0.9585	0.9400	0.9165	0.8540	0.7720	0.6751	0.5704	0.4657	0.3675	0.2808	0.0000
15	0.9780	0.9665	0.9513	0.9074	0.8444	0.7636	0.6694	0.5681	0.4667	0.3715	0.0000
16	0.9889	0.9823	0.9730	0.9441	0.8987	0.8355	0.7559	0.6641	0.5660	0.4677	0.0000
17	0.9947	0.9911	0.9857	0.9678	0.9370	0.8905	0.8272	0.7489	0.6593	0.5640	0.0000
18	0.9976	0.9957	0.9928	0.9823	0.9626	0.9302	0.8826	0.8195	0.7423	0.6550	0.0000
19	0.9989	0.9980	0.9965	0.9907	0.9787	0.9573	0.9235	0.8752	0.8122	0.7363	0.0000
20	0.9996	0.9991	0.9984	0.9953	0.9884	0.9750	0.9521	0.9170	0.8682	0.8055	0.0000
21	0.9998	0.9996	0.9993	0.9977	0.9939	0.9859	0.9712	0.9469	0.9108	0.8615	0.0000
22	0.9999	0.9999	0.9997	0.9990	0.9970	0.9924	0.9833	0.9673	0.9418	0.9047	0.0000
23	1.0000	0.9999	0.9999	0.9995	0.9985	0.9960	0.9907	0.9805	0.9633	0.9367	0.0000
24	1.0000	1.0000	1.0000	0.9998	0.9993	0.9980	0.9950	0.9888	0.9777	0.9594	0.0000
25	1.0000	1.0000	1.0000	0.9999	0.9997	0.9974	0.9938	0.9869	0.9748	0.9548	0.0000
26	1.0000	1.0000	1.0000	1.0000	0.9999	0.9995	0.9987	0.9967	0.9925	0.9848	0.0000
27	1.0000	1.0000	1.0000	1.0000	0.9999	0.9998	0.9994	0.9983	0.9959	0.9912	0.0000
28	1.0000	1.0000	1.0000	1.0000	0.9999	0.9997	0.9991	0.9978	0.9950	0.9900	0.0000
29	1.0000	1.0000	1.0000	1.0000	1.0000	0.9999	0.9996	0.9989	0.9973	0.9947	0.0000
30	1.0000	1.0000	1.0000	1.0000	1.0000	0.9999	0.9998	0.9994	0.9986	0.9976	0.0000
31	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	0.9999	0.9997	0.9993	0.9980	0.0000
32	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	0.9999	0.9996	0.0000
33	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	0.9999	0.9998	0.0000
34	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	0.9999	0.0000
35	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	0.0000