



Section: ALS Protocols
Subject: PAIN MANAGEMENT
Section #: 345.17
Issue Date: March 21, 2011
Revision Date: December 1, 2017
Approved By: 

Page 1 of 2

Michael Lozano, Jr., M.D., HCFR Medical Director

1. Intent:
 - a. Whereas many patients treated by HCFR personnel will be complaining of varying degrees of pain, it is desirable to diminish the degree of suffering that a patient experiences.
 - b. The intent of this protocol is to provide a standing order for the treatment of severe pain when the source of the pain is clear.
 - i. Examples of painful conditions that may be addressed by this protocol include, but are not limited to:
 1. Isolated orthopedic injuries
 2. Burns
 3. Kidney stones
 4. Muscle spasms
2. Contraindications:
 - a. Altered mental status (due to any internal or external cause)
 - b. Hypotension (Adult) – defined as systolic BP < 100 mmHg
 - c. Hypotension (Pediatric) – refer to age appropriate BP chart
 - d. Hypoxia, defined as SpO₂ of <95%
 - e. Head Injury
 - f. Undiagnosed abdominal Pain (contact Medic-1 for orders)
 - g. New onset back or flank pain in patients > 65 years of age
 - i. This is due to a concern for masking a potential acute abdominal aneurysm
 - ii. Contact Medic 1 for kidney stone pain management for patients ≥ 65 years of age.
3. Standards of Care:
 - a. Standard ALS/BLS supportive care
 - b. Reliable oximetry
 - c. The narcotic antidote, **naloxone** (0.1 mg/kg IV, IM, IN or SQ) must be quickly and readily available when using this protocol.
 - d. Parental consent is required, if available, before treating pediatric patients with narcotics
 - e. For medication associated nausea and vomiting, follow the **HCFR NAUSEA/VOMITING** protocol.
 - i. Prophylactic administration of ondansetron is **not** authorized under this or any protocol
4. Treatments:
 - a. Basic ALS Treatments.
 - i. Document the patient's perception of their pain severity on a scale from 0 – 10 before, during and post medication administration
 1. A visual analog scale may be appropriate for pediatric patients and those for whom a 0 – 10 scale is not feasible (i.e. language barrier).
 - ii. Document vital signs, mental status, and drug allergies prior to medication administration. Repeat and document vital signs and mental status again post medication administration.
 - iii. Document provocation, quality, region, severity, and timing (PQRST) for any patient presenting with a primary complaint of pain, not just those for given pain management measures.
 - b. Inhalational analgesia and anxiolytic agent (if available)
 - i. **Nitrous oxide**: (if available) by self-administered patient inhalation
 - ii. This can be used in all age groups

Section: ALS Protocols
Subject: PAIN MANAGEMENT
Section #: 345.17
Issue Date: March 21, 2011
Revision Date: December 1, 2017
Approved By: 

Page 2 of 2

Michael Lozano, Jr., M.D., HCFR Medical Director

- c. Analgesia agents: (use one **or** the other)
 - i. **Fentanyl:** (for normal to high BP; contraindicated if hypotensive)
 - 1. Adults: Initial dose of 100 mcg IV/IM/IN, and then 50 mcg IV/IM/IN as needed q5 minutes to a maximum dose 350 mcg.
 - 2. Pediatrics: Initial dose of 1 mcg/kg IV/IM/IN, and then 0.5 mcg/kg IV/IM/IN as needed q5 minutes to a maximum of 3.5 mcg/kg).
 - ii. **Morphine:** (for normal to high BP; contraindicated if hypotensive)
 - 1. Adults: 2 mg IV/IM/SQ, and then as needed q5 minutes to a maximum of 20 mg.
 - 2. Pediatrics: 0.1 mg/kg IV/IM/SQ (maximum 2 mg), and then as needed q5 minutes to a maximum of 20 mg.
 - d. Adjunctive agents (if needed)
 - i. **Midazolam:** (for anxiety with normal to high BP)
 - 1. Adults: 2.5 mg IV/IM/IN, and then as needed q5 min to a maximum of 10 mg.
 - 2. Pediatrics: 0.025 mg/kg (max dose of 2.5 mg) IV/IM/IN, and then as needed q5 min to a maximum of 10 mg.
 - 3. Geriatrics: 1.25 mg IV/IM/IN, and then as needed q5 min to a maximum of 5 mg.
5. Contact Medic-1:
- a. For additional doses of pain medications above the listed total maximum dosages
 - b. If source of pain is unclear or if there are extenuating or unusual circumstances about the call
 - c. For advice on situations beyond this protocol