

The quorum exponents: what reduces, what the classical bounds miss, what remains

D. Joubert

3 September 2026

Abstract

The every-scale quorum asks for two bounds on the bottom vector v of Q_L : $\delta_p = v^\top T_p v \leq e^{-7p+c}$ and $\kappa_p = \|T_p v - (v^\top T_p v)v\| \geq e^{-4p-c'}$, uniformly in the window. Both quantities are pairings of v against the prime tower T_p . The leading term is the window autocorrelation of v at lag $\log p$. Bandlimited decay (Slepian, Bernstein, integration by parts) gives powers of $1/\log p$, not e^{-7p} . The measured silence is the decay of a near-kernel, not of a desert prolate. The hole shrinks to one estimate: $\|T_p v\|$ itself is exponentially small in p . That estimate is not proved. Nothing here is RH.

1 Reduction

T_p is the contribution of the powers $p^k \leq \mu$ to the prime-side form,

$$T_p(f, f) = \sum_{p^k \leq \mu} (\log p) p^{-k/2} \Theta_f(k \log p), \quad \Theta_f(y) = \int f(x) f(x+y) dx,$$

with Θ_f supported in $[0, L]$, $L = \log \mu$, and $\Theta_f(L) = 0$. Then $\delta_p = T_p(v, v)$ and $\kappa_p^2 = \|T_p v\|^2 - \delta_p^2$. The $k = 1$ term dominates for $p \geq 5$:

$$\delta_p = \frac{\log p}{\sqrt{p}} \Theta_v(\log p) + O(p^{-1} \Theta_v(2 \log p)). \quad (1)$$

So $\delta_p \leq e^{-7p+c}$ is a pointwise bound on the autocorrelation of one explicit vector at the points $\log p$.

2 What the classical bounds give

If v were merely unit-norm in PW_τ ,

$$|\Theta_v(y)| \leq \min(1, C_N(\tau y)^{-N})$$

after N integrations by parts on a smooth cutoff. At $y = \log p$ this is a power of $1/\log p$. The Slepian that lives in the desert $(-\gamma_1, \gamma_1)$ has $\Theta(y) \sim \gamma_1 \operatorname{sinc}(\gamma_1 y/\pi)$, which at $p = 7$ is $O(1/28)$, against a measured $\delta_7 \sim 10^{-16}$. Duffin–Schaeffer on gaps loses the same exponential that *sampling-floor* already recorded for c_L .

None of these tools sees e^{-7p} . The exponent linear in p is not the Fourier decay of a function bandlimited to γ_1 .

3 What the measurements say instead

At $\mu = 11$, $-\ln \delta_p \approx 7p - 12$ for $p \geq 5$, and later $-\ln \delta_p \approx 0.19 s(\chi) p \log p$ across ζ, χ_3, χ_4 . Both δ_p and κ_p are exponentially small, so $\|T_p v\|$ itself is exponentially small: v is close to the kernel of T_p , not just orthogonal to $T_p v$ in one direction. That is stronger than silence. It means the near-kernel of Q is already almost orthogonal to every individual tower of a large prime.

This matches the leakage picture: the mass of $\hat{v}$ is expelled to the band edge, and the pairing against a lag $\log p$ — a frequency modulation of width 1 — sees that edge through a rapidly oscillating integral. Writing a bound that tracks the measured $7p$ requires a pointwise description of $\hat{v}$ on the edge, uniformly in L . The leakage profile $\ln |\hat{v}(\gamma)| \approx -\tau(\omega_{\max} - \gamma)$ is measured, not proved, and τ itself moves with μ .

4 An attempt

Θ_v is supported in $[0, L]$. The points $y = \log p$ lie *inside* the support. Exponential decay as $y \rightarrow \infty$ is the wrong picture: there is no infinity. What is asked is a dip of size e^{-7p} at each arithmetic lag $\log p \in (0, L)$.

Near the edge. Bernstein plus $\Theta_v(L) = 0$ gives $|\Theta_v(\log p)| \leq \tau \log(\mu/p)$. Useful only for $p = \mu^{1-o(1)}$. At $p = 7$, $\mu = 11$, the bound is $O(1)$ against 10^{-16} .

Contour shift. $F \in PW_\tau$ is entire of type τ , so $\int F(\omega) \overline{F(\omega)} e^{i\omega y} d\omega$ can be shifted. The shift produces a factor $e^{-\delta y}$ and a bound $e^{\tau\delta}$ on the new line — net $e^{(-\delta+\tau)y}$ at best, i.e. a power of p , and only after replacing $|F|^2$ on $\mathbb{R}$ by $F(z)\overline{F(\bar{z})}$ with control we do not have on the ground state. Super-exponential in $p = e^y$ never comes from a strip.

Fixed p , $L \rightarrow \infty$. The measured δ_p is stable from $\mu = 8$ to 16. The statement uniform in μ is therefore: the Beurling extremal v_L of the zero set in $PW_{L/2}$ has $\Theta_{v_L}(\log p)$ bounded by e^{-7p} for each fixed p , for all large L . That is a quantitative property of one sampling-extremal function attached to Γ . It is the same order of difficulty as a quantitative lower bound for c_L , already recorded as out of reach of Duffin–Schaeffer / Bernstein (*sampling-floor*, §4).

5 Two lemmas that do hold

Lemma 1 (edge). $\Theta_v(L) = 0$, hence the tower of a prime sitting at the window edge vanishes identically ($T_{11} \equiv 0$ at $\mu = 11$).

Lemma 2 (crude). $|\delta_p| \leq (\log p) p^{-1/2} \|v\|_1^2$. In particular $\delta_p = O(p^{-1/2} \log p)$.

Lemma 2 is weaker than the 2×2 criterion by an exponential factor, exactly as Bernstein was weaker than c_L by $e^{L\gamma/2}$.

Status

Tried. The three routes — decay at infinity, contour shift, edge continuity — miss e^{-7p} for the same structural reason: the lag is interior and the smallness is a dip of the Beurling extremal at arithmetic points. That dip is measured and stable in μ . It is not proved. The hole is the same hole as the quantitative floor. No RH.