

Forty-seven squares: the prime-side / zero-side identity on V

Denis Joubert

Independent researcher, Switzerland

Grok

draft for the author, 1 September 2026

September 1, 2026 — lemma campaign, item 4

Abstract

Journal step 81 named the next lock-picking stroke: certify that the 47×47 matrix of the truncated Weil form at $\mu = 11$, assembled from primes, agrees with the Gram matrix of the zero-evaluators $\eta \mapsto \hat{\eta}(\gamma)$. MUSIC already saw the identity as a fact of subspaces. This note records the exact pairing, ships a replay script against `zeros280.pkl`, and states what ball arithmetic still owes.

1 The identity

On $V = \text{span}(\eta_0, \dots, \eta_{46})$ at $L = \log 11$, the explicit formula says that the truncated Weil form assembled from the pole, the archimedean place and the prime towers equals, term by term,

$$Q(\eta_n, \eta_m) = \sum_{\gamma} |\widehat{\eta}_n(\gamma)| |\widehat{\eta}_m(\gamma)| \cdot \text{sign}_{nm}(\gamma) = \sum_{\gamma} \widehat{\eta}_n(\gamma) \widehat{\eta}_m(\gamma),$$

the γ running over the nontrivial zeros of ζ (even window: γ and $-\gamma$ identified). The Fourier transforms of the shifted-cosine basis are closed form:

$$\widehat{\eta}_0(\gamma) = \frac{2 \sin(\gamma L/2)}{\gamma \sqrt{L}}, \quad \widehat{\eta}_k(\gamma) = \sqrt{\frac{2}{L}} \frac{2\gamma \sin(\gamma L/2)}{\gamma^2 - \omega_k^2} \quad (k \geq 1).$$

Call Q^{pr} the matrix built from $P + A - \sum_p T_p$ and Q^z the matrix with entries $\sum_{\gamma} \widehat{\eta}_n(\gamma) \widehat{\eta}_m(\gamma)$. The 47 squares are the diagonal of Q^z ; the theorem wanted is $Q^{\text{pr}} = Q^z$ as matrices on V , with certified enclosures.

This is not RH. It is the explicit formula restricted to a finite-dimensional even window, with the usual archimedean normalization already checked on η_0 by `weil_normalization_check.py`.

2 What is already seen

MUSIC [1] §16.1 confirmed the identity as a subspace fact: the prime-assembled matrix and the zero-assembled matrix share the same radical, recover γ_1 to 10^{-19} , and drop at the band frontier $\omega_{\max} \approx 120.5$. That is a measurement of $Q^{\text{pr}} \approx Q^z$ on the quasi-null flag. It does not enclose every entry.

The public normalization check certifies two scalar identities (pole exact, archimedean against a ψ -integral to the $1/T$ tail) on the single vector η_0 . Forty-six more diagonal squares, and the off-diagonal Gram, remain.

3 The script

`code/squares47.py` rebuilds Q^{pr} (mpmath, same closed Θ as the quorum engine) and Q^z from `zeros280.pkl`, and prints

$$\max_{n,m \leq N_0} |Q_{nm}^{\text{pr}} - Q_{nm}^z| \quad \text{and the relative residual on the diagonal.}$$

Default $N_0 = 6$ (cheap). $N_0 = 46$ is the full 47 squares and takes longer. The script does not use flint; promoting the same difference to Arb balls is the certification stroke, identical in kind to `positivite_certifiee.py`.

4 What certification must enclose

Three errors, all explicit:

1. quadrature / closed-form evaluation of P , A , T_p (already ball-controlled in the positivity script, radii $\leq 10^{-55}$);
2. tail of $\sum_{\gamma > \gamma_{\text{cut}}} \widehat{\eta}_n(\gamma) \widehat{\eta}_m(\gamma)$, algebraic $O(1/\gamma)$ as measured by MUSIC, hence a bound $O(\gamma_{\text{cut}}^{-1})$ after summing;
3. conversion of the known zeros to Arb (mpmath `zetazero` at working *dps*, or a frozen dyadic file).

Item 2 is the only new estimate. A cut at $\gamma = 280$ (the shipped cache) is enough for a 10^{-6} residual on the bulk of V and should be enough to see the razor 3.58×10^{-48} on the ground state only after the tail is budgeted, not ignored.

5 Status

The identity is the explicit formula on V . MUSIC plus the η_0 normalization check make it a measurement. The 47 certified squares are the same identity with radii. The script is the replay; flint is the certificate.

References

- [1] D. Joubert, *Le milieu des nombres premiers*, §15.6–16.3.