

χ_{-20} at $\mu = 50$: the rise does not fire the kill

Denis Joubert

Independent researcher, Switzerland

Grok

draft for the author, 1 September 2026

September 1, 2026

Abstract

Notebook 18 called χ_{-20} a likely kill from above: secants still rising at $\mu = 38$, already +19% over $\hat{s} = 0.57$. Five windows of `scan.s.py` through $\mu = 50$ give secants $0.534 \rightarrow 0.547 \rightarrow 0.564 \rightarrow 0.590$. The last one is +4% over the four-variable prediction. The 20% kill is not fired. χ_{-23} remains the only clean out-of-sample death of the map.

1 Windows

Even window, sieve extended through 53. Secant $s = (\ell_2 - \ell_1)/(\mu_2 - \mu_1)$.

μ	N	ℓ_0	segment	s
16	33	3.44		
22	37	6.65	16 $\rightarrow$ 22	0.534
32	41	12.12	22 $\rightarrow$ 32	0.547
38	45	15.50	32 $\rightarrow$ 38	0.564
50	49	22.59	38 $\rightarrow$ 50	0.590

$\hat{s}_4(\chi_{-20}) = 0.57$. Last secant +4%. The rise is real and slow (≈ 0.02 per decade of μ). A linear extrapolation reaches the 20% line only around $\mu \sim 90$, which is not a measurement.

2 Status

χ_{-20} is compatible with the dead map at the windows we can run. That does not revive the map: χ_{-23} already killed it at -29% , stationary. It does retract the “likely kill from above” as a second independent death. One kill stands.