

Depth as sampling geometry, and the quorum as a 2×2 mechanism

D. Joubert (with the assistance of two language models)

1 September 2026 — consolidates notebook §66–71 of `riemann-weil-phenomenology`

Abstract

Two results replace two earlier readings. (1) The depth law $-\ln \lambda_{\min} = s(\chi)\mu + b$ of the truncated Weil forms, previously described by a four-variable regression on fourteen Dirichlet characters that failed out of sample, is a *geometric* property of the zero set: the floor c_L of the form is the sampling constant of the zeros for the Paley–Wiener space of type $L/2$, made exponentially small by the desert $(-\gamma_1, \gamma_1)$ and by the gaps longer than the Nyquist step $2\pi/L$. A two-coefficient law, the coefficients fitted on four scales of ζ alone ($\mu = 3, 8, 11, 16$), predicts $s(\chi)$ out of sample with median ratio 0.89 (10 of 14 characters within 20%), with no character-specific parameter. (2) The quorum theorem (every proper sub-Euler product violates positivity) has a mechanism: an unconditional 2×2 interlacing lemma, *depth $\times$ coupling*, whose three quantities we certify at $\mu = 11$ and measure at three scales; the coupling of the bottom vector to a missing prime decays in p at half the rate of its silence, which is what makes the determinant negative. Both are statements about explicitly defined forms; nothing bears on RH.

1 Depth is a sampling constant

Everything in this section is under RH (equivalently, under the identification $Q = Q_z$ of the prime-side form with the zero Gram): it describes the floor once the zeros are on the line. Section 2 uses no zeros at all. Under RH, $Q_L(f) = \sum_{\gamma} \hat{f}(\gamma)^2$ with $\hat{f} \in PW_{L/2}$, so $c_L = \inf Q_L / \|f\|^2$ is a sampling constant of the zero set, positive by Beurling (note *sampling-floor*, Theorem 1). The finite-basis minima have a floor. At $\mu = 3$ the saturation is visible: $\lambda_{\min}(N) = 1.026, 0.775, 0.641, 0.591, 0.571, 0.562 \times 10^{-7}$ for $N = 9 \dots 61$, successive ratios 1.32, 1.21, 1.08, 1.03, 1.02, converging geometrically to $c_{\log 3} \approx 5.55 \times 10^{-8}$. At $\mu = 11$ it is so far a trend: 3.59, 1.86, 1.54×10^{-48} for $N = 47, 57, 67$, ratios 1.93 then 1.21 — still a slope, decelerating toward something near 1.4×10^{-48} , not yet a saturation.

Synthetic controls (zero-side Gram, $\mu = 3$). $Q_z = \sum_{\gamma} 2\hat{c}(\gamma)\hat{c}(\gamma)^{\top}$ reproduces the prime-side floor (ratio 0.93, the truncation tail). Three fictitious zeros placed in the desert raise $\lambda_{\min}$ by 9×10^6 ; a comb with the same γ_1 and density changes it by 0.23 only; shifting the whole spectrum changes $-\ln \lambda_{\min}$ linearly in the desert width with slope 1.45 against $L = 1.10$ for Slepian’s single hole. The desert is the cause; the arithmetic fine structure touches only the prefactor.

The geometric law. With $\nu = 2\pi/L$ the Nyquist step,

$$-\ln c_L \approx a L (\gamma_1 - \nu)_+ + b L \sum_k (\gamma_{k+1} - \gamma_k - \nu)_+, \quad a = 1.69, b = 0.82,$$

the two coefficients fitted on the μ -series of ζ only ($\mu = 3, 8, 11, 16$, within 8%; Slepian’s naive values would be 2, 1). Let $\gamma^*(L)$ be the last height at which a consecutive gap exceeds ν (for

$\mu = 11$: gaps above $\nu = 2.62$ occur up to $\gamma \approx 300$ in the first 500 zeros, sporadically, while the band edge of V_{47} is $\omega_{\max} = 120.5$). As L grows, ν shrinks and the excess sum extends and grows; over $\mu = 3, 8, 11, 16$ it is empirically linear in μ , which is where the linearity of the depth law comes from in this description.

Out of sample (coefficients frozen). Predicted s = slope of the law between $\mu = 11$ and 22, against the measured s of the depth note:

χ ratio	ζ 1.22	χ_3 0.82	χ_4 0.82	χ_5 0.87	χ_7 1.08	χ_8 1.00	χ_{11} 1.17	χ_{12} 0.88
χ ratio	χ_{13} 1.35	χ_{15} 1.02	χ_{19} 0.96	χ_{21} 0.65	χ_{24}^e 1.04	χ_{24}^o 0.38	χ_{-23} 0.89	

Median 0.89; ten of fourteen within 20%; no parameter per character. The downward bias is mechanical (zero caches end near $\gamma \approx 85 < \gamma^*(22) = 138$). Of the two failures, χ_{24}^o ($\gamma_1 = 1.98$) has $\gamma_1 < \nu$ at both scales ($\nu(11) = 2.62$, $\nu(22) = 2.03$): its desert term is identically zero, and the linearized $(\gamma_1 - \nu)_+$ is the culprit — Slepian’s $1 - \lambda_0(c)$ does not switch off. χ_{21} ($\gamma_1 = 2.32$) sits between $\nu(22)$ and $\nu(11)$: the term switches on between the two scales, which explains only part of the 0.65; the rest is an unexplained residue. Where the four-variable map (trained on these characters) died out of sample (χ_{-23} : -29%), the geometric law predicts χ_{-23} to 0.89.

What the desert alone explains. Computing the Slepian constant to arbitrary precision: $1 - \lambda_0(\tau\gamma_1) = 3.35 \times 10^{-6}$, 5.42×10^{-14} , 2.93×10^{-16} at $\mu = 3, 11, 16$, i.e. 75%, 28%, 21% of the measured depth. The union of all sub-Nyquist gaps gives 74 of 110 nats at $\mu = 11$ but is not a bound in either direction. The sharp constant is c_L itself.

Conjecture 1 (candidate form). *The desert term is $-\ln(1 - \lambda_0(\frac{L}{2}\gamma_1))$, Slepian’s constant, in place of a $L(\gamma_1 - \nu)_+$. The gap term is the part not yet understood: the union concentration constant is not a bound in either direction, and the coefficient b is a fitted number. Whether a, b are absolute across L -functions is suggested by the out-of-sample table, not established.*

2 The quorum as a mechanism

Write $Q_S = Q + T_M$, T_M the towers of the missing primes M .

Lemma 2 (unconditional). *Let v be a unit vector, $w = T_M v$, $\kappa = \|w - (v^\top w)v\|$, $u = (w - (v^\top w)v)/\kappa$, and $a = v^\top Qv + v^\top T_M v$, $d = u^\top Q_S u$. If $a d < \kappa^2$ then Q_S is indefinite.*

Proof. The 2×2 principal minor of Q_S on $\{v, u\}$ is $\begin{pmatrix} a & \kappa \\ \kappa & d \end{pmatrix}$ with negative determinant, hence a negative eigenvalue; Cauchy interlacing. $\square$

No eigenvector exactness and no hypothesis on zeros is used. Taking v the bottom vector of Q : $a \approx \varepsilon + \sum_{p \in M} \delta_p$ with $\varepsilon = \lambda_{\min}$ and $\delta_p = v^\top T_p v$ the *silence* of v to p ; κ is its *coupling*. Depth $\times$ coupling: this is the content of the lemmas B1/B2 of the note *lemma-quorum-scales*, in the form in which they are provable.

Certified instance, $\mu = 11$ (code/quorum.2x2.py, Q and T_p in Arb balls). $\varepsilon = 3.5832 \times 10^{-48} \pm 3 \times 10^{-53}$. $T_{11} \equiv 0$ ($\Theta(L) = 0$ at the window edge), so the voting primes are $\{2, 3, 5, 7\}$ — the $2^4 - 1$ of the certified theorem. All fifteen proper sub-products are certified indefinite by 2×2 certificates built on the *same* v ; worst margin $M = \{7\}$: $\kappa^2 = 3.8 \times 10^{-9}$ against $a d = 1.4 \times 10^{-16}$. This re-proves the $\mu = 11$ theorem independently of its 340 witnesses.

The μ -content. High-precision measurement at three scales, criterion checked on every proper subset:

μ	ε	δ_p ($p = 2, 3, 5, 7, \dots$)	κ_p	subsets	re
8	7.5×10^{-33}	0.052, 3.1e-4, 1.3e-10, 9.4e-21	0.70, 0.53, 3.6e-3, 8.8e-9	14/14	
11	3.6×10^{-48}	0.058, 4.8e-4, 1.2e-9, 7.4e-17	0.74, 0.77, 0.051, 6.2e-5	14/14	
16	1.9×10^{-73}	0.064, 6.6e-4, 4.9e-9, 4.0e-15, 2.0e-30, 1.2e-40	0.80, 0.88, 0.28, 6.7e-3, 1.5e-9, 2.5e-15	62/62	

Two laws in p at fixed μ : $-\ln \delta_p \approx 7p - 12$ and $-\ln \kappa_p \approx 4p$ ($p \geq 5$). *The coupling decays at half the rate of the silence*; $\kappa_p^2/\delta_p \geq 9$ everywhere and grows with μ (for $p = 7$: $8 \times 10^3 \rightarrow 5 \times 10^7 \rightarrow 10^{10}$). The worst case always contains 2 ($a \approx \delta_2 \approx 0.06$ against $\kappa^2 \approx 0.5\text{--}0.9$); two preregistered guesses died (the silence grows mildly with μ ; the worst case is not the largest isolated prime).

What remains for every scale. Three unconditional analytic statements about the bottom vector, in the directions the lemma needs (silence small, coupling large): $\varepsilon(\mu) \leq e^{-s\mu+b}$ (the desert, Section 1, there under RH), $\delta_p(\mu) \leq e^{-7p+c}$, $\kappa_p(\mu) \geq e^{-4p-c'}$. What is measured is $\approx$; what is to be proved is $\leq$ and $\geq$. The first is the depth law; the other two are new, concern the autocorrelation of an explicit eigenvector at the points $\log p$, and are where an analyst can enter.

Status

Lemma proved; $\mu = 11$ instance certified; everything else measured with stated numbers. Neither section moves RH: the first describes how small the positive floor on the line is, the second why removing a prime destroys it.