## Strictly Confidential: (For Internal and Restricted use only) Senior Secondary School Term–II Examination, 2022

Marking Scheme: CHEMISTRY (Subject Code: 043)

[ Paper Code: 56/2/3]

## General Instructions: -

- You are aware that evaluation is the most important process in the actual and correct assessment of the candidates. A small mistake in evaluation may lead to serious problems which may affect the future of the candidates, education system and teaching profession. To avoid mistakes, it is requested that before starting evaluation, you must read and understand the spot evaluation guidelines carefully.
- 2. "Evaluation policy is a confidential policy as it is related to the confidentiality of the examinations conducted, Evaluation done and several other aspects. Its' leakage to public in any manner could lead to derailment of the examination system and affect the life and future of millions of candidates. Sharing this policy/document to anyone, publishing in any magazine and printing in News Paper/Website etc may invite action under IPC."
- 3. Evaluation is to be done as per instructions provided in the Marking Scheme. It should not be done according to one's own interpretation or any other consideration. Marking Scheme should be strictly adhered to and religiously followed. However, while evaluating, answers which are based on latest information or knowledge and/or are innovative, they may be assessed for their correctness otherwise and marks be awarded to them. In class-X, while evaluating two competency-based questions, please try to understand given answer and even if reply is not from marking scheme but correct competency is enumerated by the candidate, marks should be awarded.
- 4. The Head-Examiner must go through the first five answer books evaluated by each evaluator on the first day, to ensure that evaluation has been carried out as per the instructions given in the Marking Scheme. The remaining answer books meant for evaluation shall be given only after ensuring that there is no significant variation in the marking of individual evaluators.
- 5. Evaluators will mark( $\sqrt{\ }$ ) wherever answer is correct. For wrong answer 'X" be marked. Evaluators will not put right kind of mark while evaluating which gives an impression that answer is correct and no marks are awarded. **This is most common mistake which evaluators are committing.**
- 6. If a question has parts, please award marks on the right-hand side for each part. Marks awarded for different parts of the question should then be totalled up and written in the left-hand margin and encircled. This may be followed strictly.
- 7. If a question does not have any parts, marks must be awarded in the left-hand margin and encircled. This may also be followed strictly.
- 8. If a student has attempted an extra question, answer of the question deserving more marks should be retained and the other answer scored out.
- 9. No marks to be deducted for the cumulative effect of an error. It should be penalized only once.
- 10. A full scale of marks 0-35 has to be used. Please do not hesitate to award full marks if the answer deserves it.
- 11. Every examiner has to necessarily do evaluation work for full working hours i.e., 8 hours every day and evaluate 30 answer books per day in main subjects and 35 answer books per day in other subjects (Details are given in Spot Guidelines). This is in view of the reduced syllabus and number of questions in question paper.
- 12. Ensure that you do not make the following common types of errors committed by the Examiner in the past:-
  - Leaving answer or part thereof unassessed in an answer book.

- Giving more marks for an answer than assigned to it.
- Wrong totaling of marks awarded on a reply.
- Wrong transfer of marks from the inside pages of the answer book to the title page.
- Wrong question wise totaling on the title page.
- Wrong totaling of marks of the two columns on the title page.
- Wrong grand total.
- Marks in words and figures not tallying.
- Wrong transfer of marks from the answer book to online award list.
- Answers marked as correct, but marks not awarded. (Ensure that the right tick mark is correctly and clearly indicated. It should merely be a line. Same is with the X for incorrect answer.)
- Half or a part of answer marked correct and the rest as wrong, but no marks awarded.
- 13. While evaluating the answer books if the answer is found to be totally incorrect, it should be marked as cross (X) and awarded zero (0) Marks.
- 14. Any unassessed portion, non-carrying over of marks to the title page, or totalling error detected by the candidate shall damage the prestige of all the personnel engaged in the evaluation work as also of the Board. Hence, in order to uphold the prestige of all concerned, it is again reiterated that the instructions be followed meticulously and judiciously.
- 15. The Examiners should acquaint themselves with the guidelines given in the Guidelines for spot Evaluation before starting the actual evaluation.
- 16. Every Examiner shall also ensure that all the answers are evaluated, marks carried over to the title page, correctly totalled and written in figures and words.
- 17. The Board permits candidates to obtain photocopy of the Answer Book on request in an RTI application and also separately as a part of the re-evaluation process on payment of the processing charges.

## MARKING SCHEME

Senior Secondary School Examination TERM-II, 2022

## CHEMISTRY (Subject Code-043)

[ Paper Code: 56/2/3]

Q. No.	EXPECTED ANSWER / VALUE POINTS	Marks			
	SECTION—A				
1.	CH <sub>3</sub> CHOH				
	(i) CH <sub>3</sub> / Propan-2-ol				
	(ii) CH <sub>3</sub> — CH <sub>3</sub> / Ethane				
	(iii) CH <sub>3</sub> COO <sup>-</sup> / Acetate ion (Any two)	1×2			
2.	$\kappa = \frac{1}{R} \left( \frac{l}{A} \right)$	1/2			
	$1.5 \times 10^{-4} \text{ S cm}^{-1} = \frac{1}{1200 \Omega} \left( \frac{l}{A} \right)$				
	$\left(\frac{l}{A}\right) = 1.5 \times 10^{-4} \times 1200 \text{ cm}^{-1}$				
	$\left(\frac{l}{A}\right) = 0.18 \text{ cm}^{-1}$	1/2			
	$\wedge_m = \frac{\kappa}{C} \times 1000 \ \Omega^{-1} \ cm^2 \ mol^{-1}$	1/2			
	$= \frac{1 \cdot 5 \times 10^{-4}}{0 \cdot 001} \times 1000 \ \Omega^{-1} \ cm^2 \ mol^{-1}$				
	$\wedge_m = 150 \ \Omega^{-1} \ \text{cm}^2 \ \text{mol}^{-1} \ \text{or} \ 150 \ \text{S} \ \text{cm}^2 \ \text{mol}^{-1}$	1/2			
3.	(i) Rate increases 9 times	1			
	(ii) Rate increases 8 times	1			
	SECTION—B				
4.	(i) CH <sub>3</sub> COOH / Ethanoic acid				
	(ii) CH <sub>3</sub> CHO / Ethanal				
	СООН				
	(iii) Benzoic acid				
		1×3			

4.		OR			
	(b) (i) CH <sub>3</sub> COCH <sub>2</sub> CH <sub>3</sub> < CH <sub>3</sub> COCH <sub>3</sub> < CH <sub>3</sub> CH <sub>2</sub> CHO < CH <sub>3</sub> CHO				
	(ii) (CH <sub>3</sub> ) <sub>2</sub> CH-COOH < CH <sub>3</sub> COOH < Cl-CH <sub>2</sub> -COOH				
	(iii) соон	соон			
	OCH <sub>3</sub>				
		NO <sub>2</sub>			
5.	(a) (i) CHO	1			
		No₂ / m-Nitrobenzaldehyde			
	(ii)	/ Cyclohexane			
		-			
	$(iii)$ $CH_3$ — $C=$	=N—OH / Ethanal oxime	12		
	Н		1×3		
5.		OR			
		tion of propanal involves cleavage of C—H bond than C—C bond of propanone.	1		
		NaOH and $I_2$ , acetophenone gives yellow precipitate of			
		s benzophenone does not.	1		
	(iii)	$NO_{2}$			
	$CH_3$ — $C$ = $NNH$ — $NO_2$				
	CH <sub>3</sub>				
6.	$E_{\text{cell}} = (E_C^{\circ} - E_A^{\circ}) - \frac{0.059}{2} \log \frac{[Zn^{2+}]}{[Ag^+]^2}$				
	1-01				
	$= [0.80 - (-0.76)] - \frac{0.059}{2} \log \frac{(0.01)}{(0.001)^2}$				
	$=1.56-\frac{0.059}{2}\log 10^4$				
	= (1.56 - 0.118) V				
	$E_{\text{cell}} = 1 \cdot 442 \text{ V}$	(Deduct ½ marks if no or incorrect unit)	1		
7.	(a)				
	Lyophobic Sol	Lyophilic Sol			
	Solvent repelling	Solvent loving			
	Irreversible	Reversible			
	Unstable	Stable			
	Easy to coagulate	Difficult to coagulate (Any three)	1 x 3		

(b) (i) Dispersed phase = liquid and dispersion medium = liquid  (ii) Unbalanced bombardment of the colloidal particles by the molecules of dispersion medium.  (iii) Because it is exothermic in nature / weak bonds between adsorbate and adsorbent break.  8.	
dispersion medium.  (iii) Because it is exothermic in nature / weak bonds between adsorbate and adsorbent break.  8. $\mathbf{k} = \frac{2 \cdot 303}{t} \log \frac{(R)_0}{(R)}$ $\mathbf{k} = \frac{2 \cdot 303}{30} \log \frac{100}{25}$ $\mathbf{k} = \frac{2 \cdot 303}{30} \log 4 = 0.046 \text{ min}^{-1}$ $t_{1/2} = \frac{0 \cdot 693}{k}$ $= \frac{0 \cdot 693}{0.046}$ $t_{1/2} = 15 \text{ min}$ (Deduct ½ marks if no or incorrect unit) (or any other correct method)  9. (i) Variable or multiple oxidation state / ability to form complexes / they provide large surface area for adsorption (utilises (n-1) d and ns electrons for bonding).  (ii) Zn, Cd and Hg have completely filled <i>d</i> -orbitals in its ground state as well as	
(iii) Because it is exothermic in nature / weak bonds between adsorbate and adsorbent break.  8. $k = \frac{2 \cdot 303}{t} \log \frac{(R)_0}{(R)}$ $k = \frac{2 \cdot 303}{30} \log \frac{100}{25}$ $k = \frac{2 \cdot 303}{30} \log 4 = 0.046 \text{ min}^{-1}$ $t_{1/2} = \frac{0 \cdot 693}{k}$ $= \frac{0 \cdot 693}{0.046}$ $t_{1/2} = 15 \text{ min}$ (Deduct ½ marks if no or incorrect unit) (or any other correct method)  9. (i) Variable or multiple oxidation state / ability to form complexes / they provide large surface area for adsorption (utilises (n-1) d and ns electrons for bonding). (ii) Zn, Cd and Hg have completely filled d-orbitals in its ground state as well as	
adsorbent break.  8. $k = \frac{2 \cdot 303}{t} \log \frac{(R)_0}{(R)}$ $k = \frac{2 \cdot 303}{30} \log \frac{100}{25}$ $k = \frac{2 \cdot 303}{30} \log 4 = 0.046 \text{ min}^{-1}$ $t_{1/2} = \frac{0 \cdot 693}{k}$ $= \frac{0 \cdot 693}{0.046}$ $t_{1/2} = 15 \text{ min}$ (Deduct ½ marks if no or incorrect unit) (or any other correct method)  9. (i) Variable or multiple oxidation state / ability to form complexes / they provide large surface area for adsorption (utilises (n-1) d and ns electrons for bonding). (ii) Zn, Cd and Hg have completely filled <i>d</i> -orbitals in its ground state as well as	
8. $k = \frac{1 \log \frac{100}{(R)}}{t \log \frac{100}{25}}$ $k = \frac{2 \cdot 303}{30} \log \frac{100}{25}$ $k = \frac{2 \cdot 303}{30} \log 4 = 0.046 \text{ min}^{-1}$ $t_{1/2} = \frac{0 \cdot 693}{k}$ $= \frac{0 \cdot 693}{0 \cdot 046}$ $t_{1/2} = 15 \text{ min}$ (Or any other correct unit) (or any other correct method)  9. (i) Variable or multiple oxidation state / ability to form complexes / they provide large surface area for adsorption (utilises (n-1) d and ns electrons for bonding). (ii) Zn, Cd and Hg have completely filled d-orbitals in its ground state as well as	
$\mathbf{k} = \frac{2 \cdot 303}{30} \log \frac{100}{25}$ $\mathbf{k} = \frac{2 \cdot 303}{30} \log 4 = 0.046 \text{ min}^{-1}$ $t_{1/2} = \frac{0 \cdot 693}{\mathbf{k}}$ $= \frac{0 \cdot 693}{0 \cdot 046}$ $t_{1/2} = 15 \text{ min}$ (Or any other correct unit) (or any other correct method)  9. (i) Variable or multiple oxidation state / ability to form complexes / they provide large surface area for adsorption (utilises (n-1) d and ns electrons for bonding). (ii) Zn, Cd and Hg have completely filled <i>d</i> -orbitals in its ground state as well as	2
$\mathbf{k} = \frac{2 \cdot 303}{30} \log 4 = 0.046 \text{ min}^{-1}$ $t_{1/2} = \frac{0.693}{\mathbf{k}}$ $= \frac{0.693}{0.046}$ $t_{1/2} = 15 \text{ min}$ (or any other correct unit) (or any other correct method)  9. (i) Variable or multiple oxidation state / ability to form complexes / they provide large surface area for adsorption (utilises (n-1) d and ns electrons for bonding). (ii) Zn, Cd and Hg have completely filled d-orbitals in its ground state as well as	
$\mathbf{k} = \frac{2 \cdot 303}{30} \log 4 = 0.046 \text{ min}^{-1}$ $t_{1/2} = \frac{0.693}{\mathbf{k}}$ $= \frac{0.693}{0.046}$ $t_{1/2} = 15 \text{ min}$ (or any other correct unit) (or any other correct method)  9. (i) Variable or multiple oxidation state / ability to form complexes / they provide large surface area for adsorption (utilises (n-1) d and ns electrons for bonding). (ii) Zn, Cd and Hg have completely filled d-orbitals in its ground state as well as	2
$t_{1/2} = \frac{0.693}{k}$ $= \frac{0.693}{0.046}$ $t_{1/2} = 15 \text{ min}$ (Deduct ½ marks if no or incorrect unit) (or any other correct method)  9. (i) Variable or multiple oxidation state / ability to form complexes / they provide large surface area for adsorption (utilises (n-1) d and ns electrons for bonding). (ii) Zn, Cd and Hg have completely filled <i>d</i> -orbitals in its ground state as well as	
$t_{1/2} = \frac{0.693}{k}$ $= \frac{0.693}{0.046}$ $t_{1/2} = 15 \text{ min}$ (Deduct ½ marks if no or incorrect unit) (or any other correct method)  9. (i) Variable or multiple oxidation state / ability to form complexes / they provide large surface area for adsorption (utilises (n-1) d and ns electrons for bonding). (ii) Zn, Cd and Hg have completely filled <i>d</i> -orbitals in its ground state as well as	,
$= \frac{0.693}{0.046}$ $t_{1/2} = 15 \text{ min} \qquad \text{(Deduct } \frac{1}{2} \text{ marks if no or incorrect unit)} \qquad \text{(or any other correct method)}$ $9.  (i) \text{ Variable or multiple oxidation state / ability to form complexes / they provide large surface area for adsorption (utilises (n-1) d and ns electrons for bonding).}$ $(ii) \text{ Zn, Cd and Hg have completely filled } d\text{-orbitals in its ground state as well as}$	
$= \frac{0.693}{0.046}$ $t_{1/2} = 15 \text{ min} \qquad \text{(Deduct } \frac{1}{2} \text{ marks if no or incorrect unit)} \qquad \text{(or any other correct method)}$ $9.  (i) \text{ Variable or multiple oxidation state / ability to form complexes / they provide large surface area for adsorption (utilises (n-1) d and ns electrons for bonding).}$ $(ii) \text{ Zn, Cd and Hg have completely filled } d\text{-orbitals in its ground state as well as}$	
$t_{1/2} = 15 \text{ min}$ (Deduct ½ marks if no or incorrect unit) (or any other correct method)  9. (i) Variable or multiple oxidation state / ability to form complexes / they provide large surface area for adsorption (utilises (n-1) d and ns electrons for bonding). (ii) Zn, Cd and Hg have completely filled d-orbitals in its ground state as well as	2
$t_{1/2} = 15 \text{ min}$ (Deduct ½ marks if no or incorrect unit) (or any other correct method)  9. (i) Variable or multiple oxidation state / ability to form complexes / they provide large surface area for adsorption (utilises (n-1) d and ns electrons for bonding). (ii) Zn, Cd and Hg have completely filled d-orbitals in its ground state as well as	
(or any other correct method)  9. (i) Variable or multiple oxidation state / ability to form complexes / they provide large surface area for adsorption (utilises (n-1) d and ns electrons for bonding).  (ii) Zn, Cd and Hg have completely filled d-orbitals in its ground state as well as	
9. (i) Variable or multiple oxidation state / ability to form complexes / they provide large surface area for adsorption (utilises (n-1) d and ns electrons for bonding). (ii) Zn, Cd and Hg have completely filled d-orbitals in its ground state as well as	
large surface area for adsorption (utilises (n-1) d and ns electrons for bonding). (ii) Zn, Cd and Hg have completely filled d-orbitals in its ground state as well as	
(ii) Zn, Cd and Hg have completely filled d-orbitals in its ground state as well as	
in its oxidised state.	
(iii) Because of lanthanoid contraction /poor shielding effect of 4f orbitals. $1\times$	:3
<b>10.</b> (a) (i) $sp^3$ , paramagnetic $\frac{1}{2}$	1/2
(ii) $d^2 s p^3$ , diamagnetic	
(iii) $sp^3d^2$ , paramagnetic	1/2
10. OR	
(b) (i) Tetraammineaquachloridocobalt (III) chloride	
(ii) Ambidentate ligand is a ligand which has two different donor atoms	
and either of the two ligates with metal atom or ion in the complex whereas	
when a ligand ligates through two donor atoms is called a bidentate ligand.	
(iii) $[Fe(C_2O_4]_3]^{3-}$ , because of chelate effect / due to cyclic structure	1/2
11. (i) Because of the stable half-filled $3d^5$ configuration in $Mn^{2+}$ .	
(ii) Because of the irregular sum of $(\Delta_i H_1 + \Delta_i H_2)$ values and much less	
submination enthalpies for win and v.	
(iii) Because of high $\Delta_a H^0$ and low $\Delta_{hyd} H^0$	
SECTION—C	
12. (i) Aniline is a Lewis base and it reacts with AlCl <sub>3</sub> to form a salt / N of aniline	
acquires positive charge with AlCl <sub>3</sub> and hence is a deactivating group.	
$(ii)$ $(CH_3)_3N < C_2H_5NH_2 < NH_3 < C_6H_5NH_2$	
$C_2H_5 NH_2 < (CH_3)_3N < NH_3 < C_6H_5NH_2$	
(As no medium (aqueous /gaseous) or phase is given both answers to be	
considered).	

(iii) Add Hinsberg reagent (benzene sulphonyl chloride) to both the compounds. CH<sub>3</sub>CH<sub>2</sub>NH<sub>2</sub> gives ppt. that is soluble in alkali while the ppt. formed by 1 (CH<sub>3</sub>CH<sub>2</sub>)<sub>2</sub> NH is insoluble in alkali. (*iv*) (a) (I) (B)  $\frac{1}{2} + \frac{1}{2}$ (II) (A)  $CH_3CH_2NH_2$ (B) CH<sub>3</sub>CH<sub>2</sub>NHCOCH<sub>3</sub> 1/2+1/2 OR *(b)* COOH CONH<sub>2</sub> NaOBr 1 (I)NHCOCH<sub>3</sub> NHCOCH<sub>3</sub>  $NH_2$  $NH_2$ CH<sub>3</sub>COCl/Py 1 (II)FeBr<sub>3</sub> Br Br (or any other suitable method of conversion)

\* \* \*