

Exercise 5d

Question 1.

If $A = \begin{bmatrix} 2 & -3 & 5 \\ 0 & 7 & -4 \end{bmatrix}$, verify that $(A')' = A$.

Answer:

Transpose of a matrix is obtained by interchanging the rows and the columns of matrix A. It is denoted by A' .

e.g. $A_{12} = A_{21}$

$$A = \begin{bmatrix} 2 & -3 & 5 \\ 0 & 7 & -4 \end{bmatrix}$$

Hence transpose of matrix A is,

$$A' = \begin{bmatrix} 2 & 0 \\ -3 & 7 \\ 5 & -4 \end{bmatrix}$$

$$(A')' = \begin{bmatrix} 2 & -3 & 5 \\ 0 & 7 & -4 \end{bmatrix}$$

$$(A')' = A$$

Hence, Proved.

Question 2.

If $A = \begin{bmatrix} 3 & 5 \\ -2 & 0 \\ 4 & -6 \end{bmatrix}$, verify that $(2A)' = 2A'$.

Answer:

$$\text{Given } A = \begin{bmatrix} 3 & 5 \\ -2 & 0 \\ 4 & -6 \end{bmatrix}$$

To Prove: $(2A)' = 2A'$

Proof: Let us consider, $B = 2A$

$$\text{Now, } B = 2 \begin{bmatrix} 3 & 5 \\ -2 & 0 \\ 4 & -6 \end{bmatrix}$$

$$\Rightarrow \begin{bmatrix} 6 & 10 \\ -4 & 0 \\ 8 & -12 \end{bmatrix}$$

$$\text{LHS} \Rightarrow B' = \begin{bmatrix} 6 & -4 & 8 \\ 10 & 0 & -12 \end{bmatrix}$$

Again to find RHS, we will find the transpose of matrix A

$$A' = \begin{bmatrix} 3 & -2 & 4 \\ 5 & 0 & -6 \end{bmatrix}$$

$$\text{RHS} = 2A'$$

$$\Rightarrow 2 \begin{bmatrix} 3 & -2 & 4 \\ 5 & 0 & -6 \end{bmatrix}$$

$$\Rightarrow \begin{bmatrix} 6 & -4 & 8 \\ 10 & 0 & -12 \end{bmatrix}$$

$$\text{LHS} = \text{RHS}$$

Hence proved.

Question 3.

$$\text{If } A = \begin{bmatrix} 3 & 2 & -1 \\ -5 & 0 & -6 \end{bmatrix} \text{ and } B = \begin{bmatrix} -4 & -5 & -2 \\ 3 & 1 & 8 \end{bmatrix}, \text{ verify that } (A + B)' = (A' + B').$$

Answer:

$$\text{Given } A = \begin{bmatrix} 3 & 2 & -1 \\ -5 & 0 & -6 \end{bmatrix} \text{ and } B = \begin{bmatrix} -4 & -5 & -2 \\ 3 & 1 & 8 \end{bmatrix}$$

To Prove: $(A + B)' = A' + B'$

Proof: Let us consider $C = A + B$

$$C = \begin{bmatrix} 3 & 2 & -1 \\ -5 & 0 & -6 \end{bmatrix} + \begin{bmatrix} -4 & -5 & -2 \\ 3 & 1 & 8 \end{bmatrix}$$

$$\Rightarrow \begin{bmatrix} -1 & -3 & -3 \\ -2 & 1 & 2 \end{bmatrix}$$

Now LHS = C'

$$\Rightarrow \begin{bmatrix} -1 & -2 \\ -3 & 1 \\ -3 & 2 \end{bmatrix}$$

To find RHS, we will find transpose of matrix A and B

$$A' = \begin{bmatrix} 3 & -5 \\ 2 & 0 \\ -1 & -6 \end{bmatrix} \text{ And } B' = \begin{bmatrix} -4 & 3 \\ -5 & 1 \\ -2 & 8 \end{bmatrix}$$

RHS = $A' + B'$

$$\Rightarrow \begin{bmatrix} 3 & -5 \\ 2 & 0 \\ -1 & -6 \end{bmatrix} + \begin{bmatrix} -4 & 3 \\ -5 & 1 \\ -2 & 8 \end{bmatrix}$$

$$\Rightarrow \begin{bmatrix} -1 & -2 \\ -3 & 1 \\ -3 & 2 \end{bmatrix}$$

$$\text{LHS} = \text{RHS}$$

Hence proved.

Question 4.

If $P = \begin{bmatrix} 3 & 4 \\ 2 & -1 \\ 0 & 5 \end{bmatrix}$ and $Q = \begin{bmatrix} 7 & -5 \\ -4 & 0 \\ 2 & 6 \end{bmatrix}$, verify that $(P + Q)' = (P' + Q')$.

Answer:

Given $P = \begin{bmatrix} 3 & 4 \\ 2 & -1 \\ 0 & 5 \end{bmatrix}$ and $Q = \begin{bmatrix} 7 & -5 \\ -4 & 0 \\ 2 & 6 \end{bmatrix}$

To Prove: $(P + Q)' = P' + Q'$

Proof: Let us consider $R = P + Q$,

$$R = \begin{bmatrix} 3 & 4 \\ 2 & -1 \\ 0 & 5 \end{bmatrix} + \begin{bmatrix} 7 & -5 \\ -4 & 0 \\ 2 & 6 \end{bmatrix}$$

$$\Rightarrow \begin{bmatrix} 10 & -1 \\ -2 & -1 \\ 2 & 11 \end{bmatrix}$$

$$\text{LHS} = R \Rightarrow (P + Q)'$$

$$\text{LHS} = \begin{bmatrix} 10 & -2 & 2 \\ -1 & -1 & 11 \end{bmatrix}$$

To find RHS, we will first find the transpose of matrix P and Q

$$P' = \begin{bmatrix} 3 & 2 & 0 \\ 4 & -1 & 5 \end{bmatrix} \text{ And } Q' = \begin{bmatrix} 7 & -4 & 2 \\ -5 & 0 & 6 \end{bmatrix}$$

$$\text{RHS} = P' + Q'$$

$$\Rightarrow \begin{bmatrix} 3 & 2 & 0 \\ 4 & -1 & 5 \end{bmatrix} + \begin{bmatrix} 7 & -4 & 2 \\ -5 & 0 & 6 \end{bmatrix}$$

$$\Rightarrow \begin{bmatrix} 10 & -2 & 2 \\ -1 & -1 & 11 \end{bmatrix}$$

$$\text{LHS} = \text{RHS}$$

Hence proved.

Question 5.

If $A = \begin{bmatrix} 4 & 1 \\ 5 & 8 \end{bmatrix}$, show that $(A + A')$ is symmetric.

Answer:

$$\text{Given } A = \begin{bmatrix} 4 & 1 \\ 5 & 8 \end{bmatrix}$$

To Prove: $A + A'$ is symmetric. (Note: A matrix P is symmetric if $P' = P$)

Proof: We will find A' ,

$$A' = \begin{bmatrix} 4 & 5 \\ 1 & 8 \end{bmatrix}$$

Now let us take $P = A + A'$

$$P = \begin{bmatrix} 4 & 1 \\ 5 & 8 \end{bmatrix} + \begin{bmatrix} 4 & 5 \\ 1 & 8 \end{bmatrix}$$

$$\Rightarrow \begin{bmatrix} 8 & 6 \\ 6 & 16 \end{bmatrix}$$

$$\text{Now } P' = \begin{bmatrix} 8 & 6 \\ 6 & 16 \end{bmatrix}$$

$$\Rightarrow P' = P$$

Hence $A + A'$ is a symmetric matrix.

Question 6.

If $A = \begin{bmatrix} 3 & -4 \\ 1 & -1 \end{bmatrix}$, show that $(A + A')$ is skew-symmetric.

Answer:

$$\text{Given } A = \begin{bmatrix} 3 & -4 \\ 1 & -1 \end{bmatrix}$$

To prove: $A - A'$ is a skew-symmetric matrix. (Note: A matrix P is skew-symmetric if $P' = -P$)

Proof: First we will find the transpose of matrix A

$$A' = \begin{bmatrix} 3 & 1 \\ -4 & -1 \end{bmatrix}$$

Let us take $P = A - A'$

$$P = \begin{bmatrix} 3 & -4 \\ 1 & -1 \end{bmatrix} - \begin{bmatrix} 3 & 1 \\ -4 & -1 \end{bmatrix}$$

$$\Rightarrow \begin{bmatrix} 0 & -5 \\ 5 & 0 \end{bmatrix}$$

$$P = \begin{bmatrix} 0 & 5 \\ -5 & 0 \end{bmatrix}$$

$$\Rightarrow P' = P$$

Hence $A-A'$ is a skew symmetric matrix.

Question 7.

Show that the matrix $A = \begin{bmatrix} 0 & a & b \\ -a & 0 & c \\ -b & -c & 0 \end{bmatrix}$ is skew-symmetric.

HINT: Show that $A' = -A$.

Answer:

$$\text{Given } A = \begin{bmatrix} 0 & a & b \\ -a & 0 & c \\ -b & -c & 0 \end{bmatrix}$$

To Prove: A is a skew symmetric matrix.

Proof: As for a matrix to be skew symmetric $A' = -A$

We will find A' .

$$A' = \begin{bmatrix} 0 & -a & -b \\ a & 0 & -c \\ b & c & 0 \end{bmatrix}$$

$$= - \begin{bmatrix} 0 & a & b \\ -a & 0 & c \\ -b & -c & 0 \end{bmatrix}$$

$$\Rightarrow A' = -A$$

So A is A skew symmetric matrix.

Question 8.

Express the matrix $A = \begin{bmatrix} 2 & 3 \\ -1 & 4 \end{bmatrix}$ as the sum of a symmetric matrix and a skew-symmetric matrix.

Answer:

Given $A = \begin{bmatrix} 2 & 3 \\ -1 & 4 \end{bmatrix}$, As for a symmetric matrix $A' = A$ hence

$$A + A' = 2A$$

$$A = \frac{1}{2}(A + A') \Rightarrow P \text{ (Symmetric Matrix)}$$

Similarly for a skew symmetric matrix since $A' = -A$ hence

$$A - A' = 2A$$

$$A = \frac{1}{2}(A - A') \Rightarrow Q \text{ (Skew Symmetric Matrix)}$$

So a matrix can be represented as a sum of a symmetric matrix P and skew symmetric matrix Q.

First, we will find the transpose of matrix A,

$$A' = \begin{bmatrix} 2 & -1 \\ 3 & 4 \end{bmatrix}$$

Now using the above formulas,

$$P = \frac{1}{2}(A + A')$$

$$\Rightarrow \frac{1}{2} \left(\begin{bmatrix} 2 & 3 \\ -1 & 4 \end{bmatrix} + \begin{bmatrix} 2 & -1 \\ 3 & 4 \end{bmatrix} \right)$$

$$\Rightarrow \frac{1}{2} \begin{bmatrix} 4 & 2 \\ 2 & 8 \end{bmatrix}$$

$$P = \begin{bmatrix} 2 & 1 \\ 1 & 4 \end{bmatrix}$$

$$Q = \frac{1}{2} \left(\begin{bmatrix} 2 & 3 \\ -1 & 4 \end{bmatrix} - \begin{bmatrix} 2 & -1 \\ 3 & 4 \end{bmatrix} \right)$$

$$\Rightarrow \frac{1}{2} \begin{bmatrix} 0 & 4 \\ -4 & 0 \end{bmatrix}$$

$$\Rightarrow \begin{bmatrix} 0 & 2 \\ -2 & 0 \end{bmatrix}$$

Hence $A = P + Q$

$$\Rightarrow \begin{bmatrix} 2 & 1 \\ 1 & 4 \end{bmatrix} + \begin{bmatrix} 0 & 2 \\ -2 & 0 \end{bmatrix} \text{ [Matrix A as the sum of P and Q]}$$

$$\Rightarrow \begin{bmatrix} 2 & 3 \\ -1 & 4 \end{bmatrix}$$

Question 9.

Express the matrix $A = \begin{bmatrix} 3 & -4 \\ 1 & -1 \end{bmatrix}$ as the sum of a symmetric matrix and a skew-symmetric matrix.

Answer:

Given $A = \begin{bmatrix} 3 & -4 \\ 1 & -1 \end{bmatrix}$, to express as the sum of symmetric matrix P and skew symmetric matrix Q.

$$A = P + Q$$

Where $P = \frac{1}{2}(A + A')$ and $Q = \frac{1}{2}(A - A')$, we will find transpose of matrix A

$$A' = \begin{bmatrix} 3 & 1 \\ -4 & -1 \end{bmatrix}$$

Now using the above formulas

$$P = \frac{1}{2}(A + A')$$

$$\Rightarrow \frac{1}{2} \left(\begin{bmatrix} 3 & -4 \\ 1 & -1 \end{bmatrix} + \begin{bmatrix} 3 & 1 \\ -4 & -1 \end{bmatrix} \right)$$

$$\Rightarrow \frac{1}{2} \begin{bmatrix} 6 & -3 \\ -3 & -2 \end{bmatrix}$$

$$\Rightarrow \begin{bmatrix} 3 & \frac{-3}{2} \\ \frac{-3}{2} & -1 \end{bmatrix}$$

$$Q = \frac{1}{2}(A - A')$$

$$\Rightarrow \frac{1}{2} \left(\begin{bmatrix} 3 & -4 \\ 1 & -1 \end{bmatrix} - \begin{bmatrix} 3 & 1 \\ -4 & -1 \end{bmatrix} \right)$$

$$\Rightarrow \frac{1}{2} \begin{bmatrix} 0 & -5 \\ 5 & 0 \end{bmatrix}$$

$$\Rightarrow \begin{bmatrix} 0 & \frac{-5}{2} \\ \frac{5}{2} & 0 \end{bmatrix}$$

Hence $A = P + Q$

$$\Rightarrow \begin{bmatrix} 3 & \frac{-3}{2} \\ \frac{-3}{2} & -1 \end{bmatrix} + \begin{bmatrix} 0 & \frac{-5}{2} \\ \frac{5}{2} & 0 \end{bmatrix} \quad [\text{Matrix A as the sum of P and Q}]$$

$$\Rightarrow \begin{bmatrix} 3 & \frac{-8}{2} \\ \frac{2}{2} & -1 \end{bmatrix}$$

$$\Rightarrow \begin{bmatrix} 3 & -4 \\ 1 & -1 \end{bmatrix}$$

Question 10.

Express the matrix $A = \begin{bmatrix} -1 & 5 & 1 \\ 2 & 3 & 4 \\ 7 & 0 & 9 \end{bmatrix}$ as the sum of a symmetric and a skew-symmetric matrix.

Answer:

Given $A = \begin{bmatrix} -1 & 5 & 1 \\ 2 & 3 & 4 \\ 7 & 0 & 9 \end{bmatrix}$, to express as sum of symmetric matrix P and skew symmetric matrix

Q.

$$A = P + Q$$

Where $P = \frac{1}{2}(A + A')$ and $Q = \frac{1}{2}(A - A')$,

First, we find A'

$$A' = \begin{bmatrix} -1 & 2 & 7 \\ 5 & 3 & 0 \\ 1 & 4 & 9 \end{bmatrix}$$

Now using the above mentioned formulas

$$P = \frac{1}{2}(A + A')$$

$$\Rightarrow \frac{1}{2} \left(\begin{bmatrix} -1 & 5 & 1 \\ 2 & 3 & 4 \\ 7 & 0 & 9 \end{bmatrix} + \begin{bmatrix} -1 & 2 & 7 \\ 5 & 3 & 0 \\ 1 & 4 & 9 \end{bmatrix} \right)$$

$$\Rightarrow \frac{1}{2} \begin{bmatrix} -2 & 7 & 8 \\ 7 & 6 & 4 \\ 8 & 4 & 18 \end{bmatrix}$$

$$\Rightarrow \begin{bmatrix} -1 & \frac{7}{2} & 4 \\ \frac{7}{2} & 3 & 2 \\ 4 & 2 & 9 \end{bmatrix}$$

$$Q = \frac{1}{2}(A - A')$$

$$\Rightarrow \frac{1}{2} \left(\begin{bmatrix} -1 & 5 & 1 \\ 2 & 3 & 4 \\ 7 & 0 & 9 \end{bmatrix} - \begin{bmatrix} -1 & 2 & 7 \\ 5 & 3 & 0 \\ 1 & 4 & 9 \end{bmatrix} \right)$$

$$\Rightarrow \frac{1}{2} \begin{bmatrix} 0 & 3 & -6 \\ -3 & 0 & 4 \\ 6 & -4 & 0 \end{bmatrix}$$

$$\Rightarrow \begin{bmatrix} 0 & \frac{3}{2} & -3 \\ -\frac{3}{2} & 0 & 2 \\ 3 & -2 & 0 \end{bmatrix}$$

Now $A = P + Q$

$$\Rightarrow \begin{bmatrix} -1 & \frac{7}{2} & 4 \\ \frac{7}{2} & 3 & 2 \\ 4 & 2 & 9 \end{bmatrix} + \begin{bmatrix} 0 & \frac{3}{2} & -3 \\ -\frac{3}{2} & 0 & 2 \\ 3 & -2 & 0 \end{bmatrix} \quad [\text{Matrix A as sum of P and Q}]$$

$$\Rightarrow \begin{bmatrix} -1 & \frac{10}{2} & 1 \\ \frac{4}{2} & 3 & 4 \\ 7 & 0 & 9 \end{bmatrix}$$

$$A = \begin{bmatrix} -1 & 5 & 1 \\ 2 & 3 & 4 \\ 7 & 0 & 9 \end{bmatrix}$$

Question 11.

Express the matrix A as the sum of a symmetric and a skew-symmetric matrix, where

$$A = \begin{bmatrix} 3 & -1 & 0 \\ 2 & 0 & 3 \\ 1 & -1 & 2 \end{bmatrix}.$$

Answer:

Given $A = \begin{bmatrix} 3 & -1 & 0 \\ 2 & 0 & 3 \\ 1 & -1 & 2 \end{bmatrix}$, to express as sum of symmetric matrix P and skew symmetric matrix

Q

$$A = P + Q$$

Where $P = \frac{1}{2}(A + A')$ and $Q = \frac{1}{2}(A - A')$,

First we will find A' ,

$$A' = \begin{bmatrix} 3 & 2 & 1 \\ -1 & 0 & -1 \\ 0 & 3 & 2 \end{bmatrix}$$

Now using above mentioned formulas,

$$P = \frac{1}{2}(A + A')$$

$$\Rightarrow \frac{1}{2} \left(\begin{bmatrix} 3 & -1 & 0 \\ 2 & 0 & 3 \\ 1 & -1 & 2 \end{bmatrix} + \begin{bmatrix} 3 & 2 & 1 \\ -1 & 0 & -1 \\ 0 & 3 & 2 \end{bmatrix} \right)$$

$$\Rightarrow \frac{1}{2} \begin{bmatrix} 6 & 1 & 1 \\ 1 & 0 & 2 \\ 1 & 2 & 4 \end{bmatrix}$$

$$\Rightarrow \begin{bmatrix} 3 & \frac{1}{2} & \frac{1}{2} \\ \frac{1}{2} & 0 & 1 \\ \frac{1}{2} & 1 & 2 \end{bmatrix}$$

$$Q = \frac{1}{2}(A - A')$$

$$\Rightarrow \frac{1}{2} \left(\begin{bmatrix} 3 & -1 & 0 \\ 2 & 0 & 3 \\ 1 & -1 & 2 \end{bmatrix} - \begin{bmatrix} 3 & 2 & 1 \\ -1 & 0 & -1 \\ 0 & 3 & 2 \end{bmatrix} \right)$$

$$\Rightarrow \frac{1}{2} \begin{bmatrix} 0 & -3 & -1 \\ 3 & 0 & 4 \\ 1 & -4 & 0 \end{bmatrix}$$

$$\Rightarrow \begin{bmatrix} 0 & \frac{-3}{2} & \frac{-1}{2} \\ \frac{3}{2} & 0 & 2 \\ \frac{1}{2} & -2 & 0 \end{bmatrix}$$

Now $A = P + Q$

$$\Rightarrow \begin{bmatrix} 3 & \frac{1}{2} & \frac{1}{2} \\ \frac{1}{2} & 0 & 1 \\ \frac{1}{2} & 1 & 2 \end{bmatrix} + \begin{bmatrix} 0 & \frac{-3}{2} & \frac{-1}{2} \\ \frac{3}{2} & 0 & 2 \\ \frac{1}{2} & -2 & 0 \end{bmatrix} \quad [\text{Matrix A as sum of P and Q}]$$

$$\Rightarrow \begin{bmatrix} 3 & \frac{-2}{2} & 0 \\ \frac{4}{2} & 0 & 3 \\ 1 & -1 & 2 \end{bmatrix} = \begin{bmatrix} 3 & -1 & 0 \\ 2 & 0 & 3 \\ 1 & -1 & 2 \end{bmatrix}$$

Question 12.

Express the matrix $A = \begin{bmatrix} 3 & 2 & 5 \\ 4 & 1 & 3 \\ 0 & 6 & 7 \end{bmatrix}$ as sum of two matrices such that one is symmetric and the other is skew-symmetric.

Answer:

Given $A = \begin{bmatrix} 3 & 2 & 5 \\ 4 & 1 & 3 \\ 0 & 6 & 7 \end{bmatrix}$, to express as sum of symmetric matrix P and skew symmetric matrix

Q.

$$A = P + Q$$

$$\text{Where } P = \frac{1}{2}(A + A') \text{ and } Q = \frac{1}{2}(A - A'),$$

First we will find A'

$$A' = \begin{bmatrix} 3 & 4 & 0 \\ 2 & 1 & 6 \\ 5 & 3 & 7 \end{bmatrix}$$

Now using above mentioned formulas

$$P = \frac{1}{2}(A + A')$$

$$\Rightarrow \frac{1}{2} \left(\begin{bmatrix} 3 & 2 & 5 \\ 4 & 1 & 3 \\ 0 & 6 & 7 \end{bmatrix} + \begin{bmatrix} 3 & 4 & 0 \\ 2 & 1 & 6 \\ 5 & 3 & 7 \end{bmatrix} \right)$$

$$\Rightarrow \frac{1}{2} \begin{bmatrix} 6 & 6 & 5 \\ 6 & 2 & 9 \\ 5 & 9 & 14 \end{bmatrix}$$

$$\Rightarrow \begin{bmatrix} 3 & 3 & \frac{5}{2} \\ 3 & 1 & \frac{9}{2} \\ \frac{5}{2} & \frac{9}{2} & 7 \end{bmatrix}$$

$$Q = \frac{1}{2}(A - A')$$

$$\Rightarrow \frac{1}{2} \left(\begin{bmatrix} 3 & 2 & 5 \\ 4 & 1 & 3 \\ 0 & 6 & 7 \end{bmatrix} - \begin{bmatrix} 3 & 4 & 0 \\ 2 & 1 & 6 \\ 5 & 3 & 7 \end{bmatrix} \right)$$

$$\Rightarrow \frac{1}{2} \begin{bmatrix} 0 & -2 & 5 \\ 2 & 0 & -3 \\ -5 & 3 & 0 \end{bmatrix}$$

$$\Rightarrow \begin{bmatrix} 0 & -1 & \frac{5}{2} \\ 1 & 0 & \frac{-3}{2} \\ \frac{-5}{2} & \frac{3}{2} & 0 \end{bmatrix}$$

Now $A = P + Q$

$$\Rightarrow \begin{bmatrix} 3 & 3 & \frac{5}{2} \\ 3 & 1 & \frac{9}{2} \\ \frac{5}{2} & \frac{9}{2} & 7 \end{bmatrix} + \begin{bmatrix} 0 & -1 & \frac{5}{2} \\ 1 & 0 & \frac{-3}{2} \\ \frac{-5}{2} & \frac{3}{2} & 0 \end{bmatrix}$$

$$\Rightarrow \begin{bmatrix} 3 & 2 & 5 \\ 4 & 1 & 3 \\ 0 & 6 & 7 \end{bmatrix}$$

Question 13.

For each of the following pairs of matrices A and B, verify that $(AB)' = (B' A')$:

$$A = \begin{bmatrix} 1 & 3 \\ 2 & 4 \end{bmatrix} \text{ and } B = \begin{bmatrix} 1 & 4 \\ 2 & 5 \end{bmatrix}$$

Answer:

Let us take $C = AB$

$$C = \begin{bmatrix} 1 & 3 \\ 2 & 4 \end{bmatrix} \begin{bmatrix} 1 & 4 \\ 2 & 5 \end{bmatrix}$$

$$\Rightarrow \begin{bmatrix} 1+6 & 4+15 \\ 2+8 & 8+20 \end{bmatrix}$$

$$C = \begin{bmatrix} 7 & 19 \\ 10 & 28 \end{bmatrix}$$

$$\text{LHS} \Rightarrow C' = \begin{bmatrix} 7 & 10 \\ 19 & 28 \end{bmatrix}$$

To find RHS we will find transpose of matrix A and B,

$$A' = \begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix} \text{ And } B' = \begin{bmatrix} 1 & 2 \\ 4 & 5 \end{bmatrix}$$

$$\text{RHS} = B'A'$$

$$\Rightarrow \begin{bmatrix} 1 & 2 \\ 4 & 5 \end{bmatrix} \begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix}$$

$$\Rightarrow \begin{bmatrix} 1+6 & 2+8 \\ 3+16 & 8+20 \end{bmatrix}$$

$$\Rightarrow \begin{bmatrix} 7 & 10 \\ 19 & 28 \end{bmatrix}$$

$$\text{LHS} = \text{RHS}$$

Hence proved.

Question 14.

For each of the following pairs of matrices A and B, verify that $(AB)' = (B' A')$:

$$A = \begin{bmatrix} 3 & -1 \\ 2 & -2 \end{bmatrix} \text{ and } B = \begin{bmatrix} 1 & -3 \\ 2 & -1 \end{bmatrix}$$

Answer:

Let us take $C = AB$

$$C = \begin{bmatrix} 3 & -1 \\ 2 & -2 \end{bmatrix} \begin{bmatrix} 1 & -3 \\ 2 & -1 \end{bmatrix}$$

$$\Rightarrow \begin{bmatrix} 3+(-2) & -9+1 \\ 2+(-4) & -6+2 \end{bmatrix}$$

$$\Rightarrow \begin{bmatrix} 1 & -8 \\ -2 & -4 \end{bmatrix}$$

$$\text{LHS} \Rightarrow C' = \begin{bmatrix} 1 & -2 \\ -8 & -4 \end{bmatrix}$$

To find RHS we will find transpose of matrix A and B,

$$B' = \begin{bmatrix} 1 & 2 \\ -3 & -1 \end{bmatrix} \text{ And } A' = \begin{bmatrix} 3 & 2 \\ -1 & -2 \end{bmatrix}$$

$$\text{RHS} = B'A'$$

$$\Rightarrow \begin{bmatrix} 1 & 2 \\ -3 & -1 \end{bmatrix} \begin{bmatrix} 3 & 2 \\ -1 & -2 \end{bmatrix}$$

$$\Rightarrow \begin{bmatrix} 3 + (-2) & 2 + (-4) \\ -9 + 1 & -6 + 2 \end{bmatrix}$$

$$\Rightarrow \begin{bmatrix} 1 & -2 \\ -8 & -4 \end{bmatrix}$$

$$\text{LHS} = \text{RHS}$$

Hence proved.

Question 15.

For each of the following pairs of matrices A and B, verify that $(AB)' = (B' A')$:

$$A = \begin{bmatrix} -1 \\ 2 \\ 3 \end{bmatrix} \text{ and } B = [-2 \ -1 \ -4]$$

Answer:

Let us take $C = AB$

$$C = \begin{bmatrix} -1 \\ 2 \\ 3 \end{bmatrix} \begin{bmatrix} -2 & -1 & -4 \end{bmatrix}$$

$$\Rightarrow \begin{bmatrix} 2 & 1 & 4 \\ -4 & -2 & -8 \\ -6 & -3 & -12 \end{bmatrix}$$

LHS = C'

$$\Rightarrow \begin{bmatrix} 2 & -4 & -6 \\ 1 & -2 & -3 \\ 4 & -8 & -12 \end{bmatrix}$$

To find RHS we will find transpose of matrix A and B,

$$A' = \begin{bmatrix} -1 & 2 & 3 \end{bmatrix} \text{ And } B' = \begin{bmatrix} -2 \\ -1 \\ -4 \end{bmatrix}$$

RHS = $B'A'$

$$\Rightarrow \begin{bmatrix} -2 \\ -1 \\ -4 \end{bmatrix} \begin{bmatrix} -1 & 2 & 3 \end{bmatrix}$$

$$\Rightarrow \begin{bmatrix} 2 & -4 & -6 \\ 1 & -2 & -3 \\ 4 & -8 & -12 \end{bmatrix}$$

LHS = RHS

Hence proved.

Question 16.

For each of the following pairs of matrices A and B, verify that $(AB)' = (B' A')$:

$$A = \begin{bmatrix} -1 & 2 & -3 \\ 4 & -5 & 6 \end{bmatrix} \text{ and } B = \begin{bmatrix} 3 & -4 \\ 2 & 1 \\ -1 & 0 \end{bmatrix}$$

Answer:

Let us take $C = AB$

$$C = \begin{bmatrix} -1 & 2 & -3 \\ 4 & -5 & 6 \end{bmatrix} \begin{bmatrix} 3 & -4 \\ 2 & 1 \\ -1 & 0 \end{bmatrix}$$

$$\Rightarrow \begin{bmatrix} -3 + 4 + 3 & 4 + 2 + 0 \\ 12 + (-10) + (-6) & -16 + (-5) \end{bmatrix}$$

$$\Rightarrow \begin{bmatrix} 4 & 6 \\ -4 & -21 \end{bmatrix}$$

LHS = C'

$$\Rightarrow \begin{bmatrix} 4 & -4 \\ 6 & -21 \end{bmatrix}$$

To find RHS we will find transpose of matrix A and B,

$$A' = \begin{bmatrix} -1 & 4 \\ 2 & -5 \\ -3 & 6 \end{bmatrix} \text{ And } B' = \begin{bmatrix} 3 & 2 & -1 \\ -4 & 1 & 0 \end{bmatrix}$$

RHS = $B'A'$

$$\Rightarrow \begin{bmatrix} 3 & 2 & -1 \\ -4 & 1 & 0 \end{bmatrix} \begin{bmatrix} -1 & 4 \\ 2 & -5 \\ -3 & 6 \end{bmatrix}$$

$$\Rightarrow \begin{bmatrix} -3 + 4 + 3 & 12 + (-10) + (-6) \\ 4 + 2 & -16 + (-5) \end{bmatrix}$$

$$\Rightarrow \begin{bmatrix} 4 & -4 \\ 6 & -21 \end{bmatrix}$$

LHS = RHS

Hence proved.

Question 17.

If $A = \begin{bmatrix} \cos \alpha & \sin \alpha \\ -\sin \alpha & \cos \alpha \end{bmatrix}$, show that $A'A = I$.

Answer:

Given $A = \begin{bmatrix} \cos \alpha & \sin \alpha \\ -\sin \alpha & \cos \alpha \end{bmatrix}$, We will find A'

$$A' = \begin{bmatrix} \cos \alpha & -\sin \alpha \\ \sin \alpha & \cos \alpha \end{bmatrix}$$

LHS = $A'A$

$$\Rightarrow \begin{bmatrix} \cos \alpha & -\sin \alpha \\ \sin \alpha & \cos \alpha \end{bmatrix} \begin{bmatrix} \cos \alpha & \sin \alpha \\ -\sin \alpha & \cos \alpha \end{bmatrix}$$

$$\Rightarrow \begin{bmatrix} \cos^2 \alpha + \sin^2 \alpha & \cos \alpha \sin \alpha + (-\sin \alpha \cos \alpha) \\ \sin \alpha \cos \alpha + (-\cos \alpha \sin \alpha) & \sin^2 \alpha + \cos^2 \alpha \end{bmatrix}$$

$$\Rightarrow \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \text{ [Using } \cos^2 \alpha + \sin^2 \alpha = 1 \text{ and commutative law } a.b = b.a \text{ i.e. } \sin \alpha . \cos \alpha = \cos \alpha . \sin \alpha \text{)]}$$

$$\text{RHS} = I \Rightarrow \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$$

$$\text{LHS} = \text{RHS}$$

Hence proved.

Question 18.

If matrix $A = [1 \ 2 \ 3]$, write AA' .

Answer:

Given $A = [1 \ 2 \ 3]$

We will find A' to calculate AA' ,

$$A' = \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix}$$

Now

$$AA' = [1 \ 2 \ 3] \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix}$$

$$\Rightarrow [1 + 4 + 9]$$

$$\Rightarrow [14]$$