BISCHOFF
CHROMATOGRAPHY

ProntoSIL

HPLC Columns

L Notifications	Packing	Phase names
L 1	Octadecyl silane chemically bonded to porous or non-porous silica or ceramic micro-particles, 3 to 10 µm in diameter, or a monolithic rod	ProntoSIL C18 ACE-EPS ProntoSIL C18 AQ ProntoSIL C18 H ProntoSIL C18 SH Pronto I ODS-3 ProntoSIL Spheribond ODS1 ProntoSIL Spheribond ODS 2 ProntoSIL Hypersorb C18 ProntoSIL C18 BDS ProntoSIL KromaPlus C18 ProntoSIL C18 Eurobond ProntoSIL C18 CB ProntoSIL C18 AQ PLUS HAPAK ODS AB ProntoSIL Prontobond ProntoSIL C18 Basic ProntoSIL CB18 MA ProntoSIL CB18 MH
	Porous silica particles, 3 to 10 µm in diameter, or a monolithic silica rod.	ProntoSIL Silica KromaPlus Silica ProntoSIL SpheriBond SI ProntoSIL HyperSorb SI ProntoSIL CB Silica
L7	Octylsilane chemically bonded to totally or superficially porous silica particles, 3 to 10 µm in diameter, or a monolithic silica rod.	ProntoSIL C8 ACE-EPS ProntoSIL C8 SH ProntoSIL KromaPlus C8 ProntoSIL C8 BDS HIPAK C8 AB ProntoSIL CB C8 ProntoSIL Hypersorb C8 ProntoSIL SpheriBond C8
L 8	An essentially monomolecular layer of aminopropylsilane chemically bonded to totally porous silica gel support, 3 to 10 µm in diameter, or a monolithic silica rod.	ProntoSIL AMINO ProntoSIL AMINO EC ProntoSIL AMINO H ProntoSIL CB Amino ProntoSIL KromaPlus AMINO ProntoSIL Hypersorb AMINO ProntoSIL SpheriBond AMINO
L 9	Irregular or spherical, totally porous silica gel having a chemically bonded, strongly acidic cation-exchange coating, 3 to 10 µm in	ProntoSIL SpheriBond SCX
L 10	Nitrile groups chemically bonded to porous silica particles, 3 to 10 µm in diameter, or a monolithic silica rod.	ProntoSIL CN ProntoSIL CB Cyano ProntoSIL SpheriBond NITRILE ProntoSIL Hypersorb CPS (CN)
L 11	Phenyl groups chemically bonded to porous silica particles, 3 to 10 µm in diameter, or a monolithic silica rod.	ProntoSIL Phenyl ProntoSIL Phenyl Hexyl ProntoSIL CB Phenyl
L14	Silica gel having a chemicallly bonded, strongly basic quaternary ammonium anion-exchange coating, 5 to 10 µm in diameter.	ProntoSIL SpheriBond SAX
L16	Dimethylsilane chemically bonded to porous silica particles, 5 to 10 µm in diameter	ProntoSIL C1 KromaPlus C1 ProntoSIL Spheribond C1 ProntoSIL Hypersorb C1 (SAS)
L17	Strong cation-exchange resin consisting of sulfonated cross-linked styrenedivinylbenzene copolymer in the hydrogen form, 6 to 12 µm in diameter	ProntoGel H ProntoGel Su H

L Notifications	Packing	Phase names
L19	Strong cation-exchange resin consisting of sulfonated cross-linked styrenedivinylbenzene copolymer in the calcium form, about 9 µm in diameter	ProntoGel Ca ProntoGel Su Ca
L20	Dihydroxypropane groups chemically bonded to porous silica or hybrid particles, 3 to 10 µm in diameter, or a monolithic silica rod.	ProntoSIL Diol ProntoSIL CB Diol
L21	A rigid, spherical styrenedivinylbenzene copolymer, 3 to 30 µm in diameter	ProntoGel-P
L22	A cation-exchange resin made of porous polystyrene gel with sulfonic acid groups, about 10 µm in size	ProntoGel Peptide 100A^{0}
L25	Packing having the capacity to separate compounds with amolecular weight range from 100-5000 (as determined by polyethylene oxide), applied to neutral, anionic, and cationic water-soluble polymers. A polymethacrylate resin base, cross-linked with polyhydroxylated ether (surface contained some residual carboxyl functional groups) was found suitable, 3 to 10 µm in diameter, or a monolithic silica rod.	ProntoGel-AQUA-OH
L26	Butyl silane chemically bonded to totally porous silica particles, 3 to 10 µm in diameter	ProntoSIL C4 ProntoSIL KromaPlus C4 ProntoSIL CB Butyl
L34	Strong cation-exchange resin consisting of sulfonated cross-linked styrenedivinylbenzene copolymer in the lead form, 7 to 9 µm in diameter	ProntoGel-PB
L37	Packing having the capacity to separate proteins by molecular size over a range of 2,000 to 40,000 Da. It is a polymethacrylate gel	ProntoGel-AQUA-OH
L38	A methacrylate-based size-exclusion packing for water-soluble samples	ProntoGel-AQUA-OH
L39	A hydrophilic polyhydroxymethacrylate gel of totally porous spherical resin	ProntoGel-AQUA-OH
L40	Cellulose tris-3, 5-dimethylphenylcarbamate coated porous silica particles, 5 µm to 20 µm in diameter	ProntoSIL Chiral POD
L43	Pentafluorophenyl groups chemically bonded to silica particles by a propyl spacer, 2.6 to 10 µm in diameter	ProntoSIL PFP
L51	Amylose tris-3,5-dimethylphenylcarbamate-coated, porous, spherical, silica particles, 5 to 10 µm in diameter	ProntoSIL Chiral PAD
L58	Strong cation-exchange resin consisting of sulfonated cross-linked styrenedivinylbenzene copolymer in the sodium form, about 6 to 30 µm diameter	ProntoGel Su Na
L80	Cellulose tris(4-methylbenzoate)-coated, porous, spherical, silica particles, 5 µ in diameter	ProntoSIL Chiral POJ
L90	Amylose tris(S)-a-methylbenzylcarbamate	ProntoSIL Chiral PAS
L99	Amylose tris(3,5-dimethylphenylcarbamate)	ProntoSIL Chiral PIA
L119	Cellulose tris(3,5-dichlorophenylcarbamate)	ProntoSIL Chiral PIC

Bischoff chromatography offers achiral phases for supercritical fluid chromatography (SFC) applications to provide the chemist with a number of options for SFC separations. These columns are available in many chemistries. Columns include stationary phases based on silica that are coated or covalently bonded.

These columns are specifically packed in SFC compatible hardware and tested individually to guarantee performance. The low viscosity of supercritical carbon dioxide allows for separations, conversation of organic solvents and more concentrated product fractions make SFC a desirable preparative chromatographic technique for purifying chemical mixtures.

ProntoSIL SFC Semi-Prep columns are available with inner diameters from 10mm to 50mm and in lengths from 50mm to 250mm. Individual SFC documentation is included with every column.

Use ProntoSIL SFC columns for following benefits :

1. Complex samples can be separated with better resolution at a faster rate with shorter run times.

2. Sharp Peak shapes on co-ordination compound.

3. Extremely fast equilibration.

ProntoSIL SFC Bondings :

- ProntoSIL SFC CYANO
- ProntoSIL SFC DIOL
- ProntoSIL SFC Ethylpyridine
- ProntoSIL SFC SILICA
- ProntoSIL SFC PHENYL
- ProntoSIL SFC C8 ace EPS
- ProntoSIL SFC C18 SH
- ProntoSIL SFC C18 ace EPS
- ProntoSIL SFC AMINO
- ProntoSIL SFC C30



BISCHOFF
CHROMATOGRAPHY

ProntoSIL

Germany:
BISCHOFF Chromatography GmbH
Böblinger Str. 23, 71229 Leonberg
Phone: +49-7152-6064-0
Web: www.bischoff365.com

India:
B&W Separation India Pvt. Ltd.
(Unit of Bischoff chromatography Germany)
H. No. 262/A1/B1, Daulatnagar, Paud Road,
Bhugaon, Pune - 412115,
Office No. +91 78877 29825
bwteamindia@gmail.com, info@bwseparation.com
www.bwseparation.com

PRONTOGEL2363_EN