# Host-parasite DEB model equations

# Matthew Malishev<sup>1\*</sup>

 $^{1}\ Department\ of\ Biology,\ Emory\ University,\ 1510\ Clifton\ Road\ NE,\ Atlanta,\ GA,\ USA,\ 30322$ 

## Contents

Overview	 2
Reserve mobilisation	 2
Weak homeostasis	 2

Date: 2018-08-17 R version: 3.5.0

\*Corresponding author: matthew.malishev@gmail.com

### Overview

Summarised notes from DEB workshops, telecourses, lectures, and discussions.

#### Reserve mobilisation

Conductance determines mobilisation rate from reserve to structure

The larger the SA of reserve, the more mobilisation is possible and thus faster maintenance and growth because due to more SA - SA scales slower than volume-specific flows

Reserve dynamics f = 1 (max feeding rate)

$$\frac{dE}{dt} = \frac{f\{\dot{p}_{AM}\}}{L} - \frac{\dot{v}[E]}{L}$$

 $[E_M] = \max$  reserve. Reserve doesn't change.

$$= \frac{\{\dot{p}_{AM}\}}{L} - \frac{\dot{v}[E_M]}{L}$$
$$\therefore [E_M] = \frac{\{\dot{p}_{AM}\}}{\dot{v}}$$

Scaled reserve

$$e = \frac{[E]}{[E_M]}$$

$$\frac{de}{dt} = \frac{[E]/[E_M]}{dt} = \frac{f\dot{v}}{L} - \frac{e\dot{v}}{L}$$

$$=\frac{\dot{v}(fe)}{L}$$

Under steady state, reserve doesn't change

$$0 = \frac{\dot{v}(fe)}{L} \quad \text{or} \quad f = e$$

### Length

Getting maximum length  $L_M$ 

$$\frac{dV}{dt} = V\dot{r}$$

Can rewrite r using scaled reserve e

$$\dot{r} = f$$

### Weak homeostasis

Structural isomorphy implies weak homeostasis Weak homeostasis depends on ratio of reserve to structure (d[E]/dt)