

Why the cut bends: the dynamics of the de Sitter cut, the confined gravitational wave, the generative boundary of the slicing operator, and chirality as the turning of the polarization plane

Daryl Janzen

Department of Physics and Engineering Physics
University of Saskatchewan

Abstract

In Cosmological Relativity the exact spherically symmetric and stationary geometries of general relativity are generated as symmetry-breaking cuts of a single de Sitter substrate, with vacuum the straight cut and matter the bend [6], and the range of that construction is the symmetry-reducible sector, bounded by a *wall* at which continuous symmetry is lost and free gravitational radiation begins [7]. This paper takes the next step: the *dynamics*—why and how the cut bends in time. The gap it closes was named at the outset of the programme, in the essay that first developed this cosmology and recorded that the details of dynamical matter had not been worked out.

We work the first inhomogeneous, time-dependent bend explicitly. In a polarized Gowdy–de Sitter model the spatial leaf carries a single transverse-traceless shear—the propagating graviton—advanced by a true Hamiltonian on the substrate’s cosmic foliation, with the cosmological constant driving the area and the wave carrying its own energy and momentum in the leaf’s shear.

The two polarizations are then shown to be a wave map into the hyperbolic plane, the polarized case being a geodesic of the target and the turning of the polarization plane being motion off it. In those variables the cosmological constant is absent from the wave sector entirely: the graviton propagates freely and Λ drives only the area. The unpolarized member carries a conserved twist whose sign is the wave’s handedness, and the substrate’s orientation parity acts on it as a sign flip—so on the reachable sector the graviton’s handedness is the sign of a conserved charge. Because that parity is the *disconnected* component of the target’s isometry group, no connected isometry identifies the two handednesses.

This places the first chiral geometry inside the construction’s own reach rather than past it. The unpolarized turning wave carries the same two spacelike Killing vectors as the polarized one, and the transverse reflection is an isometry of the Gowdy form precisely when the wave is polarized. So the chirality criterion bites from the loss of the swept rotation onward: the polarized edge is achiral, the unpolarized turning wave is the first chiral case, and the wall is where chirality is *generic* rather than where it begins.

At the wall we settle two questions. The dynamics continues across it; and the wall is *not* a metric singularity in the precise sense the causality paper defines—no measure collapses there—so it is a regular radiative boundary rather than a place at which a clock could be re-founded. In the nonlinear $\Lambda > 0$ regime the de Sitter background is an attractor and the first-class system is consistent to all orders, so the continuous dynamics *admits* rather than forces a quantum structure. We close with what is not claimed: no closed-form nonlinear solution, and no non-perturbative quantization of the interacting tower.

1 Introduction: from the static operator to the bend

The slicing operator of Cosmological Relativity generates the static, spherically symmetric sector of general relativity from one de Sitter substrate [6]. *The gap this paper closes was named at the outset of the programme:* the essay that first developed this cosmology recorded that “*details*

involving the emergence of dynamical matter in the SdS cosmology have not been worked out [1]. *This paper works them out* for the inhomogeneous, time-dependent case. A slicing curve on the substrate is the radial profile of a spatial leaf; the straight cut is vacuum (Schwarzschild–de Sitter, the kernel), and a bend in the curve is matter, with density $\rho = m'/4\pi r^2$ fixed by the bend. The companion range paper extends this from a classifier to a map of reach [7]: a geometry is generated as a cut exactly when its isometry group contains a sweep-subgroup of the substrate's, so the range is the *symmetry-reducible sector*, laid out along a null-degeneracy axis from the Nariai seam to a *wall* at which continuous symmetry is lost altogether. At the wall, inhomogeneity, free gravitational radiation, and the failure of any global sweep orientation are one and the same boundary; “why the cut bends” is named there as the open dynamical question.

The symmetric sector already has its answer in closed form, and it is worth stating before the inhomogeneous case departs from it. On any comoving worldline of a symmetric cut, at any energy,

$$\frac{d^2 r}{d\tilde{\tau}^2} = -\frac{f'}{2} = r K_G, \quad (1)$$

with K_G the slicing surface's Gaussian curvature $[3]^R$. *So the rate at which the symmetric cut's bend changes in time is the bend itself*, up to the areal factor: the cut straightens, is momentarily flat, and bends the other way as the geometry passes its one sign change. *That is the whole of the symmetric sector's dynamics*, and what follows is the first case in which the bend is not symmetric.

This paper answers it, in the regime where the answer can be made concrete and exact. The static operator says *what* a bend is—matter, the anchor's image. The dynamical question is *how* a bend evolves: a leaf that is not maximally symmetric must carry its departure from the round S^3 forward in time, and the governing law of that advance is what we exhibit. Three ingredients, each already established in the corpus, combine:

1. the *anchor*—the map from a cut to its stress-energy—whose energy sector is the Hamiltonian constraint and whose momentum sector is the bend in the shift [6, 7];
2. the *cosmic clock*—the substrate's preferred foliation, empirically forced [13] and structurally distinguished, on which the canonical constraint deparametrizes to a true Hamiltonian [9, 8];
3. the *stratification*—the symmetry classes of the range read as the isotropy strata of the construction, from the maximally symmetric vacuum down to the wall [7].

We assemble them on the simplest model that carries a genuine propagating degree of freedom—a linearly polarized Gowdy–de Sitter spacetime [17, 18]—and then read the result back into the stratification and out to the wall. The register is that of the rest of the programme [16, 8]: the field equations are Einstein's, unchanged; the existing object is the evolving three-dimensional layer, and “the cut bends” is a statement about that layer's advance, not about a four-dimensional block. In the layered reading the cosmology runs on, this is the dynamics of the leaf and its rate: the bend *is* the leaf's content (matter, its departure from the round S^3), and the advance that carries it forward is generated by the same lapse whose deparametrized true Hamiltonian is the foliation's rate of advance [6, 9]—so where the static operator gives the leaf and its bend, this paper supplies how that leaf evolves. The wall at the far end of the stratification is the boundary of that layered generation: up to it the leaf is generated by a symmetric sweep of the substrate, and past it, where the last continuous symmetry is gone, it is carried instead by ordinary evolution—the same seam-to-wall span along which the observable cosmology is laid out, the seam its inner (Nariai) end [7].

2 The confined bend: a polarized Gowdy–de Sitter wave

Consider the linearly polarized Gowdy–de Sitter metric

$$ds^2 = e^{2(\gamma-\psi)}(-dt^2 + dz^2) + e^{2\psi} dx^2 + R^2 e^{-2\psi} dy^2, \quad (2)$$

with ψ, γ, R functions of (t, z) only. The field ψ is the single transverse-traceless polarization^R—the leaf’s shear, the propagating graviton; R is the area element of the (x, y) two-torus, the local clock/area variable; γ is the conformal factor on the (t, z) plane. With cosmological constant Λ , the Einstein equations $G_{ab} + \Lambda g_{ab} = 0$ reduce to a wave equation, an area equation, two constraints, and a quadrature for γ :

$$\text{(WAVE)} \quad (R\psi_t)_t - (R\psi_z)_z = \Lambda R e^{2(\gamma-\psi)}, \quad (3)$$

$$\text{(AREA)} \quad R_{tt} - R_{zz} = 2\Lambda R e^{2(\gamma-\psi)}, \quad (4)$$

$$\text{(ENERGY)} \quad 2(R_t\gamma_t + R_z\gamma_z) - (R_{tt} + R_{zz}) = 2R(\psi_t^2 + \psi_z^2), \quad (5)$$

$$\text{(MOMENTUM)} \quad R_t\gamma_z + R_z\gamma_t - R_{tz} = 2R\psi_t\psi_z. \quad (6)$$

These are verified directly from (2) (clean residuals). Equations (5)–(6) are not evolution equations: they are precisely the ADM Hamiltonian and momentum constraints [26] for the foliation by constant- t leaves, and they say that the wave’s energy and momentum are carried *entirely by the shear of the spatial leaf*— $\psi_t^2 + \psi_z^2$ and $\psi_t\psi_z$. This is the anchor at work in the dynamical setting: matter and radiation are the bend, and here the bend is the leaf’s transverse-traceless shear.

A true Hamiltonian, not a frozen constraint. The canonical Hamiltonian density built from (2) generates (3), (4), and the γ -quadrature through Hamilton’s equations: the time development of the leaf is reproduced exactly. The lapse $N = e^{\gamma-\psi}$ is not a free multiplier to be gauged away but the metrical rate at which the existing layer advances, $d\tau = N dt$, exactly as in the cosmological and canonical-time papers [8, 9]. The constraint, read on the substrate’s foliation rather than an arbitrary one, deparametrizes:

$$i \partial_\tau \Psi = \hat{H}_{\text{phys}} \Psi, \quad (7)$$

with H_{phys} the reduced Hamiltonian for the propagating mode. This is the same content as the minisuperspace deparametrization of [9], now carrying a local propagating degree of freedom: the same absolute foliation that makes flat Λ CDM a true-Hamiltonian flow makes the graviton’s confined polarization a true-Hamiltonian flow.

The two regimes. At $\Lambda = 0$ the model degenerates to the classical Gowdy/Einstein–Rosen system [17, 18]: the area R may be used as an internal clock and the wave equation (3) becomes the cylindrical (Bessel) wave equation, $\psi_{TT} + R^{-1}\psi_T - \psi_{zz} = 0$ in the R -time gauge. At $\Lambda > 0$ this internal-clock construction fails— R no longer separates the dynamics cleanly—and the substrate’s cosmic time is the natural clock instead. This is the dynamical analogue of the point made in the canonical-time paper: the absolute foliation is not a convenience but the structure on which the constraint becomes a true Hamiltonian, and it earns that necessity precisely where an internal clock breaks down.

2.1 The unpolarized cut: the wave map, and a conserved handedness

The cut above is *linearly polarized*, and by the criterion of §6 that makes it achiral: its single polarization is pinned to a fixed axis by the residual T^2 . The first chiral member of the reachable

sector is therefore not this one but its unpolarized generalization, which we now build—restoring the second polarization as the off-diagonal ω of the torus block,

$$ds^2 = e^{2(\gamma-\psi)}(-dt^2 + dz^2) + e^{2\psi}(dx + \omega dy)^2 + R^2 e^{-2\psi} dy^2, \quad (8)$$

with ψ, γ, ω, R functions of (t, z) and $\omega \equiv 0$ returning (2). The torus block still has determinant R^2 , so R remains the area element, and $G_{ab} + \Lambda g_{ab} = 0$ still has exactly six nonvanishing components: the conformal-constraint sector and the torus block in which the second polarization lives.

The two polarizations are a wave map into the hyperbolic plane. In the variables

$$P = 2\psi - \ln R, \quad Q = \omega, \quad (9)$$

the torus block's equations are, identically,

$$(R P_t)_t - (R P_z)_z = R e^{2P}(Q_t^2 - Q_z^2), \quad (10)$$

$$(R e^{2P} Q_t)_t - (R e^{2P} Q_z)_z = 0, \quad (11)$$

which is the harmonic-map system for a map from the (t, z) plane, with density R , into the target $dP^2 + e^{2P}dQ^2$ —a metric of Gaussian curvature -1 , the hyperbolic plane^R. *The two polarizations are therefore not two decoupled fields but one point moving on a negatively curved surface*, the polarized case being the geodesic $Q = \text{const}$ and the turning of the polarization plane being motion off it.

And Λ is absent from that sector. Both of (10)–(11) hold with no cosmological term: Λ enters only the area and conformal equations. This is worth stating because it explains the source in (3) above. With $Q \equiv 0$ the identity $(R P_t)_t - (R P_z)_z = 2[(R \psi_t)_t - (R \psi_z)_z] - (R_{tt} - R_{zz})$ is exact, so the vanishing of (10) is precisely the polarized WAVE equation minus the AREA equation, $2\Lambda R e^{2(\gamma-\psi)} - 2\Lambda R e^{2(\gamma-\psi)} = 0$. *In the wave-map variable the graviton propagates freely and Λ drives only the area*; the source in (3) is the area equation, changed variables, and it is the second polarization that forces the variable in which this is visible.

The handedness is then the disconnected component of the target's isometry group, and it carries a conserved charge. The helicity-flipping parity of §6—the transverse reflection $x \mapsto -x$ —sends $g_{xy} \mapsto -g_{xy}$ and fixes every other component of (8), which in the variables (9) is exactly $Q \mapsto -Q$ with P fixed. That is an isometry of $dP^2 + e^{2P}dQ^2$, and its differential has determinant -1 : it reverses orientation, so it lies in the component of $\text{Isom}(\mathbb{H}^2)$ that the identity component does not reach. *No connected target isometry identifies the two handednesses.* On the homogeneous reduction— z -independent, hence a three-dimensional abelian isometry with spacelike orbits, among the classes the range paper lists as reachable [7]—equation (11) integrates once,

$$c = R e^{2P} Q_t = \text{const}, \quad (12)$$

and the parity acts on this *twist* as $c \mapsto -c$. *So on the reachable sector the graviton's handedness is the sign of a conserved charge, and the substrate's orientation parity is exactly its sign flip*, with $c = 0$ the polarized, achiral cut of §2. Solutions with $c \neq 0$ exist and their polarization turns, the first integral (12) reproduced along them^R.

That reduction is homogeneous; the member the criterion is really about is the propagating one, and it too is explicit. Restoring the z -dependence, the pair (10)–(11) is an *unconstrained* wave map: the two remaining Einstein equations fix γ by a first-order quadrature whose integrability $\partial_t \gamma_z = \partial_z \gamma_t$ holds identically on shell, so every (P, Q) Cauchy datum integrates to a genuine vacuum member and the wave carries no hidden constraint. An explicit inhomogeneous single-helicity travelling datum—two propagating modes coupled through the $\mathbb{H}^2$ metric, not the homogeneous reduction—is then evolved to the roundoff floor with fourth-order convergence, and on it the polarization plane genuinely turns. Written in the orthonormal $\mathbb{H}^2$ -frame

strain-rates $(h_+, h_\times) = (P_u, e^P Q_u)$ —the plane wave’s two polarizations in the wall limit—the criterion of §6 is the winding $\chi = \oint d\arg(h_+ + ih_\times)$, and it is nonzero with a *definite* sign on the turning member, identically zero on the polarized cut $Q \equiv 0$ where the axis is fixed, and reversed by the parity $Q \mapsto -Q$: the two helicities are exhibited as parity images *on a propagating wave*, the Cauchy problem the homogeneous reduction had left open^R.

What this adds to the handedness argument of §6 is a third instrument, and the one that lies inside the construction’s own reach. The index theorem on the compact face and the polarization computation on the type-N plane wave both establish that a chirality can live only on the component no connected action reaches; but the plane wave is a geometry the operator provably cannot generate, its full isometry algebra being a five-dimensional Heisenberg algebra where $\mathfrak{so}$ ’s unipotent radical is abelian [7]. Here the same conclusion is a property of a wave map’s *target*, on a cut the operator does generate—and it upgrades the statement from a feature of a geometry to a conserved charge of a field.

And the sector that criterion was always about—a fermion on this member—is built here rather than deferred to the matter paper’s bound modes. Put a massless Dirac field on (8) and separate on the two Killing directions. The twist is then not merely analogous to a spinor’s chirality: *it is a component of the connection the spinor moves in.* The orthonormal coframe $e^0 = e^A dt$, $e^1 = e^A dz$, $e^2 = e^\psi(dx + \omega dy)$, $e^3 = R e^{-\psi} dy$ with $A = \gamma - \psi$ carries $\omega_{t23} = -\frac{1}{2} c e^{-2\psi}$ and $\omega_{z23} = -\frac{1}{2} c_z e^{-2\psi}$, with c the conserved twist (12) itself. Rescaling by $\Omega = e^{-A/2} R^{-1/2}$ —which is exactly the leaf’s own volume element, $\Omega^2 = 1/\sqrt{h}$ —removes every background-derivative term and leaves four:

$$\gamma^0 \partial_t + \gamma^1 \partial_z + i m_x \gamma^2 + i m_y \gamma^3 + i b_z \gamma^5 \gamma^1 + i b_t \gamma^5 \gamma^0 = 0, \quad (13)$$

where $m_x = e^{A-\psi} k_x$ and $m_y = e^{A+\psi} (k_y - k_x \omega)/R$ are the two Killing momenta seen as masses on the reduction and $(b_t, b_z) = (c_z, c) e^{-2\psi}/4$ is the twist, *carried on γ^5 alone*. Two things follow, and the first is what the frontier item asks for. *The principal part is $e^{-A}(\gamma^0 \partial_t + \gamma^1 \partial_z)$, independent of the twist and of the transverse momenta*, so the characteristic matrix $\gamma^0 \gamma^1$ is Hermitian with eigenvalues exactly ± 1 : the characteristics are the light cone, on the inhomogeneous member as much as the homogeneous one, and nothing is trapped. *The sector propagates*—against the matter sector’s wall modes, which bind precisely because a superpotential changes sign, and which have no counterpart here because there is no wall in z [11]. The two norms that paper found to disagree agree here, being the same integral: the factor that flattens the reduced operator is the leaf’s volume element, and this member carries no horizon in the (t, z) sector at which the two could part. *Second, the chirality is the twist’s sign.* At vanishing Killing momenta (13) splits on γ^5 into $E = \pm(k - b)$ for one chirality and $E = \pm(k + b)$ for the other—*opposite momentum shifts*—so the orientation parity is a symmetry only as the joint operation $c \mapsto -c$ with $\gamma^5 \mapsto -\gamma^5$, and neither half alone is. Evolved on a vacuum- Λ member the two chiralities accumulate a relative phase $2 \int b dt$, identically zero on the polarized cut $c = 0$ and reversing with the sign of c . *So the graviton’s handedness and the fermion’s are one datum on this background*: what §6 argues from two sectors returning the same parity is here a single field’s own dispersion relation. The locking is exact in the Killing zero mode—which the torus always carries—and broken by the transverse momenta, which are the terms that break chirality in any case^{RL}.

3 The nonlinear $\Lambda > 0$ regime: the de Sitter attractor and classical admissibility

The $\Lambda > 0$ regime is where the dynamics is most distinctively CR’s, and it can be carried through. The model has an exact isotropic de Sitter background, and that background fixes the foliation on which the nonlinear evolution runs. Isotropy is the equality of the three spatial

scale factors— $\gamma = 2\psi$ and $R = e^{2\psi}$ —and under it the homogeneous field equations (3)–(6) reduce consistently (clean residuals) to a single constraint and a single dynamical law,

$$\psi_{tt} = \psi_t^2 \quad (\text{ENERGY}), \quad 3\psi_t^2 = \Lambda e^{2\psi} \quad (\text{AREA} = \text{WAVE} = \gamma\text{-eq}), \quad (14)$$

solved by $\psi_t = \sqrt{\Lambda/3} e^\psi$. The lapse is $N = e^{\gamma-\psi} = e^\psi$, so the scale factor $a = e^\psi$ and cosmic time $d\tau = N dt = a dt$ give $a(\tau) = a_0 e^{H\tau}$ with $H^2 = \Lambda/3$ —exact de Sitter. The internal area-clock $R = t$ that organizes the $\Lambda = 0$ Gowdy system is inconsistent here (AREA then forces $0 = 2\Lambda t e^{2(\gamma-\psi)} \neq 0$); the substrate’s de Sitter cosmic time is the clock that replaces it, established positively and not merely by the failure of the alternative. The graviton is the *anisotropic departure* from this background, and the question of whether its nonlinear back-reaction *forces* a new (quantum) structure or only *admits* one—as the clean $\Lambda = 0$ case does—is the question of that departure’s evolution on the cosmic-time foliation.

In the homogeneous sector the answer is exact. Writing the shear shape $s = \psi - \frac{1}{2} \ln R$ (so $a_x/a_y = e^{2s}$, with $s = 0$ on the background), the combination $\text{AREA} - 2 \text{WAVE}$ cancels the Λ source identically and leaves a conserved quantity,

$$C_0 := R_t - 2R\psi_t = \text{const}, \quad (15)$$

the canonical shear charge^R ($C_0 = -\frac{1}{2}(p_\gamma + p_\psi)$, verified on shell). Since $2Rs_t = -C_0$, in cosmic time—with spatial volume $V = R e^{\gamma-\psi}$ —the shear rate obeys $ds/d\tau = -C_0/2V$: as the background expands, $V \rightarrow \infty$ and the anisotropy rate decays as the inverse volume. This is *cosmic no-hair* [23]—the isotropic de Sitter background is an attractor, the shape freezing as the expansion-rate anisotropy redshifts away. The homogeneous nonlinear back-reaction thus admits clean classical dynamics: a conserved charge and a monotone decay, with nothing forcing a new structure.

The propagating mode admits as well, at linear order. Linearizing the wave equation (3) for $\delta\psi(t, z)$ on the de Sitter background and passing to cosmic time gives

$$\delta\ddot{\psi} + 3H \delta\dot{\psi} + \frac{k^2}{a^2} \delta\psi + 2\Lambda \delta\psi = 0, \quad (16)$$

a de Sitter wave equation: Hubble friction $3H$, a gradient term k^2/a^2 that redshifts away, and an effective mass $m^2 = 2\Lambda = 6H^2$ (principal series) in this fixed-background (transverse-traceless) truncation. This is a massive scalar on de Sitter and admits a clean, unitary Bunch–Davies quantization [24]. Restoring the constraint back-reaction—the metric perturbations $\delta\gamma, \delta R$ the truncation drops—and reducing to the gauge-invariant combination Q yields, in conformal time, the massless minimally-coupled de Sitter Mukhanov equation [25]

$$W'' + \left(k^2 - \frac{2}{t^2}\right)W = 0, \quad (17)$$

with no $a^2 m^2$ term. The physical effective mass is therefore exactly zero^R—the apparent shift $6H^2 \rightarrow 4H^2$ is a gauge/truncation artifact, the $\propto a$ solution being pure residual gauge—and the gauge-invariant perturbation Q is bounded—its two super-horizon branches going as a^{-3} and as a constant, the latter the standard frozen mode, so that the a^{-2} decay belongs to the rescaled variable $W = a \delta\psi$ in which the equation is written rather than to Q itself^{RL}. The propagating mode is thus a healthy massless de Sitter scalar, with no ghost, tachyon, or runaway: the admissibility is established at the gauge-invariant level by computation, not merely asserted robust to an ambiguous shift.

These computations, with the canonical structure of §2 already in hand, settle the classical question. The full nonlinear Gowdy–de Sitter system is a genuine true-Hamiltonian system—Hamilton’s equations from the canonical $\mathcal{H}$ reproduce the full evolution—and it is exact vacuum- Λ general relativity, so its energy and momentum equations (5)–(6) are the first-class Hamiltonian and momentum constraints of GR. A first-class constrained system evolves consistently to

all orders by the contracted Bianchi identity: there is no classical dynamical obstruction at any order. Two exact structures sharpen this beyond the Bianchi count. First, the homogeneous shear charge $C_0 = R_t - 2R\psi_t$ extends to the full propagating system: AREA – 2 WAVE is identically a continuity equation, $\partial_t(R_t - 2R\psi_t) = \partial_z(R_z - 2R\psi_z)$ with the Λ source cancelling, so $\int(R_t - 2R\psi_t) dz$ is conserved in the full inhomogeneous model, not only the homogeneous sector. Second, the graviton’s reduced energy $e_{\text{grav}} = 2R(\psi_t^2 + \psi_z^2) \geq 0$ is positive-definite, so the propagating sector is ghost-free to all orders. These close the ghost and zero-mode run-aways exactly. The one structure they do not by themselves close is a propagating nonlinear parametric resonance, and here the natural monotone instrument fails for a precise reason: the total field energy grows as $2\Lambda a^4$, carrying the background’s own expanding energy, so no total energy is monotone. The right object is the gauge-invariant *perturbation* energy, which decays as a^{-2} with the monotone (never periodic) coefficient of (17), excluding single-mode resonance at linear order and isolating the residual to nonlinear mode–mode transfer against the de Sitter detuning. Every tractable sector—the homogeneous nonlinear back-reaction and the linearized propagating mode—returns admissibility. The verdict, recorded at the scope it is earned:

Proposition 1. *In the symmetry-reducible sector the nonlinear $\Lambda > 0$ dynamics of the de Sitter cut admits a clean, consistent classical evolution, governed by a deparametrized true Hamiltonian on the substrate’s cosmic foliation, with no obstruction at any order and unitary quantization in every regime that can be checked. No feature of the continuous dynamics forces a quantum structure; the dynamics admits quantization, as the $\Lambda = 0$ case does.*

The residual just isolated—whether the propagating nonlinear evolution no-hairs away rather than resonating into runaway—is exactly the future stability of de Sitter, and at the level of the physics it is a settled theorem rather than an open frontier. Friedrich proved the nonlinear stability of de Sitter in vacuum with $\Lambda > 0$: data near de Sitter have geodesically-complete maximal developments asymptotic to it, a proof of cosmic no-hair in the vacuum case [19]. His method—the conformal field equations, trading the global-in-time problem for a local one at $\mathcal{J}^+$ —is also why no exact energy in this model closes the question: energy is the wrong instrument, and the failure of any monotone energy above is the expected shape of that fact, not a sign of openness. Friedrich is a small-data result, exactly the perturbative regime of the propagating graviton, so it settles the in-regime all-orders stability directly; Andréasson and Ringström extend cosmic no-hair and future stability to *all* data in the T^3 -Gowdy class with $\Lambda > 0$ [20]—with Vlasov matter, a corroboration rather than a vacuum theorem—and the one non-generic exception, the Nariai branch, unstable within the Gowdy class [21], is precisely the locus the range construction already isolates as the no-pivot seam [7]. The classical propagating no-hair therefore holds for generic data, externally grounded, with the non-generic Nariai branch the genuine boundary. This is convergence across results—vacuum small-data, all-data Gowdy with matter, vacuum Nariai-genericity—rather than a single theorem covering the exact vacuum polarized Gowdy- Λ case, and the in-model computations above are the right in-model rigor the theorems ground rather than replace.

Two cautions fix the scope. This is not a constructed full non-perturbative quantization—that object is neither built here nor needed for the verdict; and it is not the claim that no structure could ever be forced, only that no *classical dynamical* obstruction exists, so that any forcing must live elsewhere than in the continuous evolution. The one remaining candidate is the discrete root structure of the construction—the horizon cubic’s S_3 permutation symmetry and its A_2 root system (§7)—a separate, discrete object rather than a feature of the continuous dynamics, and developed in the algebroid construction [12] rather than here. Within the dynamics proper the verdict stands: the cut bends by a true-Hamiltonian flow that admits quantization throughout the symmetry-reducible sector.

4 Where the bend lives: the Type-I edge of the stratification

The range paper classifies the reachable geometries by the symmetry a cut retains—read infinitesimally, the subalgebra of the substrate isometry that fixes the cut, the isotropy [7]. This isotropy is the cut-fixing *substrate* isotropy; on the symmetry-reducible sector it coincides with the generated geometry’s own isometry (the residual symmetry there is inherited from the substrate isometry that sweeps the cut), and it is this that falls, through the Type-D and Type-I classes, to zero at the wall (Type N). At the wall the two notions part company—a point we will need: the substrate isotropy is zero, while the Type-N geometry retains its own (large) plane-wave isometry that is not substrate-inherited. The isotropy is maximal at the vacuum cut (the de Sitter / FLRW stratum). The confined wave (2) sits at a definite stratum, and we can name it exactly.

Proposition 2. *The polarized Gowdy–de Sitter metric (2) with a generic profile $\psi(t, z)$ admits exactly two Killing vectors, ∂_x and ∂_y^R , and so occupies the two-Killing-vector (Type-I) stratum—the last confined stratum before the wall.*

Verification. The metric coefficients depend only on (t, z) , so ∂_x and ∂_y are manifestly Killing. For a generic propagating profile—taking, e.g., $\psi = \sin t \cos z$ with R, γ the corresponding solution—the candidate additional symmetries all fail: the Lie derivative of the metric along ∂_t , along ∂_z , and along the boost $z\partial_t + t\partial_z$ are each nonzero. The isometry is therefore exactly the two-torus $\langle \partial_x, \partial_y \rangle$, isotropy dimension two. $\square$

This places the dynamics where the range paper said the confined radiation lives: “intermediate symmetry confines radiation (the Type-I Einstein–Rosen and Gowdy waves, on the edge)” [7]. The wave is the worked first *bend* at the last confined stratum: a single residual symmetry (the T^2) still pins the polarization to a fixed global orientation, so the wave is self-consistent only while it propagates transverse to that orientation. The dynamics is then the motion of the cut along the strata: as one descends from the symmetric vacuum, the bend turns on, generated by the true Hamiltonian, and the confined Gowdy–de Sitter wave is the explicit instance at the edge.

5 The wall: a generative boundary, not a metric singularity

The loss of the last Killing vector—isotropy from two to zero—is the wall: the Type-N, freely radiating boundary where no global sweep orientation survives and the polarization must re-orient from place to place [7]. Two questions stand at the wall. Does the dynamics simply continue across it? And is the wall a *metric singularity*—a place where, as at a black-hole or cosmological horizon, a clock can be re-founded by the Null-Boundary Correspondence [2, 8]? We settle both.

A metric singularity, in the precise sense of [2], is a collapse of the metric’s *measure*: a null hypersurface along whose generators the spatial extent contracts to zero, so that two events that are null-separated ($\Delta s^2 = 0$) and spatially coincident ($\Delta x = 0$) are assigned vanishing temporal separation ($\Delta \tau = 0$)—distinct, causally ordered points the metric measures as one place at one instant. A Killing horizon is the worked sufficient case, not the definition: for Schwarzschild–de Sitter the static Killing field $\zeta = \partial_t$ has $|\zeta|^2 = g_{tt} = -f$, which vanishes at each horizon $f = 0$, so its orbits become the null generators of zero invariant spatial extent and the measure-collapse hypotheses are met, while the Kretschmann scalar $K = 48M^2/r^6 + 24/\alpha^4$ stays finite there—the finite-curvature species. Worldlines pass through, and the freed Killing direction is reassigned as the new timelike clock—this is cosmogenesis, the re-founding of a flat- Λ CDM clock on the reassigned horizon null direction [8].

Proposition 3. *The wall is not a metric singularity of either species. Its metric is non-degenerate^R—there is no null hypersurface along whose generators the spatial extent contracts*

dt, $e_1 = e^A dz$, $e_2 = e^{\psi}(dx + \omega dy)$, $e_3 = R e^{-\psi} dy$ ($A = \gamma - \psi$) the connection carries $\omega_{t,23} = -(1/2) c e^{-2\psi}$ and $\omega_{z,23} = -(1/2) c_z e^{-2\psi}$, with $c = R e^{2P} Q_t$ P11's own conserved twist – asserted as exact symbolic zeros. (2) THE FRICTION IS EXACTLY REMOVABLE and the whole reduced operator is FOUR TERMS: $\gamma^0 dt + \gamma^1 dz + i m_x \gamma^2 + i m_y \gamma^3 + i b_z \gamma^5 \gamma^1 + i b_t \gamma^5 \gamma^0 = 0$, with m_x, m_y the Killing momenta as masses and $(b_t, b_z) = (c_z, c) e^{-2\psi}/4$ the twist – carried on γ^5 alone, an AXIAL vector – asserted as an exact matrix identity. (3) IT PROPAGATES – the question the item asks. The principal part $e^{-A}(\gamma^0 dt + \gamma^1 dz)$ carries neither the twist nor the momenta, so the characteristic matrix $\gamma^0 \gamma^1$ is hermitian with eigenvalues exactly ± 1 : the characteristics ARE the light cone and nothing binds, against P14's wall modes which bind on a superpotential's sign change. Measured: packet centroid speed 1.000000. (4) THE TWO NORMS AGREE HERE. $\Omega^2 = 1/\sqrt{h}$ exactly, so the flattening factor of the reduced operator IS the leaf's own volume element and the conserved spacetime Dirac norm and the leaf proper-measure norm are the SAME integral – P14's disagreement is at horizons this member does not have. (5) THE CHIRALITY IS THE TWIST'S SIGN. At vanishing Killing momenta the operator splits on γ^5 into $E_R = \pm(k - b)$, $E_L = \pm(k + b)$, opposite shifts, so the orientation parity is a symmetry only as the JOINT operation $c \rightarrow -c$ with $\gamma^5 \rightarrow -\gamma^5$. Measured on a genuine vacuum-Lambda background: relative phase 0.326988 against $2 \int b dt = 0.326988$). *Computes*: coframe asserted to reproduce eq:unpolarized; connection asserted antisymmetric in all 64 components; the two twist identities asserted exact; the four-term Clifford form asserted an exact matrix identity; all five nonzero components of $G + \Lambda g$ asserted to vanish IDENTICALLY on the integrated background; characteristic eigenvalues asserted ± 1 ; speed asserted within $1e-4$ of 1; norm asserted conserved to $1e-9$; relative phase asserted equal to $2 \int b dt$ within $1e-4$ and asserted LARGE (> 0.3). SEVEN CONTROLS, each shown to fire: dropping ω from e_2 returns the polarised cut; the twist identity with $1/3$ for $1/2$ is FALSE; Ω without $R^{-1/2}$ leaves R -derivatives; the Clifford form with $c/3$ for $c/4$ is FALSE; doubling γ_t breaks the $[00]$ Einstein component; $k_x = 0.9$ gives speed $0.926655 < 1$, so the speed measurement CAN fail; a non-antihermitian term drifts the norm by $2.2e-1$; and $c=0$ gives relative phase 0 while $c \rightarrow -c$ reverses it *Bound*: Massless Dirac TEST field: no back-reaction, no quantisation, no mass term. Algebra general in (t, z) ; numerics on the HOMOGENEOUS reduction, where c is conserved. BOUNDARY MEASURED, not assumed: the chirality lock is EXACT in the Killing zero mode and BROKEN by transverse momenta – at $k_x = 0.9$ the phase is 0.370120 against 0.326988 – which is the expected physics, since m_x and m_y are mass terms and a mass breaks chirality; the torus always carries $k_x = k_y = 0$. NOT CLAIMED: any Standard-Model identification, any generation count, the compact-face gauge sector, or the interacting tower – separate items, still open. No new γ^5 identification is claimed; P11's is used, not re-derived *Run*: `python3 receipts/P11_dynamics_paper/C50_the_dirac_sector_propagates_and_its_chirality_is_the_twists_sign.py`

I2.the_stratification_three_of_four_are_worked [OK]
sec:gowdy (the matter dynamics is STRATIFIED by symmetry and three of four strata are worked; the unbuilt one is the stratum with no continuous isometry). *Computes*: asserts the closed form, the Gowdy exhibition and its Type-I-edge placement, the walls character, and that P11 carries zero beyond-the-wall content (13 checks) *Bound*: NOT-A-PAPER-CLAIM — narrows L-174 and routes a citation *Run*: `python3 receipts/L174_general_matter_dynamics/I2_the_stratification_three_of_four_are_worked.py`

I3.the_identification_is_general_and_the_shear_count_is_the_gap [OK]
sec:gowdy (P11s identification is a GENERAL ADM identity, so the two Killing vectors buy a COUNT and not the content; the dark region is the selection among five-component shears). *Computes*: derives $K^2 - K_{ij} K^{ij} = (2/3)\theta^2 - \sigma^2$ symbolically with no symmetry assumed, and asserts the minus sign on σ^2 (14 checks) *Bound*: NOT-A-PAPER-CLAIM — first purchase on the unworked stratum *Run*: `python3 receipts/L174_general_matter_dynamics/I3_the_identification_is_general_and_the_shear_count_is_the_gap.py`

N1.the_nonlinear_freedom_is_the_general_first_class_structure [OK]
prop:admit (L-174 (1)'s nonlinear remainder splits: the FREEDOM (σ^{TT} stays free) is the contracted Bianchi identity, symmetry-free so it covers the general leaf not only P11's Gowdy model; P11's shear-charge and reduced-energy are the separate STABILITY layer, routed to Friedrich/A-R). *Computes*: checks general no-isometry TT data preserves the momentum constraint, σ^2 is a positive $O(\epsilon^2)$ scalar sourcing the energy sector, $\nabla_\mu G^\mu{}_\nu = 0$ EXACTLY on the inhomogeneous all-orders leaf, and P11's freedom vs stability layers

at source (19 checks) *Bound*: NOT-A-PAPER-CLAIM — informs vein L-174 (1)'s nonlinear remainder and L-165 (PO-6); the conversion runs by 'F5's stated procedure *Run*: python3 receipts/L801_nonlinear_shear_freedom/N1_the_nonlinear_freedom_is_the_general_first_class_structur

I10_the_constraint_propagation_is_symmetry_free [OK]
sec:nonlinear (cc54s L-801 verified independently: the contracted Bianchi identity holds to $O(\epsilon^2)$ on a two-wave metric with NO Killing vector, so the FREEDOM half generalises and only STABILITY is Gowdy-scoped). *Computes*: builds Christoffels, Riemann, Ricci and G to second order on a no-Killing-vector metric and contracts the Bianchi identity in all four components (12 checks) *Bound*: NOT-A-PAPER-CLAIM — verifies cc54 on this tree; the vein verdict is unseated *Run*: python3 receipts/L174_general_matter_dynamics/I10_the_constraint_propagation_is_symmetry_free

S1_the_stability_bracket_closes_the_needed_cell_is_covered [OK]
sec:nonlinear (L-245 closed against the literature: the stability bracket is a 2x2 grid (symmetry x data-size) with ONE open cell — (general, all-data) = the cosmic no-hair conjecture, general GR's open problem not CR's; the cell CR needs (vacuum perturbative graviton) is covered by Friedrich, as P11 itself states). *Computes*: anchors P11's routing and its "exactly the perturbative regime of the graviton" line, builds the coverage grid with general-subsumes-symmetric, asserts the unique open cell and CR's covered cell, adds Rodnianski-Speck which the corpus does not cite (10 checks) *Bound*: NOT-A-PAPER-CLAIM — closes the bracket on L-245 (the literature horn); informs L-165 (PO-6); whether CR needs large-data is a construction question, not settled here *Run*: python3 receipts/L802_stability_bracket/S1_the_stability_bracket_closes_the_needed_cell_is_covered

S2_the_regime_question_is_settled_by_the_corpus [OK]
sec:nonlinear (the regime question is settled by the corpus rather than by a judgement: P11 places the propagating graviton in the perturbative regime, the stratus free data IS that graviton, and the wall is where it is freed). *Computes*: asserts P11s perturbative-regime sentence and cor:radiations naming at source, and rests on cc54s S1 grid which is present on this tree (10 checks) *Bound*: NOT-A-PAPER-CLAIM — closes L-245, scoped per r2505 *Run*: python3 receipts/L802_stability_bracket/S2_the_regime_question_is_settled_by_the_corpus.py

S1_the_nariai_closed_form_extends_as_a_series_in_the_horizon_separation [OK]
sec:nonlinear (A9 (PO-6(c)): the Nariai closed form extends off the degenerate member — $dS_2 \times S^2$ with both radii $r_n = \alpha/\sqrt{3} = 1/\sqrt{\Lambda}$ is the leading term, the geometry is a series in the horizon separation $\epsilon \sim \sqrt{M_N - M}$, and $f''(r_n) = -6/\alpha^2$ is its leading coefficient). *Computes*: solves $f=0$ & $f'=0$ for the Nariai member symbolically, recovers $r_n/M_N/f''(r_n)$, shows the throat is $dS_2 \times S^2$ with equal radii and the horizon separation goes as $\sqrt{M_N - M}$ (8 checks) *Bound*: NOT-A-PAPER-CLAIM — A9; informs L-165 (PO-6); the closed form extends *Run*: python3 receipts/L808_nariai_extension/S1_the_nariai_closed_form_extends_as_a_series_in_the_horizon

S1_the_L553_pontryagin_value_reproduces_at_its_own_pinned_point [OK]
sec:chirality (PO-6 (OWED #518, routed to cc54 at c54.219): STATE the (H, k, ω) at which L553's $R \sim R = 4.977310$ was evaluated so it is reproducible. The point is the DEFAULTS of L553's own numeric pontryagin_num: $(H, k, \omega) = (0.5, 1.5, 1.4)$ at $X_p = (t, x, y, z) = (1/3, 0, 0, 0.4)$, $\epsilon \sim 0$. Imports L553's function and recovers $+4.977310$ (circular+), -4.977310 (opposite handedness), 0 (linear), closed form $147 e^{(-1/6)}/25 = 4.977313$. Pins the second point so the r2794 "two parameter points, not two answers" is checkable: L-821's $+4.5$ is at $(0.5, 1, 1.5)$, this at $(0.5, 1.5, 1.4)$; the invariant is degree-4 homogeneous in (H, k, ω) , so different points give different values). *Computes*: imports L553's pontryagin_num, evaluates the three polarisations at its default $(H, k, \omega)/X_p$, checks the value, the sign flip, the closed form, and the distinctness from L-821's point (3 checks) *Bound*: NOT-A-PAPER-CLAIM — reproducibility metadata for L-553/L-821's Pontryagin figure; only the STRUCTURE $(0/\pm/\text{flip})$ is invariant, the number is a field at a chosen point *Run*: python3 receipts/L825_pontryagin_point/S1_the_L553_pontryagin_value_reproduces_at_its_own_pinned_p

P03_acceleration_is_slice_curvature [OK]
sec:curvature · rem:twocritical · the bend (on ANY member of the energy family (E cancels: $(dr/d\tau)^2 = E^2 - f$, and E is a constant of the motion) $d^2r/d\tau^2 = -f'/2 = r^*K \cdot G$ identically, so THE SIGN OF THE SLICE'S GAUSSIAN CURVATURE IS THE SIGN OF THE COMOVING ACCELERATION — and the flat locus is the deceleration-to-acceleration turnover, which on Nariai is the seam and in standard terms is $\rho_m = 2 \rho_\Lambda$ ($\cosh^2 w = 2 \Rightarrow \sinh w = 1/\sqrt{2}$, the seam's phase). A recent cosmological epoch: 7.06 Gyr at $H_0 = 73$, 7.65 at 67.4, preceding matter-Lambda equality [borrowed from P3 · P7 · P8]). *Computes*: the identity asserted; the $K \cdot G$ zero

solved and matched to $(M \alpha^2)^{1/3}$; $\text{csch}^2 w = 2 \Leftrightarrow \sinh w = 1/\sqrt{2}$ proved by algebra after simplify() failed on the closed forms; epochs at both H0 *Bound*: bound: identities on the Narai E=1 worldline; no causal claim between seam and turnover – they are one locus; dating inherits H0 *Run*: python3 receipts/P03_SdS_slicing/P03_acceleration_is_slice_curvature.py

P09_the_wall_is_not_the_loss_of_symmetry [OK]
cor:wall / cor:radiation / thm:range · sec:chirality (**THE RANGE'S BOUNDARY IS NOT THE LOSS OF CONTINUOUS SYMMETRY, AND THE PAPER'S OWN BEYOND-THE-WALL EXEMPLAR IS THE WITNESS.** ** THE TYPE-N PLANE WAVE HAS FIVE KILLING VECTORS AND IS VACUUM ** – more continuous symmetry than Schwarzschild's four, and no matter at all – so BOTH readings cor:wall gives of the boundary are falsified by the geometry cor:radiation sends beyond it. DERIVED HERE: (1) for $ds^2 = 2 du dv + H du^2 + dx^2 + dy^2$ the ansatz $\xi.f = f^a(u) d_a - (f.\dot{x}) d_v$ is Killing **iff $f.\ddot{d} = A f$, $A = [[h+,hx],[hx,-h+]]$ **, every other component of $\text{Lie}_\xi g$ vanishing identically – a 4-dim solution space plus $d_v = \text{FIVE}$, checked on a TURNING (chiral) profile with Ricci = 0; (2) $[\xi.f, \xi.g] = W(f,g) d_v$ with $dW/du = 0$ on shell $\Leftrightarrow$ the algebra is the 5-dim HEISENBERG algebra**, derived algebra the 1-dim centre; (3) **the orbits are 3-dim and NULL** (no Killing vector has a d_u component; $g(d_v, d_v) = 0$); (4) $\text{so}(5,1)$'s unipotent radical is ABELIAN** ($K_i = M_{0i} + M_{5i}$ commute pairwise, $K^3 = 0$), so no Heisenberg algebra of unipotent isometries embeds – the wave's FULL algebra is not a sweep; (5) **but its abelian R^3 subalgebras DO embed**, so the plane wave SATISFIES thm:bound/thm:range's stated condition while cor:radiation places it beyond the wall – **read literally the two are inconsistent**; (6) **the repair is thm:bound's OWN second clause**: the boundary is the loss of a symmetry whose ORBITS CAN CARRY A LEAF, the two criteria coinciding across every class the paper fills and coming apart exactly on the null orbits; (7) **and the chirality consequence runs in the construction's favour**: on the Gowdy form d_x, d_y are Killing and spacelike and the reflection $x \rightarrow -x$ is an isometry iff $g_{xy} = 0$, i.e. iff POLARIZED – so the unpolarized turning wave is chiral and lies IN the reachable sector by P9's own ground. **Gravitational chirality is achieved INSIDE the range.** [borrowed from P9 / P11]. *Computes*: the Killing condition derived symbolically and reduced to the oscillator system; Ricci asserted zero on the turning profile; the Wronskian asserted constant on shell; the orbit asserted null; $\text{so}(5,1)$ asserted 15-dimensional with the four K_i asserted pairwise-commuting and nilpotent; and the Gowdy reflection condition asserted equivalent to $Q = 0$ *Bound*: ** THIS CHANGES A CHARACTERISATION, NOT A RESULT. ** No reachable class, filling, or vacuum kernel moves; the Z₂ identification (graviton helicity and fermion γ^5 as one orientation parity) is P11's and is untouched. **And the first chiral member of the range is NAMED, not built** – the cut constructed exactly is the polarized, achiral one *Run*: python3 receipts/P09_range_paper/P09_the_wall_is_not_the_loss_of_symmetry.py

Appendix L The Knowledge Ledgers

Each claim marked ^L in the text rests on a field bake or the figure–theorem bake recorded in the named ledger, which ships with this work and carries the probes, the receipts, and the three registers kept apart — what bit, what bounced, and where the boundary is. A ledger is working apparatus, not a publication, and is cited here rather than in the bibliography. This appendix is generated from the ledger index, not maintained by hand.

`CARTAN_HOLONOMY_LEDGER.md` [field bake]

The Cartan / connections-and-holonomy field-bake ledger — what bit, what bounced, and the boundary. First of the four fields ‘L-272’'s re-survey left outstanding. ‘OWED’ 622

`FUNCTIONAL_ANALYSIS_LEDGER.md` [field bake]

The functional-analysis / unitarity field-bake ledger — the field that bounced, and the one routing fact it returned. Third of the four fields ‘L-272’'s re-survey left outstanding. ‘OWED’ 622