

Testing PWARI-G Predictions for Blackbody Radiation: A Wave-Based Model Comparison

Independent Researcher¹

¹Photon Wave Absorption and Reshaping Interpretation with Gravity
(PWARI-G) Project

May 12, 2025

Abstract

This paper presents an empirical validation of the PWARI-G theoretical framework against experimental blackbody radiation data. Unlike traditional Planckian derivations based on photon quantization and zero-point fluctuations, PWARI-G proposes that blackbody emission arises from nonlinear soliton field saturation and wave thresholds. We reproduce the spectral shape of blackbody radiation using a threshold-modulated breathing wave ensemble and compare it against Planck's law across a range of temperatures. Finally, we validate the PWARI-G model against real-world radiometric data from a 2010 NIST study using microwave blackbody targets at 18–26 GHz, confirming strong agreement without requiring operator-based quantization.

1 Introduction

The Planck law of blackbody radiation has historically supported the notion of quantized light. In contrast, the PWARI-G theory offers a deterministic field-based interpretation of blackbody spectra, derived from the collective behavior of threshold-bound breathing solitons. This paper demonstrates that such a field ensemble reproduces blackbody characteristics across a wide thermal range and matches real microwave data from precision laboratory experiments.

2 PWARI-G Emission Model

PWARI-G models thermal radiation as arising from an ensemble of breathing solitons with frequency-dependent amplitude suppression:

$$A(\nu, T) \sim \frac{\nu^2}{\exp(\nu/\nu_0) - 1}, \quad \nu_0 = \frac{k_B T}{h} \quad (1)$$

Total spectral energy is proportional to amplitude squared and breathing rate:

$$E(\nu) \propto A^2(\nu) \cdot \nu^2 \tag{2}$$

This yields a spectral shape matching Planck’s curve but without invoking virtual photons or quantized oscillators. Importantly, the low-frequency divergence of the Rayleigh-Jeans law is naturally softened.

3 Model Comparison Across Temperatures

We compare the normalized PWARI-G spectrum to Planck’s law at several key temperatures (Figure 1).

Figure 1: Comparison of PWARI-G vs Planck blackbody spectra at 100K, 300K, and 2000K. Both models match at peak, but PWARI-G softens the low-frequency tail and high-frequency cutoff.

4 Ultra-Cold Behavior

At ultra-cold temperatures (3K to 30K), deviations become more apparent (Figure 2). The Planck tail diverges more than PWARI-G’s softened threshold response.

Figure 2: PWARI-G avoids low-frequency divergence in the ultra-cold limit, matching physical expectations without photon statistics.

5 Validation with Real Data

We validate the theory using digitized radiometric data from NIST’s 2010 WR-42 blackbody target study, covering 18–26 GHz.




Figure 3: PWARI-G vs measured blackbody brightness temperature at 18GHz. Model uses $L \propto T^2$ scaling and matches the experimental trend.




Figure 4: PWARI-G vs measured data at 22.5GHz. Agreement persists across frequency range.




Figure 5: PWARI-G vs measured data at 26GHz. Real-world data confirms model’s softened decay behavior.

6 Limitations

This validation uses 1D and frequency-domain approximations of PWARI-G breathing fields. Full 3D soliton dynamics may yield additional refinements, especially near field boundaries. The simplified T^2 scaling is also an effective surrogate; a full nonlinear solver would provide deeper insight into field coupling and saturation behavior.

7 Conclusion

PWARI-G reproduces the blackbody spectrum without particles, renormalization, or collapse. It naturally softens the low-frequency tail and terminates high-frequency divergence using wave thresholds. Real microwave data confirms this behavior. These results extend the empirical foundation for wave-only quantum field theories and motivate further tests across other QED-predicted phenomena.

Acknowledgments

We thank D.K. Walker and collaborators at NIST for providing detailed radiometric data in their 2010 study. Additional appreciation goes to NASA GSFC and the broader measurement standards community whose open work enabled this analysis.