

GLM Regression (Q5L6)

Total Questions: 20

Most Correct Answers: #9

Least Correct Answers: #6

1. Poisson distribution is specified by

- 4/7 ☒ A 1 parameter
- 2/7 ☐ B 2 parameters
- 1/7 ☐ C 3 parameters
- 0/7 ☐ D Poisson distribution does not have parameters
- 0/7 ☐ E I do not know

2. The type of dependent variable in Poisson Regression is

- 1/7 ☐ A Integer
- 4/7 ☒ B Count
- 0/7 ☐ C Ratio
- 0/7 ☐ D Interval
- 0/7 ☐ E I do not know
- 2/7 ☐ F Binary

3. Overdispersion in Poisson Regression occurs when

- 5/7 ☐ A $\text{var}(Y|X) > \text{var}(Y)$
- 1/7 ☒ B $\text{var}(Y|X) > \text{mean}(Y|X)$
- 0/7 ☐ C Variance is decreasing
- 1/7 ☐ D I do not know

4. The model of Poisson Regression is

- 6/7 ☒ A $\ln(\lambda) = e^{xb}$
- 0/7 ☐ B $\ln(y) = e^{xb}$
- 0/7 ☐ C $\ln(y) = e^{xb} / (1 + e^{xb})$
- 1/7 ☐ D $\ln(\lambda) = e^{xb} / (1 + e^{xb})$
- 0/7 ☐ E I do not know

5. We can estimate Poisson Regression in R using function

- 0/7 ☐ A lm()
5/7 ☒ B glm()
0/7 ☐ C flm()
2/7 ☐ D poisson()
0/7 ☐ E I do not know

6. Which one of these is the measure for goodness of fit for Poisson Regression?(if any)

- 1/7 ☐ A Ordinal R^2
4/7 ☒ B Chi-square
0/7 ☐ C I do not know
1/7 ☐ D There are not measure for it
1/7 ☒ E Pseudo R^2

7. Which one of these is the correct interpretation of the coefficient of Poisson Regression?

- 2/7 ☐ A For a 1-unit increase in X, we expect a b1 unit increase in Y.
2/7 ☒ B For a 1-unit increase in X, we expect b1 percentage increase in Y.
3/7 ☐ C For a 1-percentage increase in X, we expect b1 percentage increase in Y.
0/7 ☐ D For a 1-percentage increase in X, we expect b1 unit increase in Y.
0/7 ☐ E I do not know

8. Count data is continuous

- 1/7 ☐ A Yes
6/7 ☒ B No
0/7 ☐ C I do not know

9. The logistic model is estimated by way of

- 1/7 ☐ A Ordinary least squares
6/7 ☒ B Maximum likelihood estimation
0/7 ☐ C Negative binomial distribution
0/7 ☐ D I do not know

10. As a result of estimation of coefficients

- 2/7 ☒ A We do not have the formula, an iterative algorithm must be used
- 2/7 ☐ B The explicit formula of coefficients exists
- 1/7 ☐ C I do not know
- 2/7 ☐ D We can obtain different values for coefficients

11. In Poisson regression...

- 2/7 ☒ A The asymptotic distribution of the maximum likelihood estimates is multivariate normal.
- 0/7 ☐ B The distribution of the maximum likelihood estimates is multivariate normal.
- 3/7 ☐ C The asymptotic distribution of the maximum likelihood estimates is multivariate Poisson distribution.
- 2/7 ☐ D I do not know

12. Pseudo R-Squared Measures are calculated based on (if any)

- 1/7 ☒ A The likelihood function
- 3/7 ☒ B Row residuals
- 0/7 ☒ C Deviance
- 0/7 ☐ D Chi-squared value
- 3/7 ☐ E I do not know

13. The formula for the raw residual is

- 5/7 ☒ A The difference between the actual response and the estimated value from the model
- 0/7 ☐ B The squared difference between the actual response and the estimated value from the model
- 2/7 ☐ C The difference between the actual response and the estimated value from the model by dividing by the standard deviation
- 0/7 ☐ D I do not know

14. Which of these is NOT the type of residuals

- 1/7 ☐ A Deviance Residual
- 1/7 ☐ B Pearson Residual
- 0/7 ☐ C Raw Residual
- 5/7 ☒ D Poisson Residual
- 0/7 ☐ E I do not know

15. In the case of intercept-only model

- 2/7 ☒ A The mean of the dependent variable equals the exponential value of intercept
- 0/7 ☐ B The mean of the dependent variable equals the intercept
- 5/7 ☐ C The mean of the dependent variable equals 0
- 0/7 ☐ D I do not know

16. $\ln(\lambda) = 0.6 - 0.2 * \text{female}$ [λ = the average number of articles] Note: $e^{(-0.2)} = 0.78$

- 3/7 ☐ A One unit increase in female brings a 0.2 decrease in $\ln(\lambda)$.
- 2/7 ☐ B Being female decreases the average number of articles by 0.78 percent
- 2/7 ☒ C Being female decreases the average number of articles by 22%
- 0/7 ☐ D I do not know

17. While running the Poisson Regression we will have never faced with the value of λ

- 3/7 ☒ A 0
- 1/7 ☐ B 1
- 2/7 ☐ C 2
- 1/7 ☐ D I do not know

18. Why does not quasi-Poisson model have AIC?

- 2/7 ☒ A Quasi-Poisson is used quasi-likelihood instead of log-likelihood estimates.
- 3/7 ☐ B Quasi-Poisson does not use iterative estimation
- 2/7 ☐ C I do not know

19. Why Poisson regression is called log-linear?

- 2/7 ☒ A Because we use a log link to estimate the logarithm of the average value of the dependent variable
- 0/7 ☐ B Because we use a log values of independent variable
- 5/7 ☐ C Because we use a log value of an independent variable is transformed to linear
- 0/7 ☐ D I do not know

20. Formulate the Null hypothesis for chi-squared and deviance test.

- 1/7 ☐ A The distance between actual and predicted values is insignificant
- 6/7 ☒ B The distance between actual and predicted values is 0
- 0/7 ☐ C There is a significant difference between actual and predicted values.
- 0/7 ☐ D I do not know