

# **Safety and Efficacy of the Xanomeline Transdermal Therapeutic System (TTS) in Patients with Mild to Moderate Alzheimer's Disease**

|   |    |
|---|----|
| 0 TITLE PAGE  | 3  |
| 1 PROTOCOL SUMMARY  | 4  |
| 2 INTRODUCTION  | 5  |
| 3 TRIAL OBJECTIVES, ENDPOINTS AND ESTIMANDS   | 6  |
| 4 TRIAL DESIGN  | 7  |
| 5 TRIAL POPULATION  | 8  |
| 6 TRIAL INTERVENTION AND CONCOMITANT THERAPY  | 10 |
| 7 DISCONTINUATION OF TRIAL INTERVENTION AND PARTICIPANT WITHDRAWAL FROM TRIAL                 | 12 |
| 8 TRIAL ASSESSMENTS AND PROCEDURES  | 13 |
| 9 STATISTICAL CONSIDERATIONS  | 15 |
| 10 GENERAL CONSIDERATIONS: REGULATORY, ETHICAL, AND TRIAL OVERSIGHT                           | 16 |
| 11 GENERAL CONSIDERATIONS: RISK MANAGEMENT AND QUALITY ASSURANCE                              | 17 |
| 12 APPENDIX: ADVERSE EVENTS AND SERIOUS ADVERSE EVENTS - DEFINITIONS, SEVERITY, AND CAUSALITY | 18 |
| 13 APPENDIX: DEFINITIONS AND SUPPORTING OPERATIONAL DETAILS                                   | 19 |
| 14 APPENDIX: GLOSSARY OF TERMS  | 20 |
| 15 APPENDIX: REFERENCES   | 21 |

# 0 TITLE PAGE

**Protocol Full Title:** \*\*\*\*\* Failed to translate reference, attribute not found \*\*\*\*\*

**Protocol Number:** H2Q-MC-LZZT

**Version:** \*\*\*\*\* Failed to translate reference, attribute not found \*\*\*\*\*

**Amendment Number:** \*\*\*\*\* Failed to translate reference, attribute not found \*\*\*\*\*

**Amendment Scope:** \*\*\*\*\* Failed to translate reference, attribute not found \*\*\*\*\*

**Compound Number(s):** \*\*\*\*\* Failed to translate reference, attribute not found \*\*\*\*\*

**Compound Name(s):** \*\*\*\*\* Failed to translate reference, attribute not found \*\*\*\*\*

**Trial Phase:** Phase II Trial

**Acronym:** \*\*\*\*\* Failed to translate reference, attribute not found \*\*\*\*\*

**Short Title:** \*\*\*\*\* Failed to translate reference, attribute not found \*\*\*\*\*

**Sponsor Name and Address:** \*\*\*\*\* Failed to translate reference, attribute not found \*\*\*\*\*

# I PROTOCOL SUMMARY

## 1.1 Protocol Synopsis

## 1.2 Trial Schema

## 1.3 Schedule of Activities

# 2 INTRODUCTION

## 2.1 Purpose of Trial

## 2.2 Summary of Benefits and Risks

# 3 TRIAL OBJECTIVES, ENDPOINTS AND ESTIMANDS

## 3.1 Primary Objectives

**Primary  
Objective**

**Primary Endpoint**

Main  
objective

Safety      The change from baseline laboratory value will be calculated as the difference between the baseline lab value and the endpoint value (i.e., the value at the specified visit) or the end of treatment observation

## 4 TRIAL DESIGN

### 4.1 Description of Trial Design

#### 4.1.1 Participant Input into Design

### 4.2 Rationale for Trial Design

#### 4.2.1 Rationale for Comparator

#### 4.2.2 Rationale for Adaptive or Novel Trial Design

#### 4.2.3 Other Trial Design Considerations

### 4.3 Access to Trial Intervention After End of Trial

### 4.4 Start of Trial and End of Trial

# 5 TRIAL POPULATION

## 5.1 Selection of Trial Population

## 5.2 Rationale for Trial Population

## 5.3 Inclusion Criteria

Patients may be included in the study only if they meet **all** the following criteria:

- [1] Males and postmenopausal females at least 50 years of age.
- [2] Diagnosis of probable AD as defined by National Institute of Neurological and Communicative Disorders and Stroke (NINCDS) and the Alzheimer's Disease and Related Disorders Association (ADRDA) guidelines (Attachment LZZT.7).
- [3] MMSE score of 10 to 23.
- [4] Hachinski Ischemic Scale score of  $\leq 4$  (Attachment LZZT.8).
- [5] CNS imaging (CT scan or MRI of brain) compatible with AD within past 1 year. The following findings are incompatible with AD:
  - a. Large vessel strokes
    - 1. Any definite area of encephalomalacia consistent with ischemic necrosis in any cerebral artery territory.
    - 2. Large, confluent areas of encephalomalacia in parieto-occipital or frontal regions consistent with watershed infarcts. The above are exclusionary. Exceptions are made for small areas of cortical asymmetry which may represent a small cortical stroke or a focal area of atrophy provided there is no abnormal signal intensity in the immediately underlying parenchyma. Only one such questionable area allowed per scan, and size is restricted to  $\leq 1$  cm in frontal/parietal/temporal cortices and  $\leq 2$  cm in occipital cortex.
  - b. Small vessel ischemia
    - 1. Lacunar infarct is defined as an area of abnormal intensity seen on CT scan or on both T1 and T2 weighted MRI images in the basal ganglia, thalamus or deep white matter which is  $\leq 1$  cm in maximal diameter. A maximum of one lacune is allowed per scan.
    - 2. Leukoariosis or leukoencephalopathy is regarded as an abnormality seen on T2 but not T1 weighted MRIs, or on CT. This is accepted if mild or moderate in extent, meaning involvement of less than 25% of cortical white matter.
  - c. Miscellaneous
    - 1. Benign small extra-axial tumors (ie, meningiomas) are accepted if they do not contact or indent the brain parenchyma.
    - 2. extra-axial arachnoid cysts are accepted if they do not indent or deform the brain parenchyma.



the caregiver

[7] Geographic proximity to investigator's site that allows adequate follow-up.

[8] A reliable caregiver who is in frequent or daily contact with the patient and who will accompany the patient to the office and/or be available by telephone at designated times, will monitor administration of prescribed medications, and will be responsible for the overall care of the patient at home. The caregiver and the patient must be able to communicate in English and willing to comply with 26 weeks of transdermal therapy.

## 5.4 Exclusion Criteria

## 5.5 Lifestyle Considerations

### 5.5.1 Meals and Dietary Restrictions

### 5.5.2 Caffeine, Alcohol, Tobacco, and Other Habits

### 5.5.3 Physical Activity

### 5.5.4 Other Activity

## 5.6 Screen Failures

# 6 TRIAL INTERVENTION AND CONCOMITANT THERAPY

## 6.1 Description of Trial Intervention

## 6.2 Rationale for Trial Intervention

## 6.3 Dosing and Administration

### 6.3.1 Trial Intervention Dose Modification

## 6.4 Treatment of Overdose

## 6.5 Preparation, Handling, Storage and Accountability

### 6.5.1 Preparation of Trial Intervention

### 6.5.2 Handling and Storage of Trial Intervention

### 6.5.3 Accountability of Trial Intervention

## 6.6 Participant Assignment, Randomisation and Blinding

### 6.6.1 Participant Assignment

### 6.6.2 Randomisation

### 6.6.3 Blinding and Unblinding

The study will be double-blind. To further preserve the blinding of the study, only a minimum number of Lilly and CRO personnel will see the randomization table and codes before the study is complete.

Emergency codes generated by a computer drug-labeling system will be available to the investigator. These codes, which reveal the patients treatment group, may be opened during the study only if the choice of follow-up treatment depends on the patient's therapy assignment.

The investigator should make every effort to contact the clinical research physician prior to unblinding a patient's therapy assignment. If a patient's therapy assignment is unblinded, Lilly must be notified immediately by telephone. After the study, the investigator must return all sealed and any opened codes.

## 6.7 Trial Intervention Compliance

ST DOCUMENT — TEST DOCUMENT — TEST DOCUMENT — Document doesn't look right? [We'll help you out!](#) — TEST DOCUMENT — TEST DOCUMENT — TEST DOCUMENT

## 6.8 Concomitant Therapy

### 6.8.1 Prohibited Concomitant Therapy

### 6.8.2 Permitted Concomitant Therapy

### 6.8.3 Rescue Therapy

### 6.8.4 Other Therapy

ST DOCUMENT — TEST DOCUMENT — TEST DOCUMENT — Document doesn't look right? [We'll help you out!](#) — TEST DOCUMENT — TEST DOCUMENT — TEST DOCUMENT

# 7 DISCONTINUATION OF TRIAL INTERVENTION AND PARTICIPANT WITHDRAWAL FROM TRIAL

## 7.1 Discontinuation of Trial Intervention

### 7.1.1 Criteria for Permanent Discontinuation of Trial Intervention

### 7.1.2 Temporary Discontinuation or Interruption of Trial Intervention

### 7.1.3 Rechallenge

## 7.2 Participant Withdrawal from the Trial

## 7.3 Lost to Follow-Up

## 7.4 Trial Stopping Rules

## **8 TRIAL ASSESSMENTS AND PROCEDURES**

### **8.1 Screening/Baseline Assessments and Procedures**

### **8.2 Efficacy Assessments and Procedures**

### **8.3 Safety Assessments and Procedures**

#### **8.3.1 Physical Examination**

#### **8.3.2 Vital Signs**

#### **8.3.3 Electrocardiograms**

#### **8.3.4 Clinical Laboratory Assessments**

#### **8.3.5 Suicidal Ideation and Behaviour Risk Monitoring**

### **8.4 Adverse Events and Serious Adverse Events**

#### **8.4.1 Definitions of AE and SAE**

#### **8.4.2 Time Period and Frequency for Collecting AE and SAE Information**

#### **8.4.3 Identifying AEs and SAEs**

#### **8.4.4 Recording of AEs and SAEs**

#### **8.4.5 Follow-up of AEs and SAEs**

#### **8.4.6 Reporting of SAEs**

#### **8.4.7 Regulatory Reporting Requirements for SAEs**

#### **8.4.8 Serious and Unexpected Adverse Reaction Reporting**

#### **8.4.9 Adverse Events of Special Interest**

#### **8.4.10 Disease-related Events or Outcomes Not Qualifying as AEs or SAEs**

## **8.5 Pregnancy and Postpartum Information**

### **8.5.1 Participants Who Become Pregnant During the Trial**

### **8.5.2 Participants Whose Partners Become Pregnant**

## **8.6 Medical Device Product Complaints for Drug/Device Combination Products**

### **8.6.1 Definition of Medical Device Product Complaints**

### **8.6.2 Recording of Medical Device Product Complaints**

### **8.6.3 Time Period and Frequency for Collecting Medical Device Product Complaints**

.

### **8.6.4 Follow-Up of Medical Device Product Complaints**

### **8.6.5 Regulatory Reporting Requirements for Medical Device Product Complaints**

## **8.7 Pharmacokinetics**

## **8.8 Genetics**

## **8.9 Biomarkers**

## **8.1 Immunogenicity Assessments**

### **8.1.1 Medical Resource Utilisation and Health Economics**

## 9 STATISTICAL CONSIDERATIONS

### 9.1 Analysis Sets

### 9.2 Analyses Supporting Primary Objective(s)

#### 9.2.1 Statistical Model, Hypothesis, and Method of Analysis

#### 9.2.2 Handling of Intercurrent Events of Primary Estimand(s)

#### 9.2.3 Handling of Missing Data

#### 9.2.4 Sensitivity Analysis

#### 9.2.5 Supplementary Analysis

### 9.3 Analysis Supporting Secondary Objective(s)

### 9.4 Analysis of Exploratory Objective(s)

### 9.5 Safety Analyses

### 9.6 Other Analyses

### 9.7 Interim Analyses

### 9.8 Sample Size Determination

### 9.9 Protocol Deviations

ST DOCUMENT — TEST DOCUMENT — TEST DOCUMENT — Document doesn't look right? [We'll help you out!](#) — TEST DOCUMENT — TEST DOCUMENT — TEST DOCUMENT

# **10 GENERAL CONSIDERATIONS: REGULATORY, ETHICAL, AND TRIAL OVERSIGHT**

## **10.1 Regulatory and Ethical Considerations**

## **10.2 Committees**

## **10.3 Informed Consent Process**

## **10.4 Data Protection**

## **10.5 Early Site Closure or Trial Termination**



# II GENERAL CONSIDERATIONS: RISK MANAGEMENT AND QUALITY ASSURANCE

## 11.1 Quality Tolerance Limits

## 11.2 Data Quality Assurance

## 11.3 Source Data

ST DOCUMENT — TEST DOCUMENT — TEST DOCUMENT — Document doesn't look right? [We'll help you out!](#) — TEST DOCUMENT — TEST DOCUMENT — TEST DOCUMENT

# **12 APPENDIX: ADVERSE EVENTS AND SERIOUS ADVERSE EVENTS - DEFINITIONS, SEVERITY, AND CAUSALITY**

## **12.1 Further Details and Clarifications on the AE Definition**

## **12.2 Further Details and Clarifications on the SAE Definition**

## **12.3 Severity**

## **12.4 Causality**

# 13 APPENDIX: DEFINITIONS AND SUPPORTING OPERATIONAL DETAILS

## 13.1 Contraception and Pregnancy Testing

### 13.1.1 Definitions Related to Childbearing Potential


### 13.1.2 Contraception

### 13.1.3 Pregnancy Testing

## 13.2 Clinical Laboratory Tests

## 13.3 Country/Region-Specific Differences

## 13.4 Prior Protocol Amendments



ST DOCUMENT — TEST DOCUMENT — TEST DOCUMENT — Document doesn't look right? [We'll help you out!](#) — TEST DOCUMENT — TEST DOCUMENT — TEST DO

# 14 APPENDIX: GLOSSARY OF TERMS

# 15 APPENDIX: REFERENCES