

AGRO-HOMEOPATHIC REPERTORY FOR PLANTS — FULL-SENTENCE VERSION

=====

SECTION 1: SOIL TYPE

=====

For alkaline soil, use Sulphur, Phosphoric acid, Ferrum phosphoricum, Nitric acid, and Ammonium carbonicum.

For sandy soil, use Silicea, Arnica montana, Acidum boracicum, Calcarea phosphoricum, Cuprum metallicum, Ferrum phosphoricum, Magnesium carbonicum, Manganum metallicum, Kalium nitricum, Kalium carbonicum, Kalium sulphuricum, Selenium, and Zincum metallicum.

SECTION 2: WEATHER / CLIMATE

=====

For cold stress, used prophylactically, give Aconite 200, Borax 12, Carbo vegetabilis 30, Psorinum 200, and Veratrum album 30.

For heat stress, used prophylactically, give Aconite 200, Arsenicum album 30, Belladonna 200, Cantharis 30, China officinalis 30, Dulcamara 30, Natrum muriaticum 200, Sol 30, Sulphur 200, and Acidum Folicum 30.

For rain, dampness, or waterlogging, give Natrum sulphuricum 30, Carbo vegetabilis 12, and Dulcamara 30.

For wind damage, give Silicea 200, Arnica montana 200, and Calendula officinalis 30.

For hail damage, give Arnica montana 200, Calendula officinalis 30, and Nux vomica 30.

SECTION 3: NITROGEN DEFICIENCY

=====

For combined Nitrogen and Potassium deficiencies, use Kalium nitricum.

For an immediate Nitrogen boost, use Ammonium nitricum.

For Nitrogen and Calcium deficiency, especially in younger plants, or for weak growth after leaching rainfall, use Calcarea nitrica.

When plants show both Nitrogen deficiency and signs of fungal infection, use Argentum nitricum.

For combined Nitrogen and Iron deficiencies, use Ferrum pernitricum.

SECTION 4: PHOSPHORUS DEFICIENCY

=====

For Phosphorus deficiency alone, which also promotes flowering and strengthens stems, use Phosphorus.

Phosphoric acid works similarly to Phosphorus and should be used when the plant lacks overall vitality.

For combined Phosphorus and Calcium deficiencies, use Calcarea phosphoricum.

For combined Phosphorus and Potassium deficiencies, use Kalium phosphoricum.

For combined Phosphorus and Magnesium deficiencies, use Magnesium phosphoricum.

SECTION 5: POTASSIUM DEFICIENCY

=====

For combined Potassium and Sulphur deficiencies, use Kalium sulphuricum.

For Potassium deficiency in young plants, use Kalium carbonicum.

For combined Potassium and Phosphorus deficiencies, use Kalium phosphoricum.

For combined Potassium and Nitrogen deficiencies, use Kalium nitricum.

SECTION 6: IRON DEFICIENCY

=====

For combined Iron and Phosphorus deficiencies, use Ferrum phosphoricum.

Ferrum muriaticum is particularly effective for iron deficiency in sandy soil.

For Iron deficiency alone, use Ferrum metallicum.

SECTION 7: SULPHUR DEFICIENCY

=====

For Sulphur deficiency, apply Sulphur after the growing season to improve overall plant health and help prevent future infestations.

For combined Sulphur and Calcium deficiencies, use Calcarea sulphuricum.

For combined Sulphur and Magnesium deficiencies, use Magnesium sulphuricum.

SECTION 8: CALCIUM DEFICIENCY

=====

Calcarea carbonicum 30 is an excellent remedy for young plants.

For young plants that grow too rapidly but remain thin and weak, use Calcarea phosphoricum.

For plants with floppy growth, easily broken stems, and tall plants that need structural support, use Calcarea fluorica.

SECTION 9: COPPER DEFICIENCY

=====

For Copper deficiency alone, use Cuprum metallicum.

For combined Copper and Sulphur deficiencies, use Cuprum sulphuricum.

SECTION 10: MANGANESE DEFICIENCY

=====

For combined Manganese and Sulphur deficiencies, which often occur together in alkaline soils, use Manganum sulphuricum.

To correct a Manganese imbalance, use Manganum metallicum.

SECTION 11: MAGNESIUM DEFICIENCY

=====

As a Magnesium booster for yellowing in younger leaves, and as a good remedy for roses, use Magnesium sulphuricum (Epsom salt); repeat a few times during the growing season and before winter.

For young plants, use Magnesium carbonicum at the beginning of the growing season.

For combined Magnesium and Phosphorus deficiencies, use Magnesium phosphoricum.

SECTION 12: BORON DEFICIENCY

=====

For Boron deficiency, use Acidum boracicum.

For combined Calcium and Boron deficiencies, use Calcarea bromatum.

For combined Sodium and Boron deficiencies, use Natrum bromatum.

SECTION 13: ZINC DEFICIENCY

=====

For Zinc deficiency alone, use Zincum metallicum.

For combined Zinc and Phosphorus deficiencies, use Zincum phosphoricum.

For combined Zinc and Sulphur deficiencies, use Zincum sulphuricum.

When standard Zinc remedies fail, use Zincum aceticum.

SECTION 14: PLANT CONDITION / STAGE

=====

For germination, soak seeds for a few hours or overnight in Arnica montana 30, Thuja occidentalis 30, Phosphoric acid 30, and Spirulina 12.

For any injury or trauma, such as transplanting, use Arnica montana 30, Aconite 200, Calendula officinalis 30, and Staphisagria 200.

For pruning, use Arnica montana 30, Calendula officinalis 30, and Staphisagria 200.

To strengthen seedlings after transplantation, apply Arnica montana 30, Silicea 12, and Phosphorus 30 once per season.

For weak, puny plants, or plants that fail to thrive, use Arnica montana 30, Calendula officinalis 30, Silicea 12, Psorinum 30, China officinalis 30, and Selenium 12.

For flowering, use Phosphorus 30 and Arnica montana 30.

To increase pollination, use *Apis mellifica* 30.

To support the production of aromatic compounds in herbs, such as pepper, rosemary, and mint, use Sulphur 12.

For fruiting, spray *Calcarea phosphoricum* 12 and Borax 12 foliarly before flowering.

For post-harvest plant recovery, use *Arnica montana* 30, *China officinalis* 30, and *Calendula officinalis* 12.

For post-harvest root and fruit rots, use *Natrum muriaticum* 200.

As a compost accelerator, which supports healthy decomposition and composting and stimulates soil microbial activity, apply *Carbo vegetabilis* 30, *Bovista* 30, *Calcarea carbonicum* 30, *Calcarea nitrica* 12, and *Ignis alcoholis* 30 weekly.

SECTION 15: CROP-SPECIFIC REMEDIES

=====

The isopathic remedy for maize is *Stigmata Maydis-Zea* 12.

The isopathic remedy for potato is *Solanum tuberosum* 12.

The constitutional remedy for potato is *Vitis vinifera* 6.

The isopathic remedy for onion is *Allium cepa* 12.

The isopathic remedy for garlic is *Allium sativum* 12.

The isopathic remedy for beetroot is *Beta vulgaris* 12.

The isopathic remedy for strawberry is *Fragaria vesca* 12.

The isopathic remedy for spinach is *Spinacea oleracea* 12.

The constitutional remedy for spinach is *Urtica dioica* 12.

The isopathic remedy for carrot is *Daucus carota* 12.

The isopathic remedy for basil is *Ocimum basilicum* 12.

The isopathic remedy for mint is *Mentha piperita* 12.

The isopathic remedy for capsicum is *Capsicum annum* 12.

For capsicum plant growth and fruit size, use Sulphur 12.

The isopathic remedy for the Brassica family is *Brassica oleracea* 12.

The isopathic remedy for radish is *Raphanus sativus* 12.

The isopathic remedy for peas is *Pisum sativum* 12.

The isopathic remedy for coriander is *Coriandrum sativum* 12.

The isopathic remedy for beans, including kidney bean, is *Phaseolus nanus* 30.

The isopathic remedy for pigeon pea is *Cajanus indicus* 30.

The isopathic remedy for sugarcane is *Saccharum officinale* 12.

The isopathic remedy for pearl millet (bajra) is *Ceanothus americanus* 12.

The isopathic remedy for squash is *Cucurbita pepo* 12.

The isopathic remedy for turmeric is *Curcuma longa* 6.

The isopathic remedy for ginger is *Zingiber officinale* 6.

The isopathic remedy for lemon is *Citrus limonum* 12.

The isopathic remedy for banana is *Musa sapientum* 12.

The isopathic remedy for papaya is *Carica papaya* 12.

The isopathic remedy for mango is *Mangifera indica* 12.

The isopathic remedy for pomegranate is *Granatum* 30.

Stigmata Maydis-Zea 12 is used as a companion remedy to beans and squash.

Ricinus communis 12 is used as a companion remedy to grapevines.

The constitutional remedy for citrus plants is *Apium graveolens* 12.

For fruiting in lemon, use *Janosia ashoka* 12.

For flowering in guava and apricot, use *Staphysagria* 200.

The constitutional remedy for tomato is *Ocimum basilicum* 12 combined with *Vitis vinifera* 6.

The isopathic remedy for tomato is *Lycopersicum esculentum* 12.

For tomato plants, a three-day treatment improves harvest and taste, and prevents blossom-end rot, brown rot, and Greenback. First, water all tomato seedlings once with *Calendula officinalis* 30. One week later, apply the three-day treatment: on Day 1 give *Calcarea carbonicum* 6, on Day 2 give *Kalium phosphoricum* 6, and on Day 3 give *Magnesium phosphoricum* 6. After one week, apply the remedies again, and repeat once more after another week, for a total of three applications.

SECTION 16: FUNGAL DISEASE

=====

As a broad-spectrum remedy for fungal disease, use *Thuja occidentalis*, *Arnica montana*, Sulphur, Ozone, *Natrum sulphuricum*, *Arsenicum album*, and *Salicylicum acidum*.

For early blight in potato, use *Arsenicum album* 30, *Natrum muriaticum* 30, *Phaseolus nanus* 30, *Salix alba* 30, Sulphur 200, *Ferrum phosphoricum* 12, and *Arnica montana* 200.

For late blight in potato, use *Arsenicum album* 30, *Thuja occidentalis* 30, *Phaseolus nanus* 30, *Salix alba* 30, *Chamomilla* 30, *Silicea* 200, and *Ferrum phosphoricum* 12.

For false smut in rice, use *Allium sativum* 12, *Azadirachta indica* 12, *Ustilago maydis* 30, *Ocimum basilicum* 12, and *Ferrum phosphoricum* 12.

For powdery mildew and downy mildew, use *Cuprum metallicum* 30, Sulphur 200, *Thuja occidentalis* 30, *Natrum sulphuricum* 30, *Silicea* 200, and *Acid aceticum* 30.

For fungus on capsicum, use *Calcarea carbonicum* 30.

For *Botrytis cinerea* (grey mould) on strawberry, use *Equisetum hyemale* 200 and Sulphur 200.

For black sigatoka, use *Arsenicum album* 30.

SECTION 17: VIRAL DISEASE

=====

As a broad-spectrum remedy for viral disease, use *Acidum ascorbicum*, *Gelsemium*, *Curcuma longa*, *Zingiber officinale*, *Salicylicum acidum*, *Natrum salicylicum*, and *Thuja occidentalis*.

For tomato leaf curl virus, use *Psorinum* 30, *Salicylicum acidum* 6, *Azadirachta indica* 12, *Acidum ascorbicum* 12, *Antimonium tartaricum* 1M, *Vitis vinifera* 6, and *Curcuma longa* 6.

For cucurbits leaf curl virus, use *Psorinum* 30, *Salicylicum acidum* 6, *Azadirachta indica* 12, *Acidum ascorbicum* 12, *Antimonium tartaricum* 1M, *Curcuma longa* 6, and *Zingiber officinale* 6.

For yellow mosaic virus, use *Acidum ascorbicum* 12, Hydrogen peroxide 12, *Salicylicum acidum* 6, *Azadirachta indica* 12, *Curcuma longa* 6, and *Zingiber officinale* 6.

For tobacco mosaic virus, use *Acidum ascorbicum* 12, *Arsenicum album* 6, *Betula Pendula* 12, *Thyroidinum* 6, *Salicylicum acidum* 6, and *Uranium nitricum* 6.

For potato virus, use *Salicylicum acidum* 6, *Natrum salicylicum* 6, *Acidum ascorbicum* 12, *Vitis vinifera* 6, *Curcuma longa* 6, and *Zingiber officinale* 6.

SECTION 18: BACTERIAL DISEASE

=====

For potato scab, use *Allium sativum* 12, *Origanum majorana* 6, *Zincum metallicum* 12, *Ustilago maydis* 30, *Argemone mexicana* 6, and *Citrus limonum* 6.

SECTION 19: PESTS

=====

As a broad-spectrum remedy for pests, use *Azadirachta indica*, *Cimicifuga racemosa*, *Ledum palustre*, *Sabadilla*, *Staphysagria*, *Latrodectus mactans*, and *Calcarea carbonicum*.

For chronic infestations, or for application at the end of the growing season to make plants resistant to insects next year, use Sulphur and Psorinum.

As a broad-spectrum remedy for caterpillars, use *Apis mellifica* 30, *Vespa crabro* 30, *Sambucus nigra* 30, Sulphur 200, and *Andrographis paniculata* 30 (Kalmegh).

For army worm, spray the maize whorls with *Piper nigrum*, *Sambucus nigra*, Cina, *Bombyx processiona*, Absinthium, Sulphur, and *Latrodectus mactans*.

For all crawling pests, and for caterpillars on cabbage, use Sulphur 200.

For scale, use *Mezereum* 200, *Cuprum metallicum* 30, *Antimonium tartaricum* 1M, *Radium bromatum* 1M, Petroleum 30, *Coccinella septempunctata* 12, and *Rhus toxicodendron* 200.

For termites, ants, and cutworms, use Camphora 30.

For aphids, use *Coccinella septempunctata*, *Staphysagria*, *Calcarea carbonicum*, *Cimicifuga racemosa*, and *Capsicum annuum*.

For leaf miner, use *Thuja occidentalis*, *Ledum palustre*, Absinthium, and *Allium sativum*.

For mites, use *Allium sativum*, *Allium cepa*, *Thuja occidentalis*, and *Bovista*.

For spider mites, use *Bovista* 30, *Petroleum* 30, *Psorinum* 200, *Sulphur* 200, and *Capsicum annuum* 30.

For brown plant hopper, use *Calcarea carbonicum* 6, *Belladonna* 12, and *Staphysagria* 200.

For potato leafhopper, use *Artemisia absinthium* 6, *Azadirachta indica* 12, and *Calcarea carbonicum* 6.

For grasshopper, use *Azadirachta indica*, *Tanacetum vulgare*, *Hypericum perforatum*, *Thuja occidentalis* 30, *Sulphur* 200, and *Latrodectus mactans* 12.

For whitefly, use *Antimonium tartaricum* 1M, *Psorinum* 200, and *Sulphur* 200.

For squash bug, use *Cantharis* 30.

For snails and slugs, use *Helix tosta* 12.

For pests with a carapace, use *Petroleum* 30.

For leafhopper, use *Andrographis paniculata* 30 (Kalmegh), *Allium sativum* 12, *Apium graveolens* 12, and *Foeniculum vulgare* 12.

For mealy bug, use *Antimonium tartaricum* 200, *Petroleum* 200, *Capsicum annuum* 30, *Acid aceticum* 30, and *Equisetum arvense* 30.

GUIDELINES FOR USING THE REPERTORY

=====

1. It is important to first ascertain the underlying cause of the symptoms observed in plants. For example, leaf curling may result from a nutrient deficiency, such as boron deficiency, a viral disease, or a combination of both. Since the appropriate homeopathic treatment depends on the underlying cause, accurate diagnosis is essential before selecting a remedy.
2. If a specific issue is listed, begin with the corresponding specific remedy. Should it fail to produce the desired results, follow it with the broad-spectrum remedy for that category.
3. For issues not specifically covered in this repertory, use the broad-spectrum remedy for the relevant category.
4. Constitutional remedy is most effective during the transition from vegetative growth to flowering and fruiting. It strengthens the plant's natural vitality and promotes fruit development.
5. Prophylactic Use: Homeopathic medicines can be applied prophylactically before the appearance of disease symptoms, particularly when a plant is at known risk of infection. Such applications are intended to stimulate the plant's natural defense mechanisms.
6. Remedies from different categories may be combined when multiple conditions are present. For instance, during a cold and damp winter, remedies from both the cold stress and dampness categories may be mixed and applied together.
7. If all the remedies listed for a particular problem are not available, it is best to proceed with spraying those that are available.

GENERAL PRECAUTIONS

=====

1. Although no standard interval has been established, it is good practice to maintain a gap of at least two to three days between the application of agro-homeopathic remedies and herbal concoctions.

2. Silicea, whether used alone or in combination with other remedies, should not be applied too frequently. Maintain a minimum interval of two weeks between successive applications.

3. As Silicea and Sulphur are antidotal to each other, a minimum interval of two weeks should be maintained between the application of Silicea and Sulphur, including any remedies or combinations containing either medicine.

4. Do not spray *Allium sativum* or *Allium cepa* on beans or peas.

POTENCY SELECTION

=====

1. Where the potency of a medicine is not specified in this repertory, follow the potency guidelines given below.

2. Potency for Soil: Use lower potencies when treating soil imbalances. Potencies such as 6X, 6C, and especially 12C have been found to be effective. Lower potencies contain more of the original substance and are generally considered better suited for treating the soil. Although some practitioners use tissue salts (typically 3X), slightly higher potencies such as 6C or 12C have often been reported to produce better results.

3. Potency for Insects and Pests: Higher potencies generally produce a more intense effect. In acute situations, such as rapidly developing insect infestations, 200C has been found to be particularly effective.

4. Potency for Diseases: Diseases can be slow or fast to develop. For persistent but slow-progressing diseases, use 30C. For rapidly spreading and severe diseases, use 200C.

5. If the specified potency is unavailable, use the closest available potency. The first rule of thumb when choosing potency is to use what you have. The main thing is giving the plant the remedy it needs.

APPLICATION FREQUENCY

=====

1. Aggressive situations: When the problem is fast-moving and devastating, apply the remedy more frequently. Treat acute symptoms for three consecutive days, then wait one week to assess the response. If the problem has improved but is not completely resolved, repeat the treatment for another three consecutive days, which is usually sufficient.

2. Prophylactic application: When a plant is at high risk of disease or pest infestation, the selected homeopathic medicine may be sprayed before symptoms appear, preferably at 7 day intervals during high-risk periods.

3. Soil deficiencies & fertilization: One treatment is usually enough for two weeks during the growing season.

4. Pests, insects & fast-moving diseases: In fast-moving, aggressive acute situations (pest and insect invasions and diseases such as blight), results should be visible within 2-4 days. If not, change the remedy. In slow-moving chronic situations, wait one week before assessing the response.

PREPARATION AND APPLICATION OF HOMEOPATHIC SPRAY

=====

1. Fill a clean bottle with approximately 200 mL of tap, spring, or rainwater.

2. Add 30 drops (or 7 globules) of the selected homeopathic remedy to the bottle. If using a combination of two or more remedies, add 15 drops of each remedy instead.

3. Wait until globules have dissolved. Close the bottle and succuss it 20 times.

4. Fill a clean spray tank with 15 L of water.

5. Pour the entire 200 mL of the succussed solution into the spray tank.
6. Stir the solution 15-20 times using a plastic or wooden spoon, stick or stirring rod. Avoid metal implements, as they may interfere with the action of the homeopathic remedy.
7. Stir in one direction (either clockwise or anticlockwise). The homeopathic spray is now ready for application.
8. The best time to spray homeopathic medicines is in the early morning or late evening. Avoid spraying during the afternoon, especially around midday, as high temperatures and intense sunlight cause the spray to dry quickly, reduce its effectiveness, and may increase the risk of leaf scorch.
9. Spray the affected parts of the plant, or apply the remedy as a foliar spray, and water the roots with the solution. Applying the remedy both as a foliar spray and to the roots enhances its overall effect. When treating the soil, water it with the remedy solution.
10. One 15 L spray tank is generally sufficient for about one-quarter acre. To treat one acre, prepare and apply approximately 60 L of spray.
11. Without alcohol as a preservative, the remedy solution will not remain effective for long and should ideally be used within a week. If it needs to be stored for a longer period, add a few tablespoons of alcohol to help preserve the solution.
12. Before and after each application, rinse the bottle, spray tank, watering can and all other equipment thoroughly with fresh water only. Do not use synthetic soaps or detergents. Make sure the spraying equipment is not contaminated with any smell or medicine of any kind. If in doubt, rinse well with the hottest water possible and leave it to dry for 24 hours before using it again.

STORING HOMEOPATHIC PREPARATIONS

=====

1. Don't store homeopathic remedies in conditions warmer than room temperature. Neither should they be kept in the fridge.

2. Don't expose them to direct sunlight.

3. Don't store them near strong-smelling substances (e.g. perfume, soap, spices, vinegar).

4. Avoid magnetic fields (radio, TV, mobile phone).

RECOMMENDED MANUFACTURERS

=====

The recommended manufacturers are SBL Pvt. Ltd.; Dr. Willmar Schwabe India Pvt. Ltd.; B. Jain Pharmaceuticals Pvt. Ltd.; Father Muller Homoeopathic Pharmaceutical Division; Bakson's Homoeopathy; Dr. Reckeweg & Co.; and St. George's Homoeopathy (for rare remedies).