

rstanarm, brms, bambi, and formula syntax

- R (and systems before R) have used a formula syntax to describe models, e.g.

rstanarm, brms, bambi, and formula syntax

- R (and systems before R) have used a formula syntax to describe models, e.g.
 - linear model: $y \sim 1 + x$ or $y \sim x$

rstanarm, brms, bambi, and formula syntax

- R (and systems before R) have used a formula syntax to describe models, e.g.
 - linear model: $y \sim 1 + x$ or $y \sim x$
 - spline model: $y \sim s(x)$

rstanarm, brms, bambi, and formula syntax

- R (and systems before R) have used a formula syntax to describe models, e.g.
 - linear model: $y \sim 1 + x$ or $y \sim x$
 - spline model: $y \sim s(x)$
 - binomial model:
`deaths ~ dosage, family = binomial`

rstanarm, brms, bambi, and formula syntax

- R (and systems before R) have used a formula syntax to describe models, e.g.
 - linear model: $y \sim 1 + x$ or $y \sim x$
 - spline model: $y \sim s(x)$
 - binomial model:
`deaths ~ dosage, family = binomial`
 - hierarchical model:
`weight ~ age + (age | ratid)`

rstanarm, brms, bambi, and formula syntax

- R (and systems before R) have used a formula syntax to describe models, e.g.
 - linear model: $y \sim 1 + x$ or $y \sim x$
 - spline model: $y \sim s(x)$
 - binomial model:
`deaths ~ dosage, family = binomial`
 - hierarchical model:
`weight ~ age + (age | ratid)`
- `rstanarm` has pre-compiled Stan models
 - no need to wait compilation time
 - limited size of the pre-compiled binaries in CRAN limits the number of model, family, prior combinations

rstanarm, brms, bambi, and formula syntax

- R (and systems before R) have used a formula syntax to describe models, e.g.
 - linear model: $y \sim 1 + x$ or $y \sim x$
 - spline model: $y \sim s(x)$
 - binomial model:
`deaths ~ dosage, family = binomial`
 - hierarchical model:
`weight ~ age + (age | ratid)`
- `rstanarm` has pre-compiled Stan models
 - no need to wait compilation time
 - limited size of the pre-compiled binaries in CRAN limits the number of model, family, prior combinations
- `brms`
 - extended formula syntax, e.g.
heteroscedastic linear model: `bf(y ~ x, sigma ~ x)`
 - user defined families
 - requires compilation of the generated Stan code

rstanarm, brms, bambi, and formula syntax

- R (and systems before R) have used a formula syntax to describe models, e.g.
 - linear model: $y \sim 1 + x$ or $y \sim x$
 - spline model: $y \sim s(x)$
 - binomial model:
`deaths ~ dosage, family = binomial`
 - hierarchical model:
`weight ~ age + (age | ratid)`
- `rstanarm` has pre-compiled Stan models
 - no need to wait compilation time
 - limited size of the pre-compiled binaries in CRAN limits the number of model, family, prior combinations
- `brms`
 - extended formula syntax, e.g.
heteroscedastic linear model: `bf(y ~ x, sigma ~ x)`
 - user defined families
 - requires compilation of the generated Stan code
- `bambi` implements formula syntax in Python

Chapter 8: Modelling accounting for data collection

Highly recommended to read. Very informative, but also a dense chapter.

- We need to model the data collection unless it is ignorable
- We need to know when data collection is ignorable

Chapter 8: Modelling accounting for data collection

Highly recommended to read. Very informative, but also a dense chapter.

- We need to model the data collection unless it is ignorable
- We need to know when data collection is ignorable
- Data collection
 - Sample surveys
 - Designed experiments
 - Randomization
 - Observational studies
 - Censoring and truncation

Chapter 14: Introduction to regression models

- Justification of conditional modeling
 - if joint model factorizes $p(y, x|\theta, \phi) = p(y|x, \theta)p(x|\phi)$
we can model just $p(y|x, \theta)$

Chapter 14: Introduction to regression models

- Justification of conditional modeling
 - if joint model factorizes $p(y, x|\theta, \phi) = p(y|x, \theta)p(x|\phi)$
we can model just $p(y|x, \theta)$
- Gaussian linear model with conjugate prior
 - the conditional posterior is multivariate normal
 - with fixed prior on weights, the joint posterior is N-Inv- χ^2

Chapter 14: Introduction to regression models

- Justification of conditional modeling
 - if joint model factorizes $p(y, x|\theta, \phi) = p(y|x, \theta)p(x|\phi)$
we can model just $p(y|x, \theta)$
- Gaussian linear model with conjugate prior
 - the conditional posterior is multivariate normal
 - with fixed prior on weights, the joint posterior is N-Inv- χ^2
 - these properties are sometimes useful and thus good to know, but with probabilistic programming less often needed

Chapter 14: Introduction to regression models

- Justification of conditional modeling
 - if joint model factorizes $p(y, x|\theta, \phi) = p(y|x, \theta)p(x|\phi)$
we can model just $p(y|x, \theta)$
- Gaussian linear model with conjugate prior
 - the conditional posterior is multivariate normal
 - with fixed prior on weights, the joint posterior is N-Inv- χ^2
 - these properties are sometimes useful and thus good to know, but with probabilistic programming less often needed
- Bit on causal analysis (see much more in ROS Ch 18–21)

Chapter 14: Introduction to regression models

- Justification of conditional modeling
 - if joint model factorizes $p(y, x|\theta, \phi) = p(y|x, \theta)p(x|\phi)$ we can model just $p(y|x, \theta)$
- Gaussian linear model with conjugate prior
 - the conditional posterior is multivariate normal
 - with fixed prior on weights, the joint posterior is N-Inv- χ^2
 - these properties are sometimes useful and thus good to know, but with probabilistic programming less often needed
- Bit on causal analysis (see much more in ROS Ch 18–21)
- Assembling matrix of explanatory variables (see also ROS Ch 10,12)
 - identifiability, collinearity, nonlinear relations, indicator and categorical variables, interactions
 - variable selection is not much discussed (extra lecture)

Chapter 14: Introduction to regression models

- Justification of conditional modeling
 - if joint model factorizes $p(y, x|\theta, \phi) = p(y|x, \theta)p(x|\phi)$ we can model just $p(y|x, \theta)$
- Gaussian linear model with conjugate prior
 - the conditional posterior is multivariate normal
 - with fixed prior on weights, the joint posterior is N-Inv- χ^2
 - these properties are sometimes useful and thus good to know, but with probabilistic programming less often needed
- Bit on causal analysis (see much more in ROS Ch 18–21)
- Assembling matrix of explanatory variables (see also ROS Ch 10,12)
 - identifiability, collinearity, nonlinear relations, indicator and categorical variables, interactions
 - variable selection is not much discussed (extra lecture)
- Regularization (see also ROS Ch 12)
 - not much discussed (see extra lectures)

Chapter 14: Introduction to regression models

- Justification of conditional modeling
 - if joint model factorizes $p(y, x|\theta, \phi) = p(y|x, \theta)p(x|\phi)$ we can model just $p(y|x, \theta)$
- Gaussian linear model with conjugate prior
 - the conditional posterior is multivariate normal
 - with fixed prior on weights, the joint posterior is N-Inv- χ^2
 - these properties are sometimes useful and thus good to know, but with probabilistic programming less often needed
- Bit on causal analysis (see much more in ROS Ch 18–21)
- Assembling matrix of explanatory variables (see also ROS Ch 10,12)
 - identifiability, collinearity, nonlinear relations, indicator and categorical variables, interactions
 - variable selection is not much discussed (extra lecture)
- Regularization (see also ROS Ch 12)
 - not much discussed (see extra lectures)
- Unequal variances and correlations

Lasso and Bayesian lasso

- Lasso is penalized maximum likelihood linear regression, with L1 one penalty where the amount penalty is adapted

Lasso and Bayesian lasso

- Lasso is penalized maximum likelihood linear regression, with L1 one penalty where the amount penalty is adapted
 - penalized maximum likelihood finds the mode given the penalty parameter, and is almost the same as maximum a posteriori

Lasso and Bayesian lasso

- Lasso is penalized maximum likelihood linear regression, with L1 one penalty where the amount penalty is adapted
 - penalized maximum likelihood finds the mode given the penalty parameter, and is almost the same as maximum a posteriori
 - when the amount of penalty is increased, marginal modes of weak effects go to zero first

Lasso and Bayesian lasso

- Lasso is penalized maximum likelihood linear regression, with L1 one penalty where the amount penalty is adapted
 - penalized maximum likelihood finds the mode given the penalty parameter, and is almost the same as maximum a posteriori
 - when the amount of penalty is increased, marginal modes of weak effects go to zero first
 - when the amount of penalty is increased, also the relevant coefficients are shrunk towards zero

Lasso and Bayesian lasso

- Lasso is penalized maximum likelihood linear regression, with L1 one penalty where the amount penalty is adapted
 - penalized maximum likelihood finds the mode given the penalty parameter, and is almost the same as maximum a posteriori
 - when the amount of penalty is increased, marginal modes of weak effects go to zero first
 - when the amount of penalty is increased, also the relevant coefficients are shrunk towards zero
 - sometimes relaxed lasso is used, where after variable selection coefficients are re-estimated

Lasso and Bayesian lasso

- Lasso is penalized maximum likelihood linear regression, with L1 one penalty where the amount penalty is adapted
 - penalized maximum likelihood finds the mode given the penalty parameter, and is almost the same as maximum a posteriori
 - when the amount of penalty is increased, marginal modes of weak effects go to zero first
 - when the amount of penalty is increased, also the relevant coefficients are shrunk towards zero
 - sometimes relaxed lasso is used, where after variable selection coefficients are re-estimated
- Bayesian lasso uses Laplace distribution as prior
 - Laplace prior is equivalent to L1 penalty

Lasso and Bayesian lasso

- Lasso is penalized maximum likelihood linear regression, with L1 one penalty where the amount penalty is adapted
 - penalized maximum likelihood finds the mode given the penalty parameter, and is almost the same as maximum a posteriori
 - when the amount of penalty is increased, marginal modes of weak effects go to zero first
 - when the amount of penalty is increased, also the relevant coefficients are shrunk towards zero
 - sometimes relaxed lasso is used, where after variable selection coefficients are re-estimated
- Bayesian lasso uses Laplace distribution as prior
 - Laplace prior is equivalent to L1 penalty
 - but the Bayesian inference includes distribution for parameters and that distribution doesn't shrink to a point at zero, even if the mode would be at zero

Lasso and Bayesian lasso


- Lasso is penalized maximum likelihood linear regression, with L1 one penalty where the amount penalty is adapted
 - penalized maximum likelihood finds the mode given the penalty parameter, and is almost the same as maximum a posteriori
 - when the amount of penalty is increased, marginal modes of weak effects go to zero first
 - when the amount of penalty is increased, also the relevant coefficients are shrunk towards zero
 - sometimes relaxed lasso is used, where after variable selection coefficients are re-estimated
- Bayesian lasso uses Laplace distribution as prior
 - Laplace prior is equivalent to L1 penalty
 - but the Bayesian inference includes distribution for parameters and that distribution doesn't shrink to a point at zero, even if the mode would be at zero
 - empirically better results obtained with more sparse priors

Lasso and Bayesian lasso


- Lasso is penalized maximum likelihood linear regression, with L1 one penalty where the amount penalty is adapted
 - penalized maximum likelihood finds the mode given the penalty parameter, and is almost the same as maximum a posteriori
 - when the amount of penalty is increased, marginal modes of weak effects go to zero first
 - when the amount of penalty is increased, also the relevant coefficients are shrunk towards zero
 - sometimes relaxed lasso is used, where after variable selection coefficients are re-estimated
- Bayesian lasso uses Laplace distribution as prior
 - Laplace prior is equivalent to L1 penalty
 - but the Bayesian inference includes distribution for parameters and that distribution doesn't shrink to a point at zero, even if the mode would be at zero
 - empirically better results obtained with more sparse priors
 - it's best to separate selection of sensible prior, good posterior inference, and the decision analysis of which variables are important

Sparse priors


Laplacian




Student-t



Strawderman-Berger




Horseshoe



from Carvalho, Polson, Scott (2009).

Regularized horseshoe




- Piironen and Vehtari (2017). Sparsity information and regularization in the horseshoe and other shrinkage priors. In Electronic Journal of Statistics, 11(2):5018-5051. [Online](#)
- `rstanarm:prior=hs()`
- `brms:prior=horseshoe()`

Projpred selection vs. Lasso

See projpred in an extra lecture


Same simulated regression data as in lecture 9,3,
 $n = 50$, $p = 500$, $p_{\text{rel}} = 150$, $\rho = 0.5$



Projpred selection vs. Lasso

See projpred in an extra lecture


Same simulated regression data as in lecture 9,3,
 $n = 50$, $p = 500$, $p_{\text{rel}} = 150$, $\rho = 0.5$



Projpred selection vs. Lasso

See projpred in an extra lecture



Same simulated regression data as in lecture 9,3,
 $n = 50$, $p = 500$, $p_{\text{rel}} = 150$, $\rho = 0.5$



Projpred selection vs. Lasso

See projpred in an extra lecture

Same simulated regression data as in lecture 9,3,
 $n = 50$, $p = 500$, $p_{\text{rel}} = 150$, $\rho = 0.5$



Chapter 15: Hierarchical linear models

- Since you know hierarchical models, theory is easy
- With probabilistic programming computation is also easy
 - BDA3 discusses some other computational issues
 - section on transformations for HMC is relevant
(see also Stan user guide 21.7 Reparameterization)

Chapter 15: Hierarchical linear models

- Since you know hierarchical models, theory is easy
- With probabilistic programming computation is also easy
 - BDA3 discusses some other computational issues
 - section on transformations for HMC is relevant
(see also Stan user guide 21.7 Reparameterization)
- Fixed, random, and mixed effects models
 - we don't recommend using these terms, but they are so popular that it's useful to know them

Chapter 15: Hierarchical linear models

- Since you know hierarchical models, theory is easy
- With probabilistic programming computation is also easy
 - BDA3 discusses some other computational issues
 - section on transformations for HMC is relevant (see also Stan user guide 21.7 Reparameterization)
- Fixed, random, and mixed effects models
 - we don't recommend using these terms, but they are so popular that it's useful to know them

$y \sim 1 + x$	fixed / population effect; pooled model
$y \sim 1 + (0 + x \mid g)$	random / group effects
$y \sim 1 + x + (1 + x \mid g)$	mixed effects; hierarchical model

- ANOVA in section 15.6 (see also `stan_aov`)

Chapter 16: Generalized linear models

- Bioassay model is an example of GLM

Chapter 16: Generalized linear models

- Bioassay model is an example of GLM
- Components: (see also ROS Ch 13–15)
 1. The linear predictor $\eta = X\beta$

Chapter 16: Generalized linear models

- Bioassay model is an example of GLM
- Components: (see also ROS Ch 13–15)
 1. The linear predictor $\eta = X\beta$
 2. The link function $g(\cdot)$ and $\mu = g^{-1}(\eta)$

Chapter 16: Generalized linear models

- Bioassay model is an example of GLM
- Components: (see also ROS Ch 13–15)
 1. The linear predictor $\eta = X\beta$
 2. The link function $g(\cdot)$ and $\mu = g^{-1}(\eta)$
 3. Outcome distribution model with location parameter μ

Chapter 16: Generalized linear models

- Bioassay model is an example of GLM
- Components: (see also ROS Ch 13–15)
 1. The linear predictor $\eta = X\beta$
 2. The link function $g(\cdot)$ and $\mu = g^{-1}(\eta)$
 3. Outcome distribution model with location parameter μ
 - the distribution can also depend on dispersion parameter ϕ

Chapter 16: Generalized linear models

- Bioassay model is an example of GLM
- Components: (see also ROS Ch 13–15)
 1. The linear predictor $\eta = X\beta$
 2. The link function $g(\cdot)$ and $\mu = g^{-1}(\eta)$
 3. Outcome distribution model with location parameter μ
 - the distribution can also depend on dispersion parameter ϕ
 - originally just exponential family distributions (e.g. Poisson, binomial, negative-binomial), which all have natural location-dispersion parameterization

Chapter 16: Generalized linear models

- Bioassay model is an example of GLM
- Components: (see also ROS Ch 13–15)
 1. The linear predictor $\eta = X\beta$
 2. The link function $g(\cdot)$ and $\mu = g^{-1}(\eta)$
 3. Outcome distribution model with location parameter μ
 - the distribution can also depend on dispersion parameter ϕ
 - originally just exponential family distributions (e.g. Poisson, binomial, negative-binomial), which all have natural location-dispersion parameterization
 - after MCMC made computation easy, GLM can refer to models where outcome distribution is not part of exponential family and dispersion parameter may have its own latent linear predictor

Chapter 16: Generalized linear models

- Bioassay model is an example of GLM
- Components: (see also ROS Ch 13–15)
 1. The linear predictor $\eta = X\beta$
 2. The link function $g(\cdot)$ and $\mu = g^{-1}(\eta)$
 3. Outcome distribution model with location parameter μ
 - the distribution can also depend on dispersion parameter ϕ
 - originally just exponential family distributions (e.g. Poisson, binomial, negative-binomial), which all have natural location-dispersion parameterization
 - after MCMC made computation easy, GLM can refer to models where outcome distribution is not part of exponential family and dispersion parameter may have its own latent linear predictor

`deaths ~ dosage, family = binomial`

Chapter 16: Generalized linear models

- Bioassay model is an example of GLM
- Components: (see also ROS Ch 13–15)
 1. The linear predictor $\eta = X\beta$
 2. The link function $g(\cdot)$ and $\mu = g^{-1}(\eta)$
 3. Outcome distribution model with location parameter μ
 - the distribution can also depend on dispersion parameter ϕ
 - originally just exponential family distributions (e.g. Poisson, binomial, negative-binomial), which all have natural location-dispersion parameterization
 - after MCMC made computation easy, GLM can refer to models where outcome distribution is not part of exponential family and dispersion parameter may have its own latent linear predictor

`deaths ~ dosage, family = binomial`

- Hierarchical GLM natural extension

Chapter 16: Generalized linear models

- Bioassay model is an example of GLM
- Components: (see also ROS Ch 13–15)
 1. The linear predictor $\eta = X\beta$
 2. The link function $g(\cdot)$ and $\mu = g^{-1}(\eta)$
 3. Outcome distribution model with location parameter μ
 - the distribution can also depend on dispersion parameter ϕ
 - originally just exponential family distributions (e.g. Poisson, binomial, negative-binomial), which all have natural location-dispersion parameterization
 - after MCMC made computation easy, GLM can refer to models where outcome distribution is not part of exponential family and dispersion parameter may have its own latent linear predictor

`deaths ~ dosage, family = binomial`

- Hierarchical GLM natural extension
- 16.3 Weakly informative priors section is excellent although the recommendation on using Cauchy has changed (see <https://github.com/stan-dev/stan/wiki/Prior-Choice-Recommendations>)

Chapter 17: Models for robust inference

- For example (see also ROS Ch 15)

normal	→	t -distribution
Poisson	→	negative-binomial
binomial	→	beta-binomial
probit	→	logistic / robit

Chapter 17: Models for robust inference

- For example (see also ROS Ch 15)
 - normal \rightarrow t -distribution
 - Poisson \rightarrow negative-binomial
 - binomial \rightarrow beta-binomial
 - probit \rightarrow logistic / robit
- Computation with MCMC easy
 - posterior can be multimodal

Chapter 17: Models for robust inference

- For example (see also ROS Ch 15)
 - normal → t -distribution
 - Poisson → negative-binomial
 - binomial → beta-binomial
 - probit → logistic / robit
- Computation with MCMC easy
 - posterior can be multimodal
 - rstanarm doesn't have t -distribution for outcome, but brms has

Chapter 18: Models for missing data

- Extends the data collection modelling from Chapter 8
- Useful terms

Chapter 18: Models for missing data

- Extends the data collection modelling from Chapter 8
- Useful terms
 - Missing completely at random (MCAR)
missingness does not depend on missing values or other observed values (including covariates)

Chapter 18: Models for missing data

- Extends the data collection modelling from Chapter 8
- Useful terms
 - Missing completely at random (MCAR)
missingness does not depend on missing values or other observed values (including covariates)
 - Missing at random (MAR)
missingness does not depend on missing values but may depend on other observed values (including covariates)

Chapter 18: Models for missing data

- Extends the data collection modelling from Chapter 8
- Useful terms
 - Missing completely at random (MCAR)
missingness does not depend on missing values or other observed values (including covariates)
 - Missing at random (MAR)
missingness does not depend on missing values but may depend on other observed values (including covariates)
 - Missing not at random (MNAR)
missingness depends on missing values

Chapter 18: Models for missing data

- Extends the data collection modelling from Chapter 8
- Useful terms
 - Missing completely at random (MCAR)
missingness does not depend on missing values or other observed values (including covariates)
 - Missing at random (MAR)
missingness does not depend on missing values but may depend on other observed values (including covariates)
 - Missing not at random (MNAR)
missingness depends on missing values
- Multiple imputation
 1. make a model predicting missing data
 2. sample repeatedly from the missing data model to generate multiple imputed data sets
 3. make usual inference for each imputed data set
 4. combine results

Chapter 18: Models for missing data

- Extends the data collection modelling from Chapter 8
- Useful terms
 - Missing completely at random (MCAR)
missingness does not depend on missing values or other observed values (including covariates)
 - Missing at random (MAR)
missingness does not depend on missing values but may depend on other observed values (including covariates)
 - Missing not at random (MNAR)
missingness depends on missing values
- Multiple imputation
 1. make a model predicting missing data
 2. sample repeatedly from the missing data model to generate multiple imputed data sets
 3. make usual inference for each imputed data set
 4. combine results
 - `mice` package is ver flexible

Chapter 18: Models for missing data

- Extends the data collection modelling from Chapter 8
- Useful terms
 - Missing completely at random (MCAR)
missingness does not depend on missing values or other observed values (including covariates)
 - Missing at random (MAR)
missingness does not depend on missing values but may depend on other observed values (including covariates)
 - Missing not at random (MNAR)
missingness depends on missing values
- Multiple imputation
 1. make a model predicting missing data
 2. sample repeatedly from the missing data model to generate multiple imputed data sets
 3. make usual inference for each imputed data set
 4. combine results
 - `mice` package is ver flexible
- `brms` can handle some missing data

Chapter 21: Gaussian process models

- Gaussian process is
 - infinite dimensional extension of normal distribution
 - useful prior for non-linear functions
 - for any finite number of variables, the marginal is multivariate normal $f_1, \dots, f_n \sim N(\mu(x_1, \dots, x_n), K(x_1, \dots, x_n))$

Chapter 21: Gaussian process models

- Gaussian process is
 - infinite dimensional extension of normal distribution
 - useful prior for non-linear functions
 - for any finite number of variables, the marginal is multivariate normal $f_1, \dots, f_n \sim N(\mu(x_1, \dots, x_n), K(x_1, \dots, x_n))$
- Often a priori $\mu = 0$

Chapter 21: Gaussian process models


- Gaussian process is
 - infinite dimensional extension of normal distribution
 - useful prior for non-linear functions
 - for any finite number of variables, the marginal is multivariate normal $f_1, \dots, f_n \sim N(\mu(x_1, \dots, x_n), K(x_1, \dots, x_n))$
- Often a priori $\mu = 0$
- Prior for smooth non-linear functions, e.g. with
$$k(x, x') = \tau^2 \exp\left(-\frac{|x-x'|^2}{2l^2}\right)$$

Chapter 21: Gaussian process models

- Gaussian process is
 - infinite dimensional extension of normal distribution
 - useful prior for non-linear functions
 - for any finite number of variables, the marginal is multivariate normal $f_1, \dots, f_n \sim N(\mu(x_1, \dots, x_n), K(x_1, \dots, x_n))$
- Often a priori $\mu = 0$
- Prior for smooth non-linear functions, e.g. with


$$k(x, x') = \tau^2 \exp\left(-\frac{|x-x'|^2}{2l^2}\right)$$

$\tau=1/2, l=2$




x

$\tau=1/4, l=1/2$



x

$\tau=1/2, l=1/2$




x

Chapter 21: Gaussian process models


- Gaussian process is
 - infinite dimensional extension of normal distribution
 - useful prior for non-linear functions
 - for any finite number of variables, the marginal is multivariate normal $f_1, \dots, f_n \sim N(\mu(x_1, \dots, x_n), K(x_1, \dots, x_n))$
- Often a priori $\mu = 0$
- Prior for smooth non-linear functions, e.g. with

$$k(x, x') = \tau^2 \exp\left(-\frac{|x-x'|^2}{2l^2}\right)$$


$\tau=1/2, l=2$



$\tau=1/4, l=1/2$



$\tau=1/2, l=1/2$




Chapter 21: Gaussian process models

- Conditional on covariance function parameter the posterior is just multivariate normal
 - need to make inference for covariance function parameters given the marginal likelihood
 - the exact computation of the marginal likelihood scales $O(N^3)$


- Easy to make additive models

$$y_t(t) = f_1(t) + f_2(t) + f_3(t) + f_4(t) + f_5(t) + \epsilon_t$$



Chapter 21: Gaussian process models

- For non-Gaussian outcome models similar extension as GLMs
- Survival model example:



GPs in Stan

- GP specific software (e.g. GPy, GPflow, GPyTorch) scale computationally better for GPs than Stan
- Stan has some built-in covariance functions
- Hilbert space basis function approximation of GPs is fast for 1D-3D (Riutort-Mayol et al., 2022)
 - Birthday example
 - Motorcycle example
- In case of non-Gaussian outcome models, sampling of latent variables can be slow (Laplace integration over the latents coming)

GPs in Stan

- GP specific software (e.g. GPy, GPflow, GPyTorch) scale computationally better for GPs than Stan
- Stan has some built-in covariance functions
- Hilbert space basis function approximation of GPs is fast for 1D-3D (Riutort-Mayol et al., 2022)
 - Birthday example
 - Motorcycle example
- In case of non-Gaussian outcome models, sampling of latent variables can be slow (Laplace integration over the latents coming)
- brms:
 - covariance matrix based computation:
 $y \sim \text{gp}(x)$
 - Hilbert space basis function approximation:
 $y \sim \text{gp}(x, k=20)$

Regression and Other Stories

- Gelman, Hill, and Vehtari (2020). Regression and Other Stories.
 - uses Bayesian inference, but maths and computation is minimal
 - focuses on different models and how think about modeling
 - a lot of different examples
 - <https://avehtari.github.io/ROS-Examples/>

Bayesian workflow

Gelman, Vehtari, Simpson, Margossian, Carpenter, Yao, Kennedy, Gabry, Bürkner, and Modrák (2020). Bayesian workflow. [arXiv:2011.01808](https://arxiv.org/abs/2011.01808)

