

Introduction to Handling Data

ECON20222 - Lecture 1

Ralf Becker and Martyn Andrews

January 2019

What is this course unit about?

- Help you implement and interpret the main estimation and inference techniques used in Economics
- Focus on:
 - ▶ causal inference
 - ▶ the main pitfalls of time-series analysis

At the end of this unit ...

You will be able to:

- Do intermediate data work in R
- Confidently apply regression analysis in R
- Apply more advanced causal inference techniques in R
- Find coding help for any new challenges in R
- Identify inference appropriate for the model being estimated
- Discuss strengths and weaknesses of particular empirical applications
- Interpret empirical results (with due caution!)

What you need to do

To learn in this unit you need to:



coding, cleaning data, struggling,
self-learning, amazement at what
you can do

answering real questions, that there
is not always a clear answer

Assessment Structure and feedback

- Online test (on the use of R) - 10%
- End-of-Term exam (MC and short answer questions) - 50%
- Group coursework - 40% (see extra info)

Aim for today

Statistics/Econometrics

- Summary Statistics
- Difference between population and sample
- Hypothesis testing
- Graphical Data Representations
- Simple regression analysis

R Coding

- Introduce you to R and RStudio
- How do I learn R
- Import data into R
- Perform some basic data manipulation
- Perform hypothesis tests
- Estimate a regression

Why Data Matter



Average GDP growth rates (High Income countries - 1946 to 2009):

Debt/GDP category	(0,30]	(30,60]	(60,90]	(90,Inf]
Avg Growth Rate (RR)	4.09%	2.87%	3.40%	-0.02%

Why Data Matter

- Reinhard and Rogoff seem to suggest that there is a level of debt (debt/GDP > 90%) beyond which higher debt levels will significantly reduce growth. **Caution: Debt <> Deficit**
- While they often provided caveats in their arguments, their results were referred to when austerity policies were justified.

For example George Osborne:

here on Channel 4: (<https://www.channel4.com/news/george-osborne-defends-austerity-plan>)

here in a Conference Speech:

(<https://conservative-speeches.sayit.mysociety.org/speech/601526>)

Average GDP growth rates (High Income countries - 1946 to 2009, RR - Reinhard and Rogoff, HAP - Herndon, Ash and Pollin):

Debt/GDP Category	(0,30]	(30,60]	(60,90]	(90,Inf]
Avg Growth Rate (RR)	4.09%	2.87%	3.40%	-0.02%
Avg Growth Rate (HAP)	4.17%	3.12%	3.22%	2.17%

Why Data Matter

Some summaries are available here

The New Yorker	The Reinhart and Rogoff Controversy: Summing Up
The Economist	The 90% question
Financial Times	Interviews with Carmen Reinhard [1 , 2 , 3 , 4]

Important issues that arise from this

- Which way is the causality? Debt to Growth or Growth to Debt? Reinhard and Rogoff are suitably careful to not associate any direct causality from the summary statistics.
- But in the political discourse such "subtleties" often get lost
- Would different summary statistics have changed the narrative?

The Plan for today

- Replicate the above summary statistics in R
- Why one can get two very different results based on the same data
- Use this example to
 - ▶ introduce you to R
 - ▶ review some summary statistics
 - ▶ review simple regression and its implementation
 - ▶ introduce some basic visualisations

Introduce R/R-Studio



- R is a statistical software package, it is open source and free
- a lot of useful functionality is added by independent researchers via packages (also for free)



- RStudio is a user interface which makes working with R easier. You need to install R before you [install RStudio](#).



- [ECLR](#) is a web-resource we have set up to support you in your R work.

Welcome to RStudio

The screenshot displays the RStudio interface with the following components:

- Source Editor:** Contains R code for data manipulation using `group_by`, `summarize`, and `print`. The code calculates mean and median growth rates for different debt categories.
- Environment:** Lists loaded packages and data objects, including `active_staff`, `admin`, `phdus`, `mscdi`, `units`, `temp`, and `units`.
- Files:** Shows the file explorer with a list of files and folders, including `ggleep.txt`, `CarmenReihart.jpg`, `Data_Intro.pdf`, `Embracoe.jpg`, `Hendson_Ash_Pollin_WP_2013.pdf`, `KennethRogoff.jpg`, `Lecture_1_-_Data_Introduction.Rmd`, `McGregor_MISchool_Depression_2013-04-18.pdf`, `RStudio.xslx`, `Lecture_1_-_Data_Introduction.pdf`, `Rimage.jpeg`, `RStudio_image.png`, `ECIR.jpg`, and `RStudio_screen.JPG`.

R Code Snippet:

```
R[1]
R[2]
R[3]
R[4]
R[5]
R[6]
R[7]
R[8]
R[9]
R[10]
R[11]
R[12]
R[13]
R[14]
R[15]
R[16]
R[17]
R[18]
R[19]
R[20]
R[21]
R[22]
R[23]
R[24]
R[25]
R[26]
R[27]
R[28]
R[29]
R[30]
R[31]
R[32]
R[33]
R[34]
R[35]
R[36]
R[37]
R[38]
R[39]
R[40]
R[41]
R[42]
R[43]
R[44]
R[45]
R[46]
R[47]
R[48]
R[49]
R[50]
R[51]
R[52]
R[53]
R[54]
R[55]
R[56]
R[57]
R[58]
R[59]
R[60]
R[61]
R[62]
R[63]
R[64]
R[65]
R[66]
R[67]
R[68]
R[69]
R[70]
R[71]
R[72]
R[73]
R[74]
R[75]
R[76]
R[77]
R[78]
R[79]
R[80]
R[81]
R[82]
R[83]
R[84]
R[85]
R[86]
R[87]
R[88]
R[89]
R[90]
R[91]
R[92]
R[93]
R[94]
R[95]
R[96]
R[97]
R[98]
R[99]
R[100]
```

Data Table:

debtcat	n	mean	median
(0,30]	13	4.08921971	4.001282
(30,60]	15	2.86594921	2.889572
(60,90]	14	3.39943999	2.857815
(90,inf]	7	-0.02421961	1.028900

Console Output:

```
4351
4352
4353
4354
4355
4356
4357
4358
4359
4360
4361
4362
4363
4364
4365
4366
4367
4368
4369
4370
4371
4372
4373
4374
4375
4376
4377
4378
4379
4380
4381
4382
4383
4384
4385
4386
4387
4388
4389
4390
4391
4392
4393
4394
4395
4396
4397
4398
4399
4400
4401
4402
4403
4404
4405
4406
4407
4408
4409
4410
4411
4412
4413
4414
4415
4416
4417
4418
4419
4420
4421
4422
4423
4424
4425
4426
4427
4428
4429
4430
4431
4432
4433
4434
4435
4436
4437
4438
4439
4440
4441
4442
4443
4444
4445
4446
4447
4448
4449
4450
4451
4452
4453
4454
4455
4456
4457
4458
4459
4460
4461
4462
4463
4464
4465
4466
4467
4468
4469
4470
4471
4472
4473
4474
4475
4476
4477
4478
4479
4480
4481
4482
4483
4484
4485
4486
4487
4488
4489
4490
4491
4492
4493
4494
4495
4496
4497
4498
4499
4500
```

Write Code Files or the Basic Workflow

- keep an original data file (usually `‘.xlsx’` or `‘.csv’`) and do not overwrite this file
- any manipulation we make to the data (data cleaning, statistical analysis etc.) is command based and we collect all these commands in a script file. R will then interpret and execute these commands. It is hence like a recipe which you present to a chef. These script files have extension `‘.r’`
- you can also learn to write Rmarkdown files (`‘.rmd’`). They combine code with normal text and output.
- When you write code you should ensure that you add comments to your code. Comments are bit of text which is ignored by R (everything after an `‘#’`) but helps you or someone else to decipher what the code does.

By following the above examples you make it easy for yourself and others to replicate your work. At the core of the Reinhard/Rogoff controversy was the ability of independent researchers to replicate their work!

Prepare your code

We start by uploading the extra packages we need in our code.

The first time you need these packages at a computer you may need to install these. Use the following code to do this

```
install.packages(c("readxl", "tidyverse", "ggplot2", "stargazer"))
```

This only needs to be done once on a particular computer. However, every time you want to use any of these packages in a code you need to make them available to your code (load them):

```
library(tidyverse)    # for almost all data handling tasks  
library(readxl)       # to import Excel data  
library(ggplot2)      # to produce nice graphs  
library(stargazer)    # to produce nice results tables
```

The data

Then we load the data from excel

```
RRData <- read_excel("RRdata.xlsx")  
RRData <- as.data.frame(RRData) # forces data.frame structure
```

and check some characteristics of the data which are now stored in RRData:

```
str(RRData) # prints some basic info on variables
```

```
## 'data.frame':    1171 obs. of  4 variables:  
## $ Year      : num  1946 1947 1948 1949 1950 ...  
## $ Country: chr  "Australia" "Australia" "Australia" "Austri  
## $ debtgdp: num  190 177 149 126 110 ...  
## $ dRGDP   : num  -3.56 2.46 6.44 6.61 6.92 ...
```

Discuss data.frame, number of obs and number of variables, their names and variable types

Variable types

These are four basic data types.

- character: `"a", "swc"`
- numeric: `2, 15.5`
- integer: `2L` (the `L` tells `R` to store this as an integer)
- logical: `TRUE, FALSE`

It is important that you know and understand differences between data types. Each variable has a particular type and some operations only work for particular datatypes. For instance, we need `num` or `int` for any mathematical operations.

In our `data.frame` three variables are `num` and one is of `chr` type.

We will encounter `logical` variables frequently. **they are very powerful**

factor variables

It is necessary to change categorical variables (here Country) to factor variables.

```
RRData$Country <- as.factor(RRData$Country)
str(RRData)
```

```
## 'data.frame':    1171 obs. of  4 variables:
## $ Year      : num  1946 1947 1948 1949 1950 ...
## $ Country: Factor w/ 20 levels "Australia","Austria",...: 1
## $ debtgdp: num  190 177 149 126 110 ...
## $ dRGDP   : num  -3.56 2.46 6.44 6.61 6.92 ...
```

- `RRData$Country` calls variable `Country` in dataframe `RRData`
- `<-` assigns what is on the right `as.factor(RRData$Country)` to the variable on the left `RRData$Country`
- `as.factor(RRData$Country)` calls a function `as.factor` and applies it to `RRData$Country`

factor variables

factor variables are variables with discrete categories. Which ones they are you can find out with the `levels()` function:

```
levels(RRData$Country)
```

```
## [1] "Australia" "Austria" "Belgium" "Canada"
## [6] "Finland" "France" "Germany" "Greece"
## [11] "Italy" "Japan" "Netherlands" "New Zealand"
## [16] "Portugal" "Spain" "Sweden" "UK"
```

Learn more about your data


Use the `stargazer` function for some initial summary stats for `num` or `int` variables

```
stargazer(RRData, type = "text")
```

```
##
## =====
## Statistic      N      Mean      St. Dev.      Min      Pctl(25) Pctl(75)      Max
## -----
## Year           1,171 1,979.155    17.944      1,946      1,964      1,995      2,009
## debtgdp         1,171  46.283     32.395       3.279      22.148     61.474    247.482
## dRGDP           1,171   3.430      2.981     -10.942      1.911      5.100     27.329
## -----
```

Scatter plot of the data

```
p1 <- ggplot(RRData, aes(debtgdp, dRGDP)) +  
  geom_point(size=0.5) +      # this produces the scatter plot  
  geom_smooth(method = "lm", se = FALSE) # adds the line  
p1
```



Point out that each dot represents one country/year's data, e.g. France in 1991. Point out line of best fit

Regression Line

The line in the previous plot is the line of best fit coming from a linear regression

$$rGDP = \alpha + \beta debtgdp + u \quad (\text{Population Model})$$

- The population model is defined by unknown parameters α and β and the unknown error terms u . We will use sample data to obtain sample estimates of these parameters.
- The error terms u contain the effects of any omitted variables and reflect that any modelled relationship will only be an approximation. The u are random variables

$$rGDP_{it} = \hat{\alpha} + \hat{\beta} debtgdp_{it} + \hat{u}_{it} \quad (\text{Estimated Sample Model})$$

Here we have two subscripts as the data have a cross-section (**i**) and a time-series dimension (**t**).

The regression line in the previous figure is represented by

$$\widehat{rGDP}_{it} = \hat{\alpha} + \hat{\beta} debtgdp_{it} \quad (\text{Regression Line})$$

Ordinary Least Squares - Simple Regression

Regression analysis is the core technique used in Econometrics. It is based on certain assumptions about the *Population Model* and the error terms u_{it} (more on this in the next few weeks).

How to estimate parameters (get $\hat{\alpha}$ and $\hat{\beta}$) using the available sample of data?

Ordinary Least Squares and the Simple Regression Model

```
mod1 <- lm(dRGDP~debtgdp, data= RRData)
summary(mod1)
```

```
##
## Call:
## lm(formula = dRGDP ~ debtgdp, data = RRData)
##
## Residuals:
##      Min       1Q   Median       3Q      Max
## -12.9958  -1.5200  -0.0774   1.5707  23.6960
##
## Coefficients:
##              Estimate Std. Error t value Pr(>|t|)
## (Intercept)  4.279290   0.148970   28.73  < 2e-16 ***
## debtgdp      -0.018355   0.002637   -6.96 5.67e-12 ***
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
## Residual standard error: 2.922 on 1169 degrees of freedom
```

OLS - nice output

```
stargazer(mod1,type="text")
```

```
##
## =====
##                               Dependent variable:
##                               -----
##                               dRGDP
## -----
## debtgdp                      -0.018***
##                               (0.003)
##
## Constant                     4.279***
##                               (0.149)
## -----
## Observations                  1,171
## R2                           0.040
## Adjusted R2                   0.039
## Residual Std. Error          2.922 (df = 1169)
## F Statistic                   48.439*** (df = 1; 1169)
## =====
```


OLS - calculation and interpretation

How were $\hat{\beta}$ and $\hat{\alpha}$ calculated?

$$\begin{aligned}\hat{\beta} &= \frac{\widehat{Cov}(dRGDP_{it}, debtgdp_{it})}{\widehat{Var}(debtgdp_{it})} \\ \hat{\alpha} &= \overline{dRGDP}_{it} - \hat{\beta} * \overline{debtgdp}_{it}\end{aligned}$$

How to interpret $\hat{\beta} = -0.018$?

An increase of one unit in **debtgdp** (=1% point) is related to a decrease of GDP growth of 0.018 units (=0.018% points - pp)

Have we established that higher debt levels **cause** lower GDP growth?

NO

OLS - estimator properties

We need to recognise that parameter estimator is a random variable.

Let's use $y_{it} = \alpha + \beta x_{it} + u_{it}$ for ease of notation:

$$\begin{aligned}\hat{\beta} &= \frac{Cov(y_{it}, x_{it})}{Var(x_{it})} \\&= \frac{Cov(\alpha + \beta x_{it} + u_{it}, x_{it})}{Var(x_{it})} \\&= \frac{Cov(\alpha, x_{it}) + Cov(\beta x_{it}, x_{it}) + Cov(u_{it}, x_{it})}{Var(x_{it})} \\[10pt]\hat{\beta} &= \frac{Cov(\alpha, x_{it})}{Var(x_{it})} + \beta \frac{Cov(x_{it}, x_{it})}{Var(x_{it})} + \frac{Cov(u_{it}, x_{it})}{Var(x_{it})} \\&= 0 + \beta \frac{Var(x_{it})}{Var(x_{it})} + \frac{Cov(u_{it}, x_{it})}{Var(x_{it})} \\&= \beta + \frac{Cov(u_{it}, x_{it})}{Var(x_{it})}\end{aligned}$$

OLS - estimator properties

What can we learn from this?

- If u_{it} is a random variable, so is $\hat{\beta}$
- Any particular value we get is a draw from a random distribution
- An estimator is unbiased if, on average, the estimates would be equal to the unknown β
at this stage the concept of unbiasedness may still be a little hazy and that is fine
- For this to happen we need to assume that $Cov(u_{it}, x_{it}) = 0$ as then $E(\hat{\beta}) = \beta$

Why do we need to assume this? Because while we do have values for x_{it} we do not have values for the unobserved error terms u_{it} . Hence we cannot test this. As you will find out this is mainly a thinking exercise and one at the core of much of what we do.

OLS - the exogeneity assumption

For $\hat{\beta}$ in $y_{it} = \alpha + \beta x_{it} + u_{it}$ to be unbiased (i.e. on average correct) we needed

$$Cov(u_{it}, x_{it}) = 0$$

This is sometimes called the **Exogeneity assumption**. The error term has to be uncorrelated to the explanatory variable x_{it}

There are a lot of reasons why this assumption may be breached.

- Simultaneity ($debtgdp \rightarrow dRGDP$ and $dRGDP \rightarrow debtgdp$)
Discuss the fact that we have to assume that causality here goes in both directions. Hence we cannot attach one one-directional causal interpretation to the estimated coefficient. If you can estimate the model the other way round
- Measurement error in x_{it} (not dealt with here)
- Omitted relevant variables or unobserved heterogeneity

Excursion - Omitted variable bias

Let's imagine I would get all your first year statistics final grades (gr_i) and how often you attended lectures during the semester (att_i). (Note, no time series dimension here. All obs from same year hence no time subscript t .)

$$gr_i = \alpha + \beta att_i + u_i$$


What value would you expect $\hat{\beta}$ to take?

positive, more lecture attendance better grade

Can we say that lecture attendances causes higher grades? What do we think is hidden in the error term? Intelligence, Attitude, Aptitude for quants. How are they related to att_i ? Presumably some positively, e.g. attitude to studies. But all of these variables should really be in the model, right? So some of their effect is going to be captured by att_i and this may bias the estimate $\hat{\beta}$ up as it now captures, say, the effect of an increase in attendance but also in attitude.

Reinhard/Rogoff Example - debtgdp


```
tempdata <- RRData %>% filter(Country %in% c("Germany", "Greece", "UK", "US"))  
ggplot(tempdata, aes(Year, debtgdp, color=Country)) +  
  geom_line(size=1)      # this produces the line plot
```



Point out the piping operator and the filter function and then ggplot and geom_line


Reinhard/Rogoff Example - dRGDP

```
tempdata <- RRData %>% filter(Country %in% c("Germany", "Greece", "UK", "US"))  
ggplot(tempdata, aes(Year, dRGDP, color=Country)) +  
  geom_line(size=1)      # this produces the line plot
```



Reinhard/Rogoff Example - dRGDP v debtgdp

```
tempdata <- RRData %>% filter(Country %in% c("Germany", "Greece", "UK", "US"))  
ggplot(tempdata, aes(debtgdp, dRGDP, color=Country)) +  
  geom_point()      # this produces the scatter plot
```



Reinhard/Rogoff Example

What have we learned?

- dept data are persistent
- growth data as well but less so
- data for different countries have different characteristics

Group the data into debt categories

Let's create the four debt categories: (0,30], (30,60], (60,90], (90,Inf].

```
RRData <- RRData %>% mutate(dgcat = cut(RRData$debtgdp, breaks=c(0,30,60,90,Inf)))
```

```
RRData %>% group_by(dgcat) %>%  
  summarise_at("dRGDP", funs(mean, median)) %>%  
  print()
```

```
## # A tibble: 4 x 3  
##   dgcat      mean median  
##   <fct>    <dbl>  <dbl>  
## 1 (0,30]    4.17    4.15  
## 2 (30,60]   3.12    3.11  
## 3 (60,90]   3.22    2.9  
## 4 (90,Inf]  2.17    2.34
```

Print out the mutate function in combination with the cut function and the pipe, then the group_by and summarise_at function

These are the statistics we saw before (not the one reported by Reinhard and Rogoff)

Why did Reinhard and Rogoff report different results?

Debt Category	(0,30]	(30,60]	(60,90]	(90,Inf]
Avg Growth Rate (RR)	4.09%	2.87%	3.40%	-0.02%
Avg Growth Rate (HAP)	4.17%	3.12%	3.22%	2.17%

Why did RR get so much lower growth for the highest debt category, (90,Inf]?

Thomas Herndon, Michael Ash and Robert Pollin (2014) replicated the work. They identified the following differences to the above analysis.

- 1 They excluded early-postwar data for New Zealand, Australia and Canada, arguing that these data are atypical for later periods, essentially they are outliers
- 2 A spreadsheet error resulted in data for the five countries (Australia, Austria, Belgium, Canada and Denmark) to not be included.
- 3 Observations are not weighted equally.

Replicate Reinhard and Rogoff's results

```
## Selective treatment of early years
RRselective <- RRData %>%
  filter(!((Year<1950 & Country=="New Zealand") |
            (Year<1951 & Country=="Australia") |
            (Year<1951 & Country=="Canada") ))

## Spreadsheet error omitting five countries
RRselective <- RRselective %>%
  filter(!( Country %in%
            c("Australia", "Austria", "Belgium", "Canada", "Denmark") ))

RRselective %>% group_by(dgcat) %>%
  summarise_at("dRGDP", funs(mean, median)) %>%
  print()
```

```
## # A tibble: 4 x 3
##   dgcat      mean median
##   <fct>    <dbl>  <dbl>
## 1 (0,30]    4.24    4.4
## 2 (30,60]   2.98    3.06
## 3 (60,90]   3.16    2.85
## 4 (90,Inf]  1.69    2.33
```

Replicate Reinhard and Rogoff's results

So the first two differences explain some but not all of the differences.
Let's implement the different weighting.

```
RRselective2 <- RRselective %>%  
  group_by(dgcat, Country) %>%           # create category and country groups  
  summarize( m1 = mean(dRGDP, na.rm = TRUE)) %>% # calculate cat average  
  group_by(dgcat) %>%                   # group again by category  
  summarize( n = n(), mean = mean(m1, na.rm = TRUE),  
             median = median(m1, na.rm = TRUE)) %>%  
  print()
```

```
## # A tibble: 4 x 4  
##   dgcat      n    mean median  
##   <fct>   <int>   <dbl>   <dbl>  
## 1 (0,30]    13   4.09     4.00  
## 2 (30,60]   15   2.87     2.89  
## 3 (60,90]   14   3.40     2.86  
## 4 (90,Inf]    7 -0.0242    1.03
```

Don't worry too much about the full details of this weighting scheme.

The combination of the three changes make a massive difference

Why data (and their treatment) matter

The combination of these changes made a significant difference to the summary statistics.

Remember, the data were very persistent.

Reinhard and Rogoff

For each country all years which fall into one of the four categories are averaged and then treated as one observation

Herndon, Ash and Pollin

Each country/year observation is treated as one independent observation.

Perhaps it is right to not treat each observation as a new piece of information. But the RR weighting scheme seems to discard a lot of information.

These results made a significant difference in the political discourse.

Outlook

Over the next weeks you will learn

- to perform more advanced statistical analysis in R, such as:
 - ▶ Hypothesis testing
 - ▶ Multivariate regression analysis
 - ▶ specification testing
- to devise methods to draw causal inference
- to understand the main pitfalls of time-series modelling and forecasting