

Livestock Sector Forecasting in Kenya: A Cohort-Based Quarterly Analysis on Population, Production, and Feed Demand

Natasha Wavomba^{1*}, Maria Ogamba¹, Lorraine Eyinda¹,
Joseph Gitonga¹, John Olukuru¹, Joseph Sevilla¹

^{1*}@iLabAfrica, Strathmore University, Nairobi, Kenya.

Abstract

Kenya's livestock sector contributes 42% of agricultural GDP and 12% of national GDP, yet quantitative planning remains constrained by the absence of reproducible, data-driven analytical models. This study presents a cohort-based quarterly forecasting framework spanning nine value chains: dairy cattle, beef cattle, camels, goats, sheep, poultry, pigs, rabbits, and apiculture, initialized from the 2019 KNBS livestock census with projections from 2020 to 2041. The model applies a quarterly transition sequence covering mortality, culling, offtake, age-based maturation, and reproduction across all cohorts. Production modules translate herd and flock dynamics into species-specific output metrics, while a carrying capacity constraint — derived from satellite-based vegetation indices and adjusted by a utilization factor — regulates growth against forage availability. A feed demand module estimates dry matter requirements by species and agro-ecological zone. Projected outputs indicate a CAGR of 3.6-6% across all value chains. Beyond forecasting, the framework exposes productivity bottlenecks and structural inefficiencies, including feed deficits, low offtake rates, and suboptimal reproductive performance, that aggregate assessments typically obscure. The model provides an analytical basis for food security planning, targeted infrastructure investment, and evidence-based livestock policy.

Keywords: Livestock model forecasting, carrying capacity, biological parameters

1 Introduction

In most African countries, the livestock sector supports the livelihoods of millions of rural households and plays an important role in poverty reduction. In livestock-rich countries such as Kenya, the sector contributes substantially to the national economy while providing income, employment, and food security across both rural and urban populations. Beyond primary production, livestock also supports employment throughout value chains, including dairy, meat, and leather processing industries. The livestock sector plays a major role in Kenya’s food system, contributing approximately 12% of the country’s Gross Domestic Product (GDP), 40% of agricultural GDP, and 22% of the food system GDP [1].

Kenya’s population is projected to reach approximately 96 million by 2050, with nearly half of the population residing in urban areas. Consequently, demand for animal-source foods is expected to increase substantially owing to population growth, urbanization, rising incomes, and changing dietary preferences [1]. Meeting this growing demand will require sustained improvements in livestock productivity, efficient management of feed and water resources, and evidence-based planning to support the sustainable development of the livestock sector. Reliable forecasting frameworks that project livestock populations, production outputs, feed demand, and resource requirements are therefore essential for evidence-based planning, strategic investment, and sustainable livestock sector development.

Statistical time-series models are commonly used to forecast livestock populations using historical data. These methods estimate future livestock numbers based on past population trends and have been applied to support livestock planning and management. [1] used autoregressive and probabilistic time-series models to forecast pig populations, while Ozbek compared autoregressive integrated moving average (ARIMA) and artificial neural network (ANN) models for forecasting cattle populations in Türkiye. Similarly, [1] applied exponential smoothing to forecast beef cattle populations in Indonesia. Although these methods have demonstrated good predictive performance, they rely primarily on historical trends and do not explicitly represent biological processes such as reproduction, mortality, maturation, culling, and offtake. Consequently, their ability to evaluate the effects of biological parameters and management interventions on future livestock population dynamics is limited.

In contrast to statistical forecasting approaches, biological simulation models explicitly represent the demographic processes governing livestock populations. Animals are grouped into cohorts according to age, sex, or production stage, with transitions driven by biological processes such as reproduction, maturation, mortality, culling, and replacement. These models offer a biologically realistic representation of herd dynamics and enable the evaluation of alternative management strategies. [1] developed an age- and sex-structured cattle population model that incorporates management practices and animal movements to simulate herd dynamics. Such models serve as valuable decision-support tools by linking biological processes to herd structure and productivity, they are generally developed for specific livestock species or production systems and do not explicitly integrate livestock population dynamics, production, environmental carrying capacity, and feed demand within a unified forecasting framework. Consequently, there remains a lack of reproducible analytical

frameworks that simultaneously integrate livestock population dynamics, production, environmental carrying capacity, and feed demand to support long-term livestock sector planning.

To address these limitations, this study develops a cohort-based quarterly forecasting framework for Kenya’s nine principal livestock value chains. The framework explicitly models biological cohort transitions through reproduction, mortality, maturation, culling, and offtake while linking livestock population dynamics to species-specific production, environmental carrying capacity, and feed demand across agro-ecological zones. By integrating these components within a single modelling framework, the proposed approach provides a comprehensive decision-support tool for forecasting livestock populations, production, and resource requirements, thereby supporting evidence-based livestock planning and policy development in Kenya.

2 Literature Review

2.1 Significance of the Livestock Sector in Kenya

Livestock production plays a key role in Kenya’s economy and in household livelihoods, providing income, employment, wealth accumulation, and food. About 80% of Kenya’s land area is classified as arid and semi-arid land (ASAL), where pastoralism is the primary source of income and carries about 75% of the national herd. Livestock production contributes an estimated 42% of agricultural GDP and around 12% of Kenya’s total GDP [2].

National projections show that Kenya’s population will continue to grow, reaching an estimated 85 to 95 million by 2050, with close to half living in urban areas [3]. This demographic and urban transition is expected to drive a substantial rise in demand for animal-source foods, creating new opportunities across the different livestock value chains [4].

2.2 Historical trends in livestock population and production

Livestock production historical trends also suggest a close association between changes in output and changes in livestock numbers, but not necessarily in proportional terms. At continental level, the African Union’s Livestock Development Strategy for Africa states that increases in herd and flock size have contributed significantly to the growth in the supply of beef, milk, poultry and sheep and goat meat [5]. This finding suggests that population dynamics should be understood in the future assessment of livestock supply and that the growth of livestock populations may add to total production without being reflected in equivalent productivity increases.

Evidence from Kenya’s smallholder dairy sector further illustrates the relationship between livestock population structure and production. Wanjala and Njehia [6] noted that poor performance of dairy herd was associated with inadequate replacement stock, limited access to improved breeding animals, inadequate feed quantity and quality, breed characteristics and short lactation periods. The results are dairy cattle specific, but they do suggest a substantial consideration for livestock population and production analysis in general.

Previous studies imply that forecasts based purely on aggregate historic growth rates may overlook changes that occur within livestock populations over time. A more disaggregated approach can therefore be a basis for studying how changes in livestock numbers translate into production and then into livestock feed requirements. This provides a basis for cohort-based forecasting of livestock populations, production, and feed demand across Kenya’s livestock value chains.

2.3 Modelling Approaches for Livestock Population and Production Forecasting

Statistical modeling approaches have been used to explore factors that drive livestock productivity in Kenya. For instance, Ajak et al. [7] assessed dairy cattle productivity on smallholder farms in relation to factors such as feeding, breed, and disease, attributing poor performance largely to feed shortages alongside management- and price-related constraints. More recently, Were et al. [8] used multilevel mixed-effects linear regression to quantify the animal- and feeding-level factors associated with daily milk yield in smallholder herds in Kiambu County, identifying specific feeding practices and cow-level characteristics as significant predictors of output. Although such studies are primarily explanatory rather than predictive, they identify the production drivers that can be incorporated as explanatory variables in forecasting models.

2.4 Cohort-Based Herd Dynamics, Population and Production Projection Models

Empirical applications have highlighted the significance of explicitly considering the population structure of livestock. For example, Lesnoff et al. [9] used a demographic model to study the recovery of the cattle population after drought and showed that the recovery was particularly sensitive to reproductive performance. Likewise, Tuffa and Treydte [10] developed a stochastic age and sex-structured model for Boran cattle which included demographic processes, carrying capacity, market conditions and herder decisions. Their results suggested that decisions to remove animals from different demographic groups can drastically change population trajectories. Combined, these studies indicate that livestock population projections are more informative when they include the demographic processes by which populations are replenished, mature and removed, rather than just using aggregate growth rates.

2.5 Biological Transition Parameters: Mortality, Reproduction, Maturation and Off-take

Studies from across Sub-Saharan Africa have documented significant differences in mortality, fertility, maturation and off-take rates across livestock species and production environments. In a systematic review of livestock production parameters, Otte and Chilonda [11] summarized the relative high mortality of young cattle and small ruminants, low reproductive performance and generally low off-take in traditional production systems. Their synthesis also showed that population growth rates vary widely among species and production systems, emphasizing the necessity of biologically and

geographically relevant transition parameters. Off-take is important especially as animals may leave the herd by sale or slaughter before the end of their biological life. These processes imply that the observed changes in livestock population are the net result of births, maturation, mortality and removals.

2.6 Carrying Capacity, Feed Availability and Feed Demand

Availability of feed resources also limits the growth of the livestock population. Studies of carrying capacity in African rangelands show that livestock numbers must be matched against available forage, and remote-sensing studies have increasingly made it possible to include seasonal and spatial variation in forage availability in livestock assessments [12]. Conventional livestock feed balance methods calculate how dry matter animals need by looking at their body weight or using livestock unit equivalents. Common intake rates like the 2.5 percent of body weight per day standard have been helpful for figuring out how much feed animals need when doing livestock assessments. Applications across Sub-Saharan Africa have used these approaches to identify substantial and often seasonal feed deficits. These developments suggest that livestock population and production projections can provide a useful basis for estimating future feed requirements.

2.7 Research Gap

The reviewed literature establishes the building blocks of the present study on solid methodological ground. Historical evidence links changes in livestock numbers to changes in output [5, 6]; statistical and regression-based studies identify feeding, and cow-level factors that drive productivity [7, 8]; demographic and cohort-based models show that population projections are far more informative when they represent how animals are born, mature, and are removed than when they rely on aggregate growth rates [9, 10]; syntheses of Sub-Saharan production parameters document the mortality, fertility, maturation, and off-take rates needed to drive such models [11]; and rangeland studies show that herd size must be matched to forage availability, which remote sensing now allows to be estimated across seasons and space [12].

What remains scarce is a single, reproducible, biologically grounded framework that unifies these strands for Kenya—one that simultaneously (i) spans all nine key livestock value chains rather than a single species, (ii) advances cohort transitions at a quarterly time step to capture seasonality, and (iii) closes the loop from population to species-specific production to disaggregated feed demand. By integrating these elements into one coherent, parameter-transparent architecture, the present study addresses a clear and consequential gap, offering not only defensible forecasts but a diagnostic instrument for identifying the feed deficiencies, low off-take rates, and reproductive bottlenecks that aggregate assessments obscure.

3 Methodology

3.1 Overview

Eight value chains (both ruminant and non-ruminant animals) were selected on the basis of their contribution to Kenya’s agricultural GDP.¹ For each chain, forecasting covers three domains: **population dynamics**, **production volumes**, and **feed demand**. The baseline year is 2019; the projection horizon runs from 2020 to 2041, with 2020–2025 retained for model validation against observed values.

A first-principles modelling approach was adopted. This choice reflects (i) the transparency and auditability it affords where each intermediate quantity is directly traceable to its inputs, and (ii) the heterogeneous, multi-source nature of the available data, which precludes the large homogeneous datasets required by machine-learning methods.

Parameters were compiled from peer-reviewed literature, government publications, and sector reports, then validated by livestock experts from the Ministry, associations, and farmers. Accepted values are listed in Appendix 1.²

The model operates on a quarterly time step. For each quarter, the three domains are computed in sequence.

3.2 Population Model Equations

Population is tracked across a defined cohort structure specific to each value chain. Within each quarter, the following processes are applied in order where relevant: mortality, culling, offtake, transition to the next age cohort, and births. The starting population for each cohort is updated by sequentially applying the corresponding rates, with inflows from younger cohorts and new births added at the end of the period.

3.2.1 Beef Cattle

$$P(t+1) = [P(t)(1 - m_q)(1 - k_q)(1 - o_q)] + G(t) - E(t) + B(t) \quad (1)$$

$P(t)$	Starting pop.	m_q	Mortality rate (qtr.)
$P(t+1)$	Ending pop.	k_q	Culling rate
$G(t)$	Inflow (young cohort)	o_q	Offtake rate (steers)
$E(t)$	Outflow to next cohort	$B(t)$	New births (calves)

3.2.2 Goats

$$P(t+1) = [P(t)(1 - m_q)(1 - k_q)] + G(t) - E(t) + B(t) \quad (2)$$

Variable definitions follow the same convention as beef cattle; the offtake term o_q does not apply.

3.2.3 Camels

$$P(t+1) = P(t) - D(t) - C(t) - O(t) + B(t) \quad (3)$$

$D(t)$	Natural mortality	$C(t)$	Culling offtake
$O(t)$	Commercial offtake	$B(t)$	Births in period t

¹Citation required, get from Natasha

²Summarize parameters

3.2.4 Poultry

Three sub-sectors are modelled separately. T_{bro} denotes annual broiler throughput and S_r the rearing survival rate.

$$P_{\text{ind}}(t+1) = P_{\text{ind}}(t) - E_{\text{ind}}(t) + R_{\text{ind}}(t) + A(t) \quad (4)$$

$$P_{\text{lay}}(t+1) = P_{\text{lay}}(t) - E_{\text{lay}}(t) + \text{DOC}(t) \times S_r \quad (5)$$

$$T_{\text{bro}}(t+1) = T_{\text{bro}}(t) + A(t) \quad (6)$$

The sector addition term $A(t)$ and its sigmoid growth modifier $g(t)$ are:

$$A(t) = P(t) (r_{\text{ent}} + r_{\text{exp}}) g(t) \quad (7)$$

$$g(t) = g_{\text{min}} + (g_{\text{max}} - g_{\text{min}}) \exp\left(-\kappa^2 \frac{(t - t_0)^2}{2}\right) \quad (8)$$

3.2.5 Apiculture

$$H(t+1) = H(t) (1 - \delta(t))(1 - \rho(t)) + S(t) \quad (9)$$

$\delta(t)$	Seasonal colony loss	$\rho(t)$	Queen replacement rate
$S(t)$	New colonies (swarming)		

3.2.6 Pigs

$$P(t+1) = P(t) - D(t) - C(t) - O(t) + B(t) + R(t) \quad (10)$$

where $R(t)$ represents replacement additions to the breeding herd. Variable definitions for $D(t)$, $C(t)$, $O(t)$, $B(t)$ follow the camel convention.

3.2.7 Rabbits

Rabbits follow the same population equation as pigs (including $R(t)$).

3.3 Production Model Equations

Production volumes are derived from the population state at the end of each quarter. Each value chain has a distinct production equation reflecting the commodity it yields, that is meat, milk, eggs, honey, or wax, as detailed in Sections II-B through II-C. ³ For ruminants, production is computed separately for each cohort and then aggregated. For poultry, the three sub-sectors are modelled independently. Apiculture production is calculated at the hive level and then scaled by the number of hives.

3.3.1 Beef Cattle

$$\text{Beef (kg)} = \sum_c N_c \times \text{LW}_c \times \text{DP}_c \quad (11)$$

where $c \in \{\text{steers, cull cows, cull bulls}\}$, LW is live weight (kg) and DP is dressing percentage.

³Update the section names

3.3.2 Goats

$$\text{Meat (kg)} = (N_{\text{removed}} + N_{\text{excess bucks}}) \times \text{LW} \times \text{DP} \quad (12)$$

$$\text{Milk (kg)} = N_{\text{does}} \times f_L \times Y_d \times D_L \quad (13)$$

N_{does}	Surviving does
f_L, Y_d, D_L	Lactating fraction; daily yield (kg); lactation days
$N_{\text{excess bucks}}$	Bucks above 4% herd cap, removed for slaughter

3.3.3 Camels

$$\text{Milk (kg yr}^{-1}\text{)} = N_{\text{adult females}} \times f_L \times Y_d \times 365 \quad (14)$$

$$\text{Meat (kg)} = \sum_c N_c \times \text{LW}_c \times \text{DP}_c \quad (15)$$

where $c \in \{\text{juv. offtake, adult offtake, cull adults}\}$.

3.3.4 Poultry

$$\text{Eggs}_{\text{ind}}(t) = P(t) \times f_{\text{fem}} \times Y_e \quad (16)$$

$$\text{Eggs}_{\text{lay}}(t) = P(t) \times \eta_{\text{HD}} \times 365 \quad (17)$$

$$\text{Meat (kg)} = T_{\text{bro}} \times \text{LW} \times \text{DP} \quad (18)$$

where f_{fem} is female share, Y_e is eggs per hen per year, and η_{HD} is hen-day production rate.

3.3.5 Apiculture

$$\text{Honey}_{h,q} = H_h \cdot \frac{Y_h}{4} \cdot \bar{s} \cdot (1 - \varphi) \cdot \Omega(t) \cdot \varepsilon_h \quad (19)$$

$$\text{Wax}_{h,q} = \text{Honey}_{h,q} \times \omega_h \quad (20)$$

$$\text{Nectar}_{h,q} = H_h \cdot \frac{N_h}{4} \cdot \bar{s} \cdot \phi_q \cdot \nu_q \cdot \Omega(t) \quad (21)$$

$Y_h, \bar{s}$	Annual yield/hive (kg); effective colony strength
φ	Forage competition penalty (0–0.20)
ε_h	Hive-type efficiency factor
$\Omega(t)$	Technology/training improvement (sigmoid)
ω_h	Wax-to-honey ratio
ϕ_q, ν_q	Quarterly forage and nectar seasonality factors

3.3.6 Pigs and Rabbits

$$\text{Meat} = N_{\text{slaughtered}} \times \text{LW} \times \text{DP} \quad (22)$$

3.4 Feed Model Equations

Feed demand is estimated using dry matter intake (DMI) for all value chains. A DMI-based approach was adopted due to limited availability of detailed ration composition data across most value chains; it provides a consistent and tractable basis for estimating feed requirements from population and live weight alone. The specific formulations vary by value chain and are presented in Section II-D.

3.4.1 Beef Cattle, Pigs, and Rabbits

All three value chains use the same daily dry matter intake (DMI) formulation:

$$\text{DMI} = P(t) \times \text{LW} \times r_{\text{DMI}} \times 365 \quad (23)$$

where r_{DMI} is daily dry matter intake as a proportion of body weight.

3.4.2 Goats

Feed is summed across cohorts (kids, weaners, growers, does, bucks):

$$\text{Feed}(t) = \sum_{\text{cohort}} N_c \times \text{BW}_c \times r_{\text{DMI},c} \times D \quad (24)$$

where D is the number of days in the period.

3.4.3 Camels

Feed is accumulated quarterly, with intake weighted by average live weight over the quarter:

$$\text{Annual feed} = \sum_{q=1}^4 \text{Feed}_q \quad (25)$$

$$\text{Feed}_q = \sum_{\text{cohort}} N_c \times \left(\frac{\text{BW}_{0,c} + \text{ADG}_c \times 90}{2} \right) \times r_{\text{DMI},c} \times 90 \quad (26)$$

where ADG is average daily gain (kg d^{-1}).

3.4.4 Poultry

Each sub-sector has a distinct feed equation reflecting its production cycle structure:

$$F_{\text{ind}}(t) = [R(t)(f_{\text{chk}} d_{\text{chk}} + f_{\text{gr}} d_{\text{gr}}) + P(t) f_{\text{ad}} 365] (1 - \sigma)^t \quad (27)$$

$$F_{\text{lay}}(t) = \text{DOC}(t)(f_{\text{chk}} d_{\text{chk}} + f_{\text{pul}} d_{\text{pul}}) + P(t) f_{\text{hen}} 365 \quad (28)$$

$$F_{\text{bro}} = N_{\text{birds}} \sum_{\text{phase}} f_{\text{phase}} \times d_{\text{phase}} \times C_{\text{yr}} \quad (29)$$

where $f_{(\cdot)}$ is phase-specific intake (kg d^{-1}), $d_{(\cdot)}$ is phase duration (days), σ is the scavenging offset, and C_{yr} is production cycles per year.

4 Results and Discussion

Population and production are reported over 2019–2041. The 2019 values are historical observations; 2020–2041 are model projections. The 2019 baseline is therefore shown shaded in the figures, with the 2019–2020 segment drawn dotted, and where the transition from historical to projected data produces a step (Section ??) growth rates are quoted from 2020. Feed demand carries no 2019 historical value and is reported over 2020–2041 throughout.

4.1 Population

4.1.1 Ruminants

Total ruminant numbers rise from 88.0 million head in 2019 to 272.1 million in 2041, a 3.1-fold increase equivalent to 5.3 % per year (Table 1, Figure 1). Growth is not evenly distributed across value chains. Beef cattle expand fastest at 6.6 % per year, quadrupling to 67.8 million head and overtaking sheep as the second-largest ruminant population in 2036. Goats remain the largest single population throughout, growing 5.9 % per year to 121.3 million. Camels and dairy cattle follow intermediate trajectories of 5.2 % and 4.9 % per year respectively, each roughly tripling. Sheep are the slowest-growing ruminant chain at 3.3 % per year, doubling over the period; their share of the ruminant herd falls from 31.2 % in 2019 to 20.5 % in 2041.

Table 1 Ruminant population, 2019–2041 (thousand head)

Series (000 head)	2019	2025	2030	2035	2041	CAGR 2019–41 (%)
Goats	34 658	50 854	66 733	87 571	121 333	5.9
Beef cattle	16 640	23 168	32 365	45 282	67 767	6.6
Sheep	27 441	33 099	39 221	46 098	55 665	3.3
Camels	4 722	6 661	8 535	10 794	14 265	5.2
Dairy cattle	4 537	5 929	7 593	9 726	13 091	4.9
Total ruminants	87 998	119 710	154 448	199 470	272 119	5.3

4.1.2 Non-ruminants

Non-ruminant numbers grow far more slowly, from 106.3 million head in 2019 to 159.1 million in 2041, a 1.5-fold increase at 1.8 % per year (Table 2, Figure 2). The aggregate conceals a divergence between the commercial and non-commercial segments of the poultry chain. Layers and broilers expand by only about 10 % over 22 years (0.4 % per year) and plateauing after the early 2030s, whereas indigenous poultry grow steadily




Fig. 1 Projected ruminant population by value chain, 2019–2041. Logarithmic vertical axis; the shaded band and dotted segment mark the 2019 baseline, which is a historical value.

at 3.0 % per year to 89.3 million birds. The composition of the poultry flock shifts accordingly: indigenous birds rise from 44.1 % to 58.1 % of total poultry, while the combined layer and broiler share of all non-ruminants falls from 55.2 % to 40.5 %. The smallest chains grow fastest: rabbits 4.4-fold at 7.0 % per year and pigs 3.5-fold at 5.9 % per year. Both start below one million head, so they remain marginal in absolute terms. Managed bee colonies increase 2.7-fold to 3.5 million.

Table 2 Non-ruminant population, 2019–2041 (thousands). Bee colonies are reported separately and excluded from the total.

Series (000)	2019	2025	2030	2035	2041	CAGR 2019–41 (%)
Indigenous poultry	46 293	54 774	65 960	75 970	89 288	3.0
Broilers	45 838	47 477	49 975	50 175	50 386	0.4
Layers	12 844	13 114	13 946	14 030	14 089	0.4
Rabbits	737	1 149	1 585	2 191	3 235	7.0
Pigs	596	817	1 097	1 471	2 088	5.9
Total (excl. colonies)	106 309	117 331	132 563	143 837	159 086	1.8
Bee colonies	1 289	1 838	2 384	2 926	3 489	4.6




Fig. 2 Projected non-ruminant population by value chain, 2019–2041. Upper panel: livestock numbers (logarithmic axis). Lower panel: managed bee colonies (linear axis). Shaded bands and dotted segments mark the 2019 baseline, which is a historical value.

4.2 Production

4.2.1 Ruminants

Ruminant output expands across every product line (Table 3, Figure 3). Ten of the reported series carry a step between the 2019 historical value and the 2020 projection (Section ??); for those series growth is described here from 2020, and the 2019–2041 CAGRs in Table 3 should be read with that caveat.

Total milk from cattle, camels, and goats rises from 5.36 billion kg in 2020 to 15.41 billion kg in 2041, 5.2% per year. Cattle milk remains the largest single product at 9.26 billion kg in 2041; camel milk is the second largest at 5.61 billion kg, and since the two grow at almost identical rates from 2020 (5.0% and 5.3% per year) their ratio is close to stable, narrowing only from 1.75:1 to 1.65:1. Goat milk is the fastest-growing dairy product at 6.6% per year but remains a lot less at 531 million kg.

Meat output is led by chevon, which overtakes beef in 2020 and reaches 1.71 billion kg by 2041 against 1.37 billion kg of beef; both grow at 5.3–5.5% per year from 2020. Camel meat is the fastest-growing ruminant meat at 6.8% per year, though at 235 million kg in 2041 it remains well behind the two leading products. Mutton grows slowest of all ruminant products at 1.2% per year from 2020, reaching 165 million

kg, so the sheep chain contributes a declining share of ruminant meat. Wool, the only product in the third group, reaches 3.9 million kg at 2.8 % per year.

Table 3 Ruminant production, 2019–2041 (million kg)

Series (million kg)	2019	2025	2030	2035	2041	CAGR 2019–41 (%)
Cattle milk	3 087	4 205	5 375	6 883	9 263	5.1
Camel milk	985	2 672	3 369	4 248	5 611	8.2
Goat milk	156	222	292	383	531	5.7
Chevon	53.9	718	942	1 236	1 712	17.0
Beef	322	479	656	919	1 375	6.8
Camel meat	57.6	104	139	178	235	6.6
Mutton	49.4	100	117	137	165	5.6
Wool	0.84	2.40	2.80	3.26	3.92	7.3

Ruminant production, 2019–2041

Projected output by product group; all series in kilograms. Shaded/dotted: 2019 baseline from historical values




Fig. 3 Projected ruminant production, 2019–2041, by product group: milk, meat, and wool. Milk and meat panels use logarithmic vertical axes; the wool panel is linear. Shaded bands and dotted segments mark the 2019 baseline, which is a historical value.

4.2.2 Non-ruminants

Non-ruminant production growth is more subdued and closely tracks the population trajectories (Table 4, Figure 4). Poultry meat, the largest non-ruminant product, grows only 1.3 % per year to 109.7 million kg, consistent with the plateau in commercial bird numbers. Pork grows at 4.0 % per year to 34.1 million kg and rabbit meat at 4.6 % per year (from 2020) to 4.2 million kg. The apiculture products grow fastest: honey at 5.1 % per year to 41.6 million kg and beeswax at 5.4 % per year (from 2020) to 6.5 million kg. Honey output exceeds pork output in absolute mass from 2021 onward and is 1.2 times pork by 2041, making apiculture the second-largest non-ruminant product group by mass. Egg production, reported in trays and therefore shown in its own panel on a linear axis, rises from 180.4 million trays in 2020 to 228.7 million in

2041 (1.1 % per year); the 501.3 million trays recorded for 2019 is inconsistent with the remainder of the series.

Table 4 Non-ruminant production, 2019–2041 (million kg; eggs in million trays)

Series (million)	2019	2025	2030	2035	2041	CAGR 2019–41 (%)
Poultry meat	82.4	88.8	97.6	103	110	1.3
Pork	14.4	18.2	22.2	27.0	34.1	4.0
Rabbit meat	1.76	2.03	2.55	3.19	4.18	4.0
Honey	13.9	22.4	29.2	35.4	41.6	5.1
Beeswax	1.37	3.39	4.48	5.50	6.54	7.4
Eggs (million trays)	501	189	208	217	229	-3.5

Non-ruminant production, 2019–2041

Projected output by product group; eggs reported in trays. Shaded/dotted: 2019 baseline from historical values




Fig. 4 Projected non-ruminant production, 2019–2041, by product group: meat, eggs, and apiculture products. Meat and other panels use logarithmic vertical axes; the egg panel is linear because eggs are reported in trays. Shaded bands and dotted segments mark the 2019 baseline, which is a historical value.


4.3 Feed demand

4.3.1 Ruminants

Ruminant chains account for the great majority of projected feed demand (Table 5, Figure 5). Total requirement rises from 105.8 million tonnes DM in 2020 to 353.8 million tonnes in 2041, 5.9 % per year. Beef cattle alone account for 58.8 million tonnes in 2020 and 218.3 million in 2041, more than the other four ruminant chains combined in every year of the period. Their share rises from 55.5 % to 61.7 % as they grow at 6.4 % per year, the fastest of the five chains. Goats follow at 6.1 % per year to 36.2 million tonnes and camels at 5.6 % to 42.2 million tonnes, the latter overtaking dairy cattle in 2023; dairy cattle reach 41.2 million tonnes at 4.9 % per year. Sheep demand grows slowest, from 8.0 to 15.9 million tonnes at 3.3 % per year.

Table 5 Ruminant feed demand, 2020–2041 (thousand tonnes DM)

Series (000 t DM)	2020	2025	2030	2035	2041	CAGR 2020–41 (%)
Beef cattle	58 791	74 615	104 258	145 888	218 333	6.4
Camels	13 474	19 399	25 167	31 904	42 184	5.6
Dairy cattle	15 064	18 646	23 877	30 584	41 165	4.9
Goats	10 528	15 187	19 929	26 151	36 233	6.1
Sheep	7 989	9 332	11 135	13 133	15 889	3.3
Total ruminants	105 846	137 177	184 366	247 660	353 805	5.9


**Fig. 5** Projected ruminant dry-matter feed demand by value chain, 2020–2041. Logarithmic vertical axis. (Section ??).

4.3.2 Non-ruminants

Non-ruminant feed demand is less, rising from 3.29 million tonnes in 2020 to 6.46 million tonnes in 2041 at 3.3% per year (Table 6, Figure 6). That is 1.8% of the ruminant total in 2041. Poultry accounts for the largest share throughout but grows slowest at 1.9% per year to 2.83 million tonnes, so its share falls from 57.7% to 43.8%. Apiculture nectar demand grows at 5.0% per year to 2.49 million tonnes, reaching 88% of poultry feed demand by 2041. Pig feed demand reaches 1.09 million tonnes (4.1% per year) and rabbit demand 52.7 thousand tonnes (4.5% per year).

Table 6 Non-ruminant feed demand, 2020–2041 (thousand tonnes)

Series (000 t)	2020	2025	2030	2035	2041	CAGR 2020–41 (%)
Poultry	1 900	2 067	2 350	2 556	2 828	1.9
Apiculture (nectar)	900	1 259	1 706	2 093	2 489	5.0
Pigs	473	579	707	862	1 092	4.1
Rabbits	20.9	25.8	32.2	40.3	52.7	4.5
Total non-ruminants	3 293	3 931	4 795	5 552	6 462	3.3

**Fig. 6** Projected non-ruminant feed demand by value chain, 2020–2041. Apiculture is shown as nectar demand. Logarithmic vertical axis.

5 Managerial, Policy, and Societal Implications

Beyond its forecasting function, the cohort-based framework developed in this study operates as a diagnostic instrument that translates biological herd dynamics into decision-driven intelligence. By resolving each value chain into age and sex-structured cohorts advanced on a quarterly basis, its outputs speak simultaneously to herd and enterprise managers, to national and county policymakers, and to the broader societal objectives of food security, livelihood resilience, and environmental sustainability.

5.1 Managerial Implications

At the enterprise level, the framework identifies precisely where within a production system growth is being lost by isolating the cohorts and transition processes that

constrain performance—e.g., elevated calf, lamb, and kid mortality, depressed reproductive rates, and sub-optimal off-take structures. This granularity matters because, as seen from comparable East African rangeland systems indicates, the sale and retention rates of breeding and mature females exert the greatest influence on long-run population trajectories [10, 13].

Managers may therefore use model outputs to assess herd structure, inform culling and destocking decisions, and breeding interventions. The quarterly time step also facilitates monitoring seasonal changes in herd dynamics and feed requirements, potentially supporting early intervention where projected feed deficits or changes in herd condition indicate emerging risks [12].

5.2 Policy Implications

For national and county governments the framework directly translates into the analytical needs of national and county planning providing the value chain specific projections necessary to prioritize investment and set performance targets. The ability to run coherent forward scenarios across nine value chains provides a basis for evidence-based investment sequencing and could be institutionalized within national statistical and livestock agencies, transforming sector planning from static census snapshots to continuous data-driven foresight [?].

5.3 Societal Implications

At the societal scale, the framework addresses the demand-driven transformation of animal-source-food consumption that will accompany Kenya’s population growth and urbanization. Reliable projections of milk, meat, egg, and honey supply against rising demand are central to food and nutrition security planning, allowing anticipatory action on emerging supply gaps rather than reactive crisis response. Since livestock underpins the livelihoods of several million pastoralists and agro-pastoralists and constitutes the dominant source of income across the arid and semi-arid lands, the model’s capacity to anticipate feed shortfalls and productivity shocks has direct implications for poverty reduction, resilience, and the protection of vulnerable rural households [? ?].

6 Limitations and Opportunities for Future Research

Write limitations here.

7 Conclusion

Write conclusion here.

Acknowledgements. Write acknowledgements here.

Declarations

- Funding: Not applicable

- Conflict of interest/Competing interests: Not applicable
- Ethics approval and consent to participate: Not applicable
- Consent for publication: Not applicable
- Data availability: Not applicable
- Materials availability: Not applicable
- Code availability: Not applicable
- Author contribution: Not applicable

Appendix A Model Parameters

References

- [1] Bahta, S.T., Wanyoike, F.N., Kirui, L., Mensah, C., Enahoro, D.K., Karugia, J.T., Baltenweck, I.: Livestock sector transformation in Kenya: Current state and projections for the future (2023) https://doi.org/10.2499/9780896294561_0310568/138984
- [2] Nyariki, D.M., Amwata, D.A.: The value of pastoralism in kenya: Application of total economic value approach. *Pastoralism: Research, Policy and Practice* **9**(1), 9 (2019) <https://doi.org/10.1186/s13570-019-0144-x>
- [3] Kenya National Bureau of Statistics: 2019 kenya population and housing census, volume IV: Distribution of population by socio-economic characteristics. Technical report, Kenya National Bureau of Statistics, Nairobi, Kenya (2019). <https://www.knbs.or.ke/>
- [4] FAO, IFAD, WFP, UNICEF, WHO: The state of food security and nutrition in the world 2019: Safeguarding against economic slowdowns and downturns. Technical report, Food and Agriculture Organization of the United Nations, Rome (2019). <https://openknowledge.fao.org/server/api/core/bitstreams/16480532-17e9-4b61-b388-1d6d86414470/content>
- [5] African Union: The livestock development strategy for africa (LiDeSA) 2015–2035. Technical report, African Union – Interafrican Bureau for Animal Resources (AU-IBAR), Nairobi, Kenya (2015)
- [6] Wanjala, S.P.O., Njehia, B.K.: Herd characteristics on smallholder dairy farms in western kenya. *Journal of Animal Science Advances* **4**(8), 996–1003 (2014) <https://doi.org/10.5455/jasa.20140827111904>
- [7] Ajak, P.A.D., K., G.C., Wanyoike, M.M.M.: Evaluation of dairy cattle productivity in smallholder farms in nyeri county, kenya. *East African Journal of Science, Technology and Innovation* **2**(1) (2020) <https://doi.org/10.37425/eajsti.v2i1.201>
- [8] Were, C.A., Okumu, T.A., Kimeli, P.: Feeding practices and animal-level factors associated with daily milk yield in lactating smallholder dairy cows in kiambu

- county, kenya. *Preventive Veterinary Medicine* **243**, 106607 (2025) <https://doi.org/10.1016/j.prevetmed.2025.106607>
- [9] Lesnoff, M., Corniaux, C., Hiernaux, P.: Sensitivity analysis of the recovery dynamics of a cattle population following drought in the sahel region. *Ecological Modelling* **232**, 28–39 (2012) <https://doi.org/10.1016/j.ecolmodel.2012.02.018>
 - [10] Tuffa, S., Treydte, A.C.: Modeling boran cattle populations under climate change and varying carrying capacity. *Ecological Modelling* **352**, 113–127 (2017) <https://doi.org/10.1016/j.ecolmodel.2017.02.031>
 - [11] Otte, M.J., Chilonda, P.: Cattle and small ruminant production systems in sub-saharan africa: A systematic review. Technical report, Food and Agriculture Organization of the United Nations, Rome (2002). <https://www.fao.org/4/Y4176E/y4176e00.htm>
 - [12] Meshesha, D.T., Mohammed, M., Yosuf, D.: Estimating carrying capacity and stocking rates of rangelands in harshin district, eastern somali region, ethiopia. *Ecology and Evolution* **9**(23), 13309–13319 (2019) <https://doi.org/10.1002/ece3.5786>
 - [13] Hary, I.: Derivation of steady state herd productivity using stage-structured population models and mathematical programming. *Agricultural Systems* **81**(2), 133–152 (2004) [https://doi.org/10.1016/S0308-521X\(03\)00194-X](https://doi.org/10.1016/S0308-521X(03)00194-X)