

Microplastic Contamination in Deep-Ocean Ecosystems: Distribution Patterns, Biological Uptake, and Remediation Prospects

Findings from a 36-Month Global Survey (2021–2024)

Sofia Andersson
Kwame Asante-Mensah
Yuki Tanaka
Rachel O'Brien

Institute of Marine Sciences, University of Bergen
Woods Hole Oceanographic Institution
Corresponding author: s.andersson@uib.no

Abstract

Microplastic pollution has emerged as one of the most pervasive environmental challenges of the 21st century, with recent evidence demonstrating contamination extending into hadal zones exceeding 10,000 meters depth. This paper presents findings from a 36-month global survey spanning 14 oceanic regions, employing a standardized trawl-and-core sampling methodology combined with Fourier-transform infrared spectroscopy (FTIR) for particle characterization. We document median concentrations of 4,200 particles/m³ in mesopelagic zones and 890 particles/kg in sediment cores, with polyethylene terephthalate (PET) and polystyrene (PS) comprising 67% of identified polymer types. Biological uptake analysis across 23 species reveals trophic magnification factors of 1.8–3.2 across three food-web levels. We further evaluate four candidate remediation technologies — bioreactor filtration, magnetic polymer capture, UV photocatalysis, and engineered microbial degradation — finding that no current technology achieves scalable deep-ocean remediation. Our data indicate that without dramatic reductions in plastic production, deep-ocean microplastic concentrations will increase by an estimated 47% by 2040.

Keywords: microplastics, ocean pollution, marine ecology, trophic magnification, remediation, deep-sea

1. Introduction

1.1 Global Plastic Production and Leakage

Annual global plastic production reached 400.3 million metric tons in 2022, with estimates suggesting that 8–12 million metric tons enter marine environments each year through riverine discharge, coastal erosion, atmospheric deposition, and direct dumping. Once in the ocean, UV radiation and mechanical action fragment macro-plastics into microplastics (particles 1 nm–5 mm) and nanoplastics (<1 nm), which resist conventional filtration and persist in the environment for

centuries.

1.2 Knowledge Gaps in Deep-Ocean Research

While surface and coastal microplastic contamination has been extensively documented, the deep ocean — which constitutes 95% of the biosphere by volume — remains relatively understudied due to the logistical and financial barriers associated with deep-sea sampling. Hadal trench sediments sampled opportunistically have yielded unexpectedly high concentrations, suggesting that deep-ocean basins may function as ultimate sinks for buoyant particles that aggregate with organic matter and sink through a process termed the biological pump.

2. Materials and Methods

2.1 Sampling Design

Sampling was conducted across 14 oceanic regions: North and South Atlantic, North and South Pacific, Indian Ocean, Arctic Ocean, Antarctic Circumpolar Current, and six marginal seas. At each station, we deployed a Manta trawl (330-micron mesh) for surface sampling, a vertically-integrated Multi-Level Trawl System (MLTS) for mesopelagic profiling, and gravity cores for sediment analysis. Samples were processed within 4 hours of collection to minimize contamination from shipboard sources.

2.2 Particle Characterization

All particles were characterized using a Bruker ALPHA II FTIR spectrometer with an ATR accessory. Polymer type, particle morphology (fragment, fiber, film, pellet), size distribution, and weathering index were recorded for each particle. Quality control protocols included procedural blanks, field blanks, and inter-laboratory comparisons with two partner institutions.

3. Results

3.1 Spatial Distribution

Microplastic concentrations showed strong spatial heterogeneity, with highest surface-water concentrations observed in the North Pacific Subtropical Gyre (median: 18,400 particles/m³) and lowest in the Southern Ocean (median: 340 particles/m³). Notably, mesopelagic concentrations were not well-predicted by surface concentrations, indicating that vertical transport processes other than simple settling — including zooplankton diel vertical migration and aggregate formation — play important roles in redistributing particles.

3.2 Biological Uptake and Trophic Transfer

Gastrointestinal content analysis of 1,847 individuals across 23 species (8 fish, 7 invertebrate, 4 cetacean, 4 seabird) found microplastics in 78% of mesopelagic fish, 64% of surface-dwelling seabirds, and 91% of filter-feeding invertebrates. Trophic magnification factors (TMF) calculated from species pairs with well-characterized predator-prey relationships ranged from 1.8 (copepod to sardine) to 3.2 (krill to Antarctic fur seal), indicating progressive bioaccumulation with trophic level.

4. Remediation Technology Assessment

We evaluated four remediation approaches based on technical readiness level, scalability, cost per cubic meter treated, and potential ecological side effects. Bioreactor filtration, which uses membrane systems with pore sizes of 0.1–1.0 micron, achieves >99% particle removal in controlled settings but is currently limited to volumes of <10,000 m³/day and cannot be deployed in open-ocean settings. Magnetic polymer capture using iron oxide-coated particles shows promise for localized hotspot treatment but introduces secondary contamination risks. UV photocatalysis achieves degradation of PS and PET but requires light penetration that limits its applicability below the photic zone. Engineered microbial degradation using *Ideonella sakaiensis* strains has demonstrated PET degradation rates of 0.13 mg/cm²/day under laboratory conditions, but in situ applications face challenges of dilution, temperature sensitivity, and competitive exclusion by native microbiota.

5. Projections and Policy Implications

Using a mass-balance transport model incorporating current production trends, degradation rates, and sinking fluxes, we project that deep-ocean microplastic concentrations will increase by 47% (95% CI: 38–61%) by 2040 under a business-as-usual scenario. Only scenarios involving a 60% or greater reduction in primary plastic production by 2030 yield stabilization of deep-ocean inventories within this century. These findings underscore the urgency of binding international agreements on plastic production limits, extending the scope of existing treaties such as the Basel Convention on hazardous wastes.

6. Conclusions

This survey provides the most comprehensive quantitative characterization of deep-ocean microplastic contamination to date, confirming that hadal and mesopelagic zones represent major global reservoirs of plastic pollution. The trophic magnification data demonstrate that deep-sea organisms are not isolated from surface pollution but are actively exposed through vertical transport pathways. Critically, no currently available remediation technology offers scalable solutions for deep-ocean cleanup. Preventing further contamination through upstream production controls is both more feasible and more cost-effective than remediation-based approaches.

References

-
- [1] Jambeck, J. R., et al. (2015). Plastic waste inputs from land into the ocean. *Science*, 347(6223), 768–771.
 - [2] Jamieson, A. J., et al. (2019). Microplastics and synthetic particles ingested by deep-sea amphipods in six of the deepest marine ecosystems on Earth. *Royal Society Open Science*, 6(2), 180667.
 - [3] Choy, C. A., et al. (2019). The vertical distribution and biological transport of marine microplastics across the epipelagic and mesopelagic water column. *Scientific Reports*, 9(1), 7843.
 - [4] Rochman, C. M., et al. (2021). Rethinking microplastics as a diverse contaminant suite. *Environmental Toxicology and Chemistry*, 38(4), 703–711.

- [5] Yoshida, S., et al. (2016). A bacterium that degrades and assimilates poly(ethylene terephthalate). *Science*, 351(6278), 1196–1199.
- [6] UNEP. (2023). *Turning off the Tap: How the World Can End Plastic Pollution and Create a Circular Economy*. United Nations Environment Programme, Nairobi.