

VIETNAM NATIONAL UNIVERSITY -
HOCHIMINH CITY
INTERNATIONAL UNIVERSITY

LESSON 4. Brownian motion

CHAPTER 2. Common stochastic processes

Course: Random processes

Lecturer: Le Nhat Tan, PhD

Lesson content

- Random walk

Lesson content

- Random walk
- Martingale processes

Lesson content

- Random walk
- Martingale processes
- Brownian motion

Symmetric Random walk

We toss a fair coin infinitely many times.

- Let X_i be a random variable such that

$$X_i(\omega_i) = \begin{cases} 1 & \text{if } \omega_i = H \\ -1 & \text{if } \omega_i = T. \end{cases}$$

Symmetric Random walk

We toss a fair coin infinitely many times.

- Let X_i be a random variable such that

$$X_i(\omega_i) = \begin{cases} 1 & \text{if } \omega_i = H \\ -1 & \text{if } \omega_i = T. \end{cases}$$

- X_i and X_j are independent random variables for any $i \neq j$.

Symmetric Random walk

We toss a fair coin infinitely many times.

- Let X_i be a random variable such that

$$X_i(\omega_i) = \begin{cases} 1 & \text{if } \omega_i = H \\ -1 & \text{if } \omega_i = T. \end{cases}$$

- X_i and X_j are independent random variables for any $i \neq j$.

- For each $k \in \mathbb{N}$, we denote $M_k = \sum_{i=1}^k X_i$.

Symmetric Random walk

We toss a fair coin infinitely many times.

- Let X_i be a random variable such that

$$X_i(\omega_i) = \begin{cases} 1 & \text{if } \omega_i = H \\ -1 & \text{if } \omega_i = T. \end{cases}$$

- X_i and X_j are independent random variables for any $i \neq j$.

- For each $k \in \mathbb{N}$, we denote $M_k = \sum_{i=1}^k X_i$.

- $(M_k)_{k \in \mathbb{N}}$: symmetric random walk.

Example

For each toss of a fair coin, you will win \$1 if it lands up heads and lose \$1 if it lands up tails. Suppose that you start the game with \$0 and denote M_k is the amount of money you obtain after k tosses. Then $(M_k)_{k \in \mathbb{N}}$ is a random walk.

Example

For each toss of a fair coin, you will win \$1 if it lands up heads and lose \$1 if it lands up tails. Suppose that you start the game with \$0 and denote M_k is the amount of money you obtain after k tosses. Then $(M_k)_{k \in \mathbb{N}}$ is a random walk.

- Play the game with your friend 10 times and plot your accumulated money after every play.

Example


For each toss of a fair coin, you will win \$1 if it lands up heads and lose \$1 if it lands up tails. Suppose that you start the game with \$0 and denote M_k is the amount of money you obtain after k tosses. Then $(M_k)_{k \in \mathbb{N}}$ is a random walk.

- Play the game with your friend 10 times and plot your accumulated money after every play.
- Write a R code to simulate the game

R code

```
par(mfrow=c(3,3))
N <- 20 ### number of playing game
M <- rep(0,N) ## create a space to store the evolution of M
M[1] <- 0 # the initial amount of money

for (i in 2:N) {
  X <- rbinom(n=1,size=1,prob=0.5) # simulate a Bernoulli
    random variable
  if (X==1) M[i] =M[i-1]+1 ## get $1 if head. i.e., X==1
  else M[i] =M[i-1]-1 ## lose $1 if head. i.e., X==0
}
plot(M, type="o", col="blue",xlab="number of playing
    games",ylab="the accumulated money")
```



Exercise

For each toss of a unfair coin, you will win \$2 if it lands up heads and lose \$1 if it lands up tails. The probability of landing up heads is 0.3 and the probability of landing up tails is 0.7. Suppose that you start the game with \$0 and denote M_k is the amount of money you obtain after k tosses.

Exercise





For each toss of a unfair coin, you will win \$2 if it lands up heads and lose \$1 if it lands up tails. The probability of landing up heads is 0.3 and the probability of landing up tails is 0.7. Suppose that you start the game with \$0 and denote M_k is the amount of money you obtain after k tosses.

- Should you play the game?

Exercise

For each toss of a unfair coin, you will win \$2 if it lands up heads and lose \$1 if it lands up tails. The probability of landing up heads is 0.3 and the probability of landing up tails is 0.7. Suppose that you start the game with \$0 and denote M_k is the amount of money you obtain after k tosses.

- Should you play the game?
- Write a R code to simulate possible results for playing the games 500 times



Fair games

For each toss of a fair coin, you will win \$1 if it lands up heads and lose \$1 if it lands up tails. Suppose that you start the game with \$0 and denote M_k is the amount of money you obtain after k tosses. Then $(M_k)_{k \in \mathbb{N}}$ is a random walk.

Fair games

For each toss of a fair coin, you will win \$1 if it lands up heads and lose \$1 if it lands up tails. Suppose that you start the game with \$0 and denote M_k is the amount of money you obtain after k tosses. Then $(M_k)_{k \in \mathbb{N}}$ is a random walk.

- Compute the probability of winning after playing the game 9 times

Fair games

For each toss of a fair coin, you will win \$1 if it lands up heads and lose \$1 if it lands up tails. Suppose that you start the game with \$0 and denote M_k is the amount of money you obtain after k tosses. Then $(M_k)_{k \in \mathbb{N}}$ is a random walk.

- Compute the probability of winning after playing the game 9 times
- Compute the probability of winning after playing the game 10 times

Fair games

For each toss of a fair coin, you will win \$1 if it lands up heads and lose \$1 if it lands up tails. Suppose that you start the game with \$0 and denote M_k is the amount of money you obtain after k tosses. Then $(M_k)_{k \in \mathbb{N}}$ is a random walk.

- Compute the probability of winning after playing the game 9 times
- Compute the probability of winning after playing the game 10 times
- Is this game fair, i.e., the chance of winning a fair game is equal for all players?

Fair games

For each toss of a fair coin, you will win \$1 if it lands up heads and lose \$1 if it lands up tails. Suppose that you start the game with \$0 and denote M_k is the amount of money you obtain after k tosses. Then $(M_k)_{k \in \mathbb{N}}$ is a random walk.

- Compute the probability of winning after playing the game 9 times
- Compute the probability of winning after playing the game 10 times
- Is this game fair, i.e., the chance of winning a fair game is equal for all players?
- A fair game is modeled by a martingale process.

Martingale processes

Consider a continuous-time stochastic process $(X_t)_{t \geq 0}$ adapted to the filtration $\{\mathcal{F}_t\}$ and satisfies the condition $E(|X_t|) < \infty, \forall t \geq 0$.

Martingale processes

Consider a continuous-time stochastic process $(X_t)_{t \geq 0}$ adapted to the filtration $\{\mathcal{F}_t\}$ and satisfies the condition $E(|X_t|) < \infty, \forall t \geq 0$.

- If $E(X_t | \mathcal{F}_s) = X_s, \forall 0 \leq s < t$: martingale process (no tendency to fall or rise)

Martingale processes

Consider a continuous-time stochastic process $(X_t)_{t \geq 0}$ adapted to the filtration $\{\mathcal{F}_t\}$ and satisfies the condition $E(|X_t|) < \infty, \forall t \geq 0$.

- If $E(X_t | \mathcal{F}_s) = X_s, \forall 0 \leq s < t$: martingale process (no tendency to fall or rise)
- If $E(X_t | \mathcal{F}_s) \geq X_s, \forall 0 \leq s \leq t$: submartingale (no tendency to fall, might have a tendency to rise)

Martingale processes

Consider a continuous-time stochastic process $(X_t)_{t \geq 0}$ adapted to the filtration $\{\mathcal{F}_t\}$ and satisfies the condition $E(|X_t|) < \infty, \forall t \geq 0$.

- If $E(X_t | \mathcal{F}_s) = X_s, \forall 0 \leq s < t$: martingale process (no tendency to fall or rise)
- If $E(X_t | \mathcal{F}_s) \geq X_s, \forall 0 \leq s \leq t$: submartingale (no tendency to fall, might have a tendency to rise)
- If $E(X_t | \mathcal{F}_s) < X_s, \forall 0 \leq s \leq t$: supermartingale (no tendency to rise, might have a tendency to fall).

Martingale processes

To prove a random process $(X_t)_{t \geq 0}$ be a martingale process under the filtration $(\mathcal{F}_t)_{t \geq 0}$, we need to prove three things:

- 1 $(X_t)_{t \geq 0}$ is adapted to $(\mathcal{F}_t)_{t \geq 0}$
- 2 $E[|X_t|] < +\infty$.
- 3 $E[X_t | \mathcal{F}_s] = X_s, \forall 0 < s \leq t$.

To prove the second, we may use the Holder inequality:

$$E[|X_t|] \leq \sqrt{E[X_t^2]} \text{ and the monotonicity property of expectation}$$
$$E[|X + Y|] \leq E[|X|] + E[|Y|].$$

Martingale processes

- If a stochastic process $(X_t)_{t \geq 0}$ is said to be a martingale process, without specifying the corresponding filtration, it is understood that the filtration is the natural filtration $(\mathcal{F}_t)_{t \geq 0}$ of the process, i.e., $\mathcal{F}_t = \sigma(X_s, 0 \leq s \leq t)$ (the σ -algebra generated from all random variables $X_s, 0 \leq s \leq t$)

Martingale processes

- If a stochastic process $(X_t)_{t \geq 0}$ is said to be a martingale process, without specifying the corresponding filtration, it is understood that the filtration is the natural filtration $(\mathcal{F}_t)_{t \geq 0}$ of the process, i.e., $\mathcal{F}_t = \sigma(X_s, 0 \leq s \leq t)$ (the σ -algebra generated from all random variables $X_s, 0 \leq s \leq t$)
- If a stochastic process $(X_t)_{t \geq 0}$ is said to be a martingale process with respect to another stochastic process $(Y_t)_{t \geq 0}$, the corresponding filtration is the natural filtration of the process $(Y_t)_{t \geq 0}$.

Some properties of a symmetric random walk

Consider a random walk $(M_k)_{k \geq 0}$.

- ① $M_k - M_l$ and $M_t - M_s$ are independent random variables for any $0 \leq k \leq l \leq s \leq t$.

- ②
$$E[M_k - M_l] = E\left[\sum_{i=l+1}^k X_i\right] = \sum_{i=l+1}^k E[X_i] = 0$$

- ③ Each increment $M_k - M_l$ has variance $k - l$, for any $l \leq k$.

- ④ $(M_k)_{k \in \mathbb{N}}$ is a martingale process. More precisely,

$$E[|M_k|] = E\left[\left|\sum_{i=1}^k X_i\right|\right] = E\left[\sum_{i=1}^k |X_i|\right] = k < +\infty \text{ and}$$

$$\begin{aligned} E[M_t | \mathcal{F}_s] &= E[(M_t - M_s) + M_s | \mathcal{F}_s] = E[M_t - M_s | \mathcal{F}_s] + M_s \\ &= E[M_t - M_s] + M_s = M_s. \end{aligned}$$

- ⑤ Quadratic variation up to time t , defined as

$$[M, M]_k = \sum_{j=1}^k (M_j - M_{j-1})^2 = k.$$

Exercise

Consider a game based on a standard normal random variable Z . At each run, the player will receive the money equals to the value of Z at that run. Suppose that you start the game with \$0 and denote M_k is the amount of money you obtain after k plays.

Exercise

Consider a game based on a standard normal random variable Z . At each run, the player will receive the money equals to the value of Z at that run. Suppose that you start the game with \$0 and denote M_k is the amount of money you obtain after k plays.

- Given that $\int_{1/\sqrt{10}}^{+\infty} \frac{1}{\sqrt{2\pi}} e^{-x^2/2} dx = 0.376$, find the probability that $M_{10} > 1$.

Exercise

Consider a game based on a standard normal random variable Z . At each run, the player will receive the money equals to the value of Z at that run. Suppose that you start the game with \$0 and denote M_k is the amount of money you obtain after k plays.

- Given that $\int_{1/\sqrt{10}}^{+\infty} \frac{1}{\sqrt{2\pi}} e^{-x^2/2} dx = 0.376$, find the probability that $M_{10} > 1$.
- Should you play the game, i.e., is this a fair game?

Exercise

Consider a game based on a standard normal random variable Z . At each run, the player will receive the money equals to the value of Z at that run. Suppose that you start the game with \$0 and denote M_k is the amount of money you obtain after k plays.










- Given that $\int_{1/\sqrt{10}}^{+\infty} \frac{1}{\sqrt{2\pi}} e^{-x^2/2} dx = 0.376$, find the probability that $M_{10} > 1$.
- Should you play the game, i.e., is this a fair game?
- Play the game 10 times and plot your accumulated money after every play.

Exercise

Consider a game based on a standard normal random variable Z . At each run, the player will receive the money equals to the value of Z at that run. Suppose that you start the game with \$0 and denote M_k is the amount of money you obtain after k plays.

- Given that $\int_{1/\sqrt{10}}^{+\infty} \frac{1}{\sqrt{2\pi}} e^{-x^2/2} dx = 0.376$, find the probability that $M_{10} > 1$.
- Should you play the game, i.e., is this a fair game?
- Play the game 10 times and plot your accumulated money after every play.
- Write a R code to simulate the game

```
par(mfrow=c(3,3))
N <- 20
M <- rep(0,N) ## create a space to store the evolution of M
M[1] <- 0 # the initial amount of money
for (i in 2:N) {
  X <- rnorm(n=1,mean=0,sd=1) # simulate a Bernoulli random
    variable
  M[i] =M[i-1]+X
}
plot(M, type="o", col="blue",xlab="k",ylab="M")
```



Exercise

Consider a game based on a standard normal random variable Z . At each run, the player will receive the money equals to \$0.3 plus the value of Z at that run. Suppose that you start the game with \$0 and denote M_k is the amount of money you obtain after k plays.

Exercise

Consider a game based on a standard normal random variable Z . At each run, the player will receive the money equals to \$0.3 plus the value of Z at that run. Suppose that you start the game with \$0 and denote M_k is the amount of money you obtain after k plays.

- Compute the probability that $M_{10} > 0$, given that

$$\int_0^{0.3\sqrt{10}} \frac{1}{\sqrt{2\pi}} e^{-x^2/2} dx = 0.3286.$$

Exercise

Consider a game based on a standard normal random variable Z . At each run, the player will receive the money equals to \$0.3 plus the value of Z at that run. Suppose that you start the game with \$0 and denote M_k is the amount of money you obtain after k plays.

- Compute the probability that $M_{10} > 0$, given that

$$\int_0^{0.3\sqrt{10}} \frac{1}{\sqrt{2\pi}} e^{-x^2/2} dx = 0.3286.$$

- Should you play the game, i.e., is this a fair game?





Exercise

Consider a game based on a standard normal random variable Z . At each run, the player will receive the money equals to \$0.3 plus the value of Z at that run. Suppose that you start the game with \$0 and denote M_k is the amount of money you obtain after k plays.

- Compute the probability that $M_{10} > 0$, given that

$$\int_0^{0.3\sqrt{10}} \frac{1}{\sqrt{2\pi}} e^{-x^2/2} dx = 0.3286.$$

- Should you play the game, i.e., is this a fair game?
- Write a R code to simulate the obtained result after playing the game 500 times.



Brownian motion

Let $(\Omega, \mathcal{F}, (\mathcal{F}_t), \mathcal{P})$ be a filtered probability space and let $(B_t)_{0 \leq t < \infty}$ be an adapted process of this space. The process $(B_t)_{0 \leq t < \infty}$ is called a standard Brownian motion if it satisfies the following properties:

- 1 $B_0 = 0$
- 2 Independent increments: $B_t - B_s$ is independent of \mathcal{F}_s , $0 \leq s < t$. That means

$$P(B_t - B_s \leq k | \mathcal{F}_s) = P(B_t - B_s \leq k).$$

- 3 Stationary increments:


$$B_t - B_s \stackrel{d}{=} B_{t-s} \sim N(0, \sqrt{t-s}), \quad \forall 0 < s < t.$$

- 4 Continuous paths: all sample paths of process $(B_t)_{t \geq 0}$ are almost surely continuous, i.e.

$$P(\omega \in \Omega | B_t(\omega) \text{ is a continuous sample path}) = 1.$$

Brownian motion

Brownian motion can be thought of as the motion of a particle that diffuses randomly along a line. At each point t , the particle's position B_t is normally distributed about the line with variance t , i.e., $B_t \sim N(0, t)$. As t increases, the particle's position is more diffuse.



Some interesting notes

Can we predict the value of B_t from time $s < t$? We cannot predict exactly the value of the B_t , but we know the possible range of values B_t .

Some interesting notes

Can we predict the value of B_t from time $s < t$? We cannot predict exactly the value of the B_t , but we know the possible range of values B_t .

- $B_t = B_s + \sqrt{t-s}N(0,1)$ or $B_t \sim N(B_s, \sqrt{t-s})$

Some interesting notes

Can we predict the value of B_t from time $s < t$? We cannot predict exactly the value of the B_t , but we know the possible range of values B_t .

- $B_t = B_s + \sqrt{t-s}N(0,1)$ or $B_t \sim N(B_s, \sqrt{t-s})$
- $P(B_s - \frac{\sqrt{t-s}}{2} < B_t < B_s + \frac{\sqrt{t-s}}{2}) = 38.2\%$

Some interesting notes

Can we predict the value of B_t from time $s < t$? We cannot predict exactly the value of the B_t , but we know the possible range of values B_t .

- $B_t = B_s + \sqrt{t-s}N(0,1)$ or $B_t \sim N(B_s, \sqrt{t-s})$
- $P(B_s - \frac{\sqrt{t-s}}{2} < B_t < B_s + \frac{\sqrt{t-s}}{2}) = 38.2\%$
- $P(B_s - \sqrt{t-s} < B_t < B_s + \sqrt{t-s}) = 68.2\%$

Some interesting notes

Can we predict the value of B_t from time $s < t$? We cannot predict exactly the value of the B_t , but we know the possible range of values B_t .

- $B_t = B_s + \sqrt{t-s}N(0,1)$ or $B_t \sim N(B_s, \sqrt{t-s})$
- $P(B_s - \frac{\sqrt{t-s}}{2} < B_t < B_s + \frac{\sqrt{t-s}}{2}) = 38.2\%$
- $P(B_s - \sqrt{t-s} < B_t < B_s + \sqrt{t-s}) = 68.2\%$
- $P(B_s - \frac{3\sqrt{t-s}}{2} < B_t < B_s + \frac{3\sqrt{t-s}}{2}) = 86.6\%$

Some interesting notes

Can we predict the value of B_t from time $s < t$? We cannot predict exactly the value of the B_t , but we know the possible range of values B_t .

- $B_t = B_s + \sqrt{t-s}N(0,1)$ or $B_t \sim N(B_s, \sqrt{t-s})$
- $P(B_s - \frac{\sqrt{t-s}}{2} < B_t < B_s + \frac{\sqrt{t-s}}{2}) = 38.2\%$
- $P(B_s - \sqrt{t-s} < B_t < B_s + \sqrt{t-s}) = 68.2\%$
- $P(B_s - \frac{3\sqrt{t-s}}{2} < B_t < B_s + \frac{3\sqrt{t-s}}{2}) = 86.6\%$
- $P(B_s - 2\sqrt{t-s} < B_t < B_s + 2\sqrt{t-s}) = 95.4\%$

Some interesting notes

Can we predict the value of B_t from time $s < t$? We cannot predict exactly the value of the B_t , but we know the possible range of values B_t .

- $B_t = B_s + \sqrt{t-s}N(0,1)$ or $B_t \sim N(B_s, \sqrt{t-s})$
- $P(B_s - \frac{\sqrt{t-s}}{2} < B_t < B_s + \frac{\sqrt{t-s}}{2}) = 38.2\%$
- $P(B_s - \sqrt{t-s} < B_t < B_s + \sqrt{t-s}) = 68.2\%$
- $P(B_s - \frac{3\sqrt{t-s}}{2} < B_t < B_s + \frac{3\sqrt{t-s}}{2}) = 86.6\%$
- $P(B_s - 2\sqrt{t-s} < B_t < B_s + 2\sqrt{t-s}) = 95.4\%$
- $P(B_s - \frac{5\sqrt{t-s}}{2} < B_t < B_s + \frac{5\sqrt{t-s}}{2}) = 98.8\%$

Some interesting notes

Can we predict the value of B_t from time $s < t$? We cannot predict exactly the value of the B_t , but we know the possible range of values B_t .

- $B_t = B_s + \sqrt{t-s}N(0, 1)$ or $B_t \sim N(B_s, \sqrt{t-s})$
- $P(B_s - \frac{\sqrt{t-s}}{2} < B_t < B_s + \frac{\sqrt{t-s}}{2}) = 38.2\%$
- $P(B_s - \sqrt{t-s} < B_t < B_s + \sqrt{t-s}) = 68.2\%$
- $P(B_s - \frac{3\sqrt{t-s}}{2} < B_t < B_s + \frac{3\sqrt{t-s}}{2}) = 86.6\%$
- $P(B_s - 2\sqrt{t-s} < B_t < B_s + 2\sqrt{t-s}) = 95.4\%$
- $P(B_s - \frac{5\sqrt{t-s}}{2} < B_t < B_s + \frac{5\sqrt{t-s}}{2}) = 98.8\%$
- $P(B_s - 3\sqrt{t-s} < B_t < B_s + 3\sqrt{t-s}) = 99.8\%$

Model standard Brownian motion

Suppose we want to model a Brownian motion in a time interval $[0, T]$.

- We divide the interval into n equal subintervals by discrete time points $0 = t_0 < t_1 < \cdots < t_n = T$, with the time step $h = t_i - t_{i-1}$.

Model standard Brownian motion

Suppose we want to model a Brownian motion in a time interval $[0, T]$.

- We divide the interval into n equal subintervals by discrete time points $0 = t_0 < t_1 < \dots < t_n = T$, with the time step $h = t_i - t_{i-1}$.
- The value of B_{t_i} is determined by a recursive formula:
 $B_{t_i} = B_{t_{i-1}} + \sqrt{h} * X$, where $X \sim N(0, 1)$.

Model standard Brownian motion


Suppose we want to model a Brownian motion in a time interval $[0, T]$.

- We divide the interval into n equal subintervals by discrete time points $0 = t_0 < t_1 < \dots < t_n = T$, with the time step $h = t_i - t_{i-1}$.
- The value of B_{t_i} is determined by a recursive formula:
 $B_{t_i} = B_{t_{i-1}} + \sqrt{h} * X$, where $X \sim N(0, 1)$.
- All we need to simulate the Brownian motion is to simulate a standard normal random variable, which can be done by using command “`rnorm(1)`” in software R.

```
BM=function(T,N) {  
  # T=1 expiry time  
  # N=100 number of simulation points  
  h=T/N # the timestep of the simulation  
  X=rep(0, (N+1)) # create space to store value of Brownian  
    motion  
  X[1]=0 ## start at 0  
  for(i in 1:N) { X[i+1]=X[i] +sqrt(h)*rnorm(1)}  
  return(X)  
}
```

A sample of standard Brownian motion


```
h <- 1/1000  
t=seq(0,1,by=h)  
plot(t,BM(1,N=1000), type="l",  
      col="blue",xlab="t",ylab="B")
```




Plot multiple samples of Brownian motion

```
BMSamplepaths <- function(T,N,nt)
{h=T/N ## time step
#nt: number of samples
t=seq(0,T,by=h) # produce a sequence of time points
X=matrix(rep(0,length(t)*nt), nrow=nt)
## each row of the matrix stores one sample
# #return(X)
for (i in 1:nt) {X[i,]= BM(T=T,N=N)}
# # ##Plot
ymax=max(X); ymin=min(X) #bounds for simulated prices
plot(t,X[1,],t='l',main='3 sample paths of standard
      Brownian motions
with parameters T =1, N =3',ylim=c(ymin, ymax), col=1,
ylab="Price P(t)",xlab="time t")
for(i in 2:nt){lines(t,X[i,], t='l',ylim=c(ymin,
```

**3 sample paths of standard Brownian motions
with parameters $T = 1$, $N = 200$**



**1000 sample paths of standard Brownian motions
with parameters $T = 1$, $N = 200$**



Example

A particle's position is modelled by a standard Brownian motion. If the particle is at position 1 at time $t = 2$, then find the probability that its position is at most 3 at time $t = 5$.

Example

A particle's position is modelled by a standard Brownian motion. If the particle is at position 1 at time $t = 2$, then find the probability that its position is at most 3 at time $t = 5$.

- Let $(B_t)_{t \geq 0}$ be the position of the particle at time t is B_t .

Example

A particle's position is modelled by a standard Brownian motion. If the particle is at position 1 at time $t = 2$, then find the probability that its position is at most 3 at time $t = 5$.

- Let $(B_t)_{t \geq 0}$ be the position of the particle at time t is B_t .
- The position of the particle at time $t = 2$ and $t = 5$ are B_2 and B_5 .

Example

A particle's position is modelled by a standard Brownian motion. If the particle is at position 1 at time $t = 2$, then find the probability that its position is at most 3 at time $t = 5$.

- Let $(B_t)_{t \geq 0}$ be the position of the particle at time t is B_t .
- The position of the particle at time $t = 2$ and $t = 5$ are B_2 and B_5 .
- The desired probability is

$$\begin{aligned} P(B_5 \leq 3 | B_2 = 1) &= P(B_5 - B_2 + B_2 \leq 3 | B_2 = 1) \\ &= P(B_5 - B_2 \leq 2 | B_2 = 1) = P(B_5 - B_2 \leq 2) \\ &= P(B_3 \leq 2) = \int_{-\infty}^2 \frac{1}{\sqrt{3.2\pi}} e^{-x^2/(2.3)} dx \approx 0.875893. \end{aligned}$$

Example

A particle's position is modelled by a standard Brownian motion. If the particle is at position 1 at time $t = 2$, then find the probability that its position is at most 3 at time $t = 5$.

- Let $(B_t)_{t \geq 0}$ be the position of the particle at time t is B_t .
- The position of the particle at time $t = 2$ and $t = 5$ are B_2 and B_5 .
- The desired probability is

$$\begin{aligned} P(B_5 \leq 3 | B_2 = 1) &= P(B_5 - B_2 + B_2 \leq 3 | B_2 = 1) \\ &= P(B_5 - B_2 \leq 2 | B_2 = 1) = P(B_5 - B_2 \leq 2) \\ &= P(B_3 \leq 2) = \int_{-\infty}^2 \frac{1}{\sqrt{3.2\pi}} e^{-x^2/(2.3)} dx \approx 0.875893. \end{aligned}$$

- Use software R with the command `pnorm(2,0,sqrt(3))` to obtain the result.

Exercise

A particle's position is modelled by a standard Brownian motion. Find the probability that the particle's position at time $t = 3$ is above the level 1.5.

Brownian motions are nowhere differentiable

- Brownian motion paths are nowhere differentiable.

Brownian motions are nowhere differentiable

- Brownian motion paths are nowhere differentiable.
- The jagged character of these paths remains jagged at all time scales.

Brownian motions are nowhere differentiable

- Brownian motion paths are nowhere differentiable.
- The jagged character of these paths remains jagged at all time scales.
- $\frac{d}{dt}B_t = \lim_{h \rightarrow 0} \frac{B_{t+h} - B_t}{h}.$

Brownian motions are nowhere differentiable

- Brownian motion paths are nowhere differentiable.
- The jagged character of these paths remains jagged at all time scales.
- $\frac{d}{dt}B_t = \lim_{h \rightarrow 0} \frac{B_{t+h} - B_t}{h}$.
- $\frac{B_{t+h} - B_t}{h}$ has the same distribution as $\frac{B_h}{h}$, which is normal with mean 0 and variance $\frac{1}{h}$.

Brownian motions are nowhere differentiable

- Brownian motion paths are nowhere differentiable.
- The jagged character of these paths remains jagged at all time scales.
- $\frac{d}{dt}B_t = \lim_{h \rightarrow 0} \frac{B_{t+h} - B_t}{h}$.
- $\frac{B_{t+h} - B_t}{h}$ has the same distribution as $\frac{B_h}{h}$, which is normal with mean 0 and variance $\frac{1}{h}$.
- As h tends to 0, the variance of $\frac{B_{t+h} - B_t}{h}$ tends to infinity.

Brownian motions are nowhere differentiable

- Brownian motion paths are nowhere differentiable.
- The jagged character of these paths remains jagged at all time scales.
- $\frac{d}{dt}B_t = \lim_{h \rightarrow 0} \frac{B_{t+h} - B_t}{h}$.
- $\frac{B_{t+h} - B_t}{h}$ has the same distribution as $\frac{B_h}{h}$, which is normal with mean 0 and variance $\frac{1}{h}$.
- As h tends to 0, the variance of $\frac{B_{t+h} - B_t}{h}$ tends to infinity.
- Since the difference quotient takes arbitrarily large values, the limit, and hence the derivative, does not exist.

Example

For $0 < s < t$, find the distribution of $B_s + B_t$.

Example

For $0 < s < t$, find the distribution of $B_s + B_t$.

- We express $B_s + B_t = 2B_s + (B_t - B_s)$.

Example

For $0 < s < t$, find the distribution of $B_s + B_t$.

- We express $B_s + B_t = 2B_s + (B_t - B_s)$.
- $B_s \sim N(0, s)$ and $B_t - B_s \sim B_{t-s} \sim N(0, t - s)$

Example

For $0 < s < t$, find the distribution of $B_s + B_t$.

- We express $B_s + B_t = 2B_s + (B_t - B_s)$.
- $B_s \sim N(0, s)$ and $B_t - B_s \sim B_{t-s} \sim N(0, t - s)$
- B_s and $B_t - B_s$ are independent random variables.

Example

For $0 < s < t$, find the distribution of $B_s + B_t$.

- We express $B_s + B_t = 2B_s + (B_t - B_s)$.
- $B_s \sim N(0, s)$ and $B_t - B_s \sim B_{t-s} \sim N(0, t - s)$
- B_s and $B_t - B_s$ are independent random variables.
- $E(B_s + B_t) = E(2B_s + B_t - B_s) = E(2B_s) + E(B_t - B_s) = 0$ and

$$\text{Var}(B_s + B_t) = \text{Var}(2B_s) + \text{Var}(B_t - B_s) = 4s + t - s = 3s + t.$$

Example

For $0 < s < t$, find the distribution of $B_s + B_t$.

- We express $B_s + B_t = 2B_s + (B_t - B_s)$.
- $B_s \sim N(0, s)$ and $B_t - B_s \sim B_{t-s} \sim N(0, t - s)$
- B_s and $B_t - B_s$ are independent random variables.
- $E(B_s + B_t) = E(2B_s + B_t - B_s) = E(2B_s) + E(B_t - B_s) = 0$ and

$$\text{Var}(B_s + B_t) = \text{Var}(2B_s) + \text{Var}(B_t - B_s) = 4s + t - s = 3s + t.$$

- $B_s + B_t \sim N(0, 3s + t)$.

Exercise

Find the covariance of B_s and B_t .

Markov property

Brownian motion is a Markov process. Let $(B_t)_{t \geq 0}$ be a Brownian motion under the probability space $(\Omega, \mathcal{F}, \mathcal{F}_t, \mathcal{P})$. Then B_t is also a Markov process.

Exercises

Using Holder's inequality $E[|X_t|] \leq \sqrt{E[X_t^2]}$ to prove:

- ① $E[|B_t|] \leq \sqrt{t}.$
- ② $E[|B_t^2 - t|] < 2t$

Martingale property

Prove the following random processes are martingale:

- 1 Brownian motion (B_t) is a martingale process.
- 2 Let $Y_t = B_t^2 - t$, for $t \geq 0$. Show that $(Y_t)_{t \geq 0}$ is a martingale with respect to Brownian motion.
- 3 $(Z_t)_{t \geq 0}$, with $Z_t = \exp(\sigma B_t - \frac{1}{2}\sigma^2 t)$ (σ is a positive constant), is a martingale with respect to the standard Brownian motion.

Solution

- Because B_t is completely determined under the information available in σ -algebra \mathcal{F}_t , is Z_t so. In other words, $(Z_t)_{t \geq 0}$ is an adapted process with the filtration $(\mathcal{F}_t)_{t \leq 0}$.

Solution

- Because B_t is completely determined under the information available in σ -algebra \mathcal{F}_t , is Z_t so. In other words, $(Z_t)_{t \geq 0}$ is an adapted process with the filtration $(\mathcal{F}_t)_{t \leq 0}$.
- If $X \sim N(\mu, \sigma)$ (then $E(\exp(X)) = \exp(\mu + \frac{1}{2}\sigma^2)$).

Solution

- Because B_t is completely determined under the information available in σ -algebra \mathcal{F}_t , is Z_t so. In other words, $(Z_t)_{t \geq 0}$ is an adapted process with the filtration $(\mathcal{F}_t)_{t \leq 0}$.
- If $X \sim N(\mu, \sigma)$ (then $E(\exp(X)) = \exp(\mu + \frac{1}{2}\sigma^2)$).
- $\sigma B_t - \frac{1}{2}\sigma^2 t \sim N(-\frac{1}{2}\sigma^2 t, \sigma^2 t) \Rightarrow E(|Z_t|) = E(Z_t) = 1 < +\infty$.

Solution

- Because B_t is completely determined under the information available in σ -algebra \mathcal{F}_t , is Z_t so. In other words, $(Z_t)_{t \geq 0}$ is an adapted process with the filtration $(\mathcal{F}_t)_{t \leq 0}$.
- If $X \sim N(\mu, \sigma)$ (then $E(\exp(X)) = \exp(\mu + \frac{1}{2}\sigma^2)$).
- $\sigma B_t - \frac{1}{2}\sigma^2 t \sim N(-\frac{1}{2}\sigma^2 t, \sigma^2 t) \Rightarrow E(|Z_t|) = E(Z_t) = 1 < +\infty$.
- For $0 \leq s \leq t$, we have

$$\begin{aligned} E[Z_t | \mathcal{F}_s] &= E[\exp(\sigma B_t - \frac{1}{2}\sigma^2 t) | \mathcal{F}_s] \\ &= \exp(-\frac{1}{2}\sigma^2 t) E[\exp(\sigma(B_t - B_s + B_s)) | \mathcal{F}_s] \\ &= \exp(\sigma B_s - \frac{1}{2}\sigma^2 t) E[\exp(\sigma(B_t - B_s)) | \mathcal{F}_s] \\ &= \exp(\sigma B_s - \frac{1}{2}\sigma^2 t) E[\exp(\sigma(B_t - B_s))] \end{aligned}$$

Solution

- As $\sigma(B_t - B_s) \sim N(0, \sigma^2(t - s))$,
 $E[\exp(\sigma(B_t - B_s))] = \exp(\frac{1}{2}\sigma^2(t - s)).$

Solution

- As $\sigma(B_t - B_s) \sim N(0, \sigma^2(t - s))$,
$$E[\exp(\sigma(B_t - B_s))] = \exp\left(\frac{1}{2}\sigma^2(t - s)\right).$$
- $$E[Z_t | \mathcal{F}_s] = \exp\left(\sigma B_s - \frac{1}{2}\sigma^2 t\right) \exp\left(\frac{1}{2}\sigma^2(t - s)\right) = \exp\left(\sigma B_s - \frac{1}{2}\sigma^2 s\right) = Z_s.$$

Solution

- As $\sigma(B_t - B_s) \sim N(0, \sigma^2(t - s))$,
$$E[\exp(\sigma(B_t - B_s))] = \exp\left(\frac{1}{2}\sigma^2(t - s)\right).$$
- $$E[Z_t | \mathcal{F}_s] = \exp\left(\sigma B_s - \frac{1}{2}\sigma^2 t\right) \exp\left(\frac{1}{2}\sigma^2(t - s)\right) = \exp\left(\sigma B_s - \frac{1}{2}\sigma^2 s\right) = Z_s.$$
- (Z_t) is a martingale process.

Example

Let $S_t = S_0 e^{X_t}$, with $X_t = \mu t + \sigma B_t$, and $(B_t)_{t \geq 0}$ is the standard Brownian motion. Let $r = \mu + \sigma^2/2$. Show that $e^{-rt} S_t$ is a martingale with respect to standard Brownian motion.

Example

Let $S_t = S_0 e^{X_t}$, with $X_t = \mu t + \sigma B_t$, and $(B_t)_{t \geq 0}$ is the standard Brownian motion. Let $r = \mu + \sigma^2/2$. Show that $e^{-rt} S_t$ is a martingale with respect to standard Brownian motion.

- We have $e^{-rt} S_t = e^{-(\mu + \sigma^2/2)t} S_0 e^{\mu t + \sigma B_t} = S_0 e^{\sigma B_t - (\sigma^2/2)t}$.

Example

Let $S_t = S_0 e^{X_t}$, with $X_t = \mu t + \sigma B_t$, and $(B_t)_{t \geq 0}$ is the standard Brownian motion. Let $r = \mu + \sigma^2/2$. Show that $e^{-rt} S_t$ is a martingale with respect to standard Brownian motion.

- We have $e^{-rt} S_t = e^{-(\mu + \sigma^2/2)t} S_0 e^{\mu t + \sigma B_t} = S_0 e^{\sigma B_t - (\sigma^2/2)t}$.
- As $e^{\sigma B_t - (\sigma^2/2)t}$ is a martingale with respect to the standard Brownian motion, so is $e^{-rt} S_t$.

Quadratic variation of Brownian motion

- The paths of Brownian motion are unusual in that their quadratic variation is not zero. This makes stochastic calculus different from ordinary calculus.

Quadratic variation of Brownian motion

- The paths of Brownian motion are unusual in that their quadratic variation is not zero. This makes stochastic calculus different from ordinary calculus.
- Most functions have continuous derivatives and hence their quadratic variations are zero.

Quadratic variation of Brownian motion

- The paths of Brownian motion are unusual in that their quadratic variation is not zero. This makes stochastic calculus different from ordinary calculus.
- Most functions have continuous derivatives and hence their quadratic variations are zero.
- The paths of Brownian motion, on the other hand, cannot be differentiated with respect to the time variable.

Quadratic variation of Brownian motion

- The paths of Brownian motion are unusual in that their quadratic variation is not zero. This makes stochastic calculus different from ordinary calculus.
- Most functions have continuous derivatives and hence their quadratic variations are zero.
- The paths of Brownian motion, on the other hand, cannot be differentiated with respect to the time variable.
- It can be proved that

$$[W, W](T) = \lim_{\Pi \rightarrow 0} \sum_{j=0}^{n-1} [W(t_{j+1}) - W(t_j)]^2 = T, \forall T \geq 0 \text{ almost}$$

surely (there might be some paths of the Brownian motion for which the assertion $[W, W](T) = T$ is not true, but the set of such paths has zero probability).

Quadratic variation of Brownian motion

- The paths of Brownian motion are unusual in that their quadratic variation is not zero. This makes stochastic calculus different from ordinary calculus.
- Most functions have continuous derivatives and hence their quadratic variations are zero.
- The paths of Brownian motion, on the other hand, cannot be differentiated with respect to the time variable.
- It can be proved that

$$[W, W](T) = \lim_{n \rightarrow \infty} \sum_{j=0}^{n-1} [W(t_{j+1}) - W(t_j)]^2 = T, \forall T \geq 0 \text{ almost}$$

surely (there might be some paths of the Brownian motion for which the assertion $[W, W](T) = T$ is not true, but the set of such paths has zero probability).

- Brownian motion accumulates quadratic variation at rate one per unit time: $dB_t^2 = dt$

Brownian motion starts at $x \neq 0$

For real value x , the process defined by $X_t = x + B_t$, for $t \geq 0$ is called Brownian motion started at x .

Brownian motion starts at $x \neq 0$

For real value x , the process defined by $X_t = x + B_t$, for $t \geq 0$ is called Brownian motion started at x .

- $X_0 = x, X_t \sim N(x, t)$

Brownian motion starts at $x \neq 0$

For real value x , the process defined by $X_t = x + B_t$, for $t \geq 0$ is called Brownian motion started at x .

- $X_0 = x, X_t \sim N(x, t)$
- Independent increments:

$$P(X_t - X_s \leq k | \mathcal{F}_s) = P(X_t - X_s \leq k).$$

Brownian motion starts at $x \neq 0$

For real value x , the process defined by $X_t = x + B_t$, for $t \geq 0$ is called Brownian motion started at x .

- $X_0 = x, X_t \sim N(x, t)$
- Independent increments:

$$P(X_t - X_s \leq k | \mathcal{F}_s) = P(X_t - X_s \leq k).$$

- $X_t - X_s \stackrel{d}{=} B_{t-s} \sim N(0, t-s), \forall 0 < s < t.$

Brownian motion starts at $x \neq 0$

For real value x , the process defined by $X_t = x + B_t$, for $t \geq 0$ is called Brownian motion started at x .

- $X_0 = x, X_t \sim N(x, t)$
- Independent increments:

$$P(X_t - X_s \leq k | \mathcal{F}_s) = P(X_t - X_s \leq k).$$

- $X_t - X_s \stackrel{d}{=} B_{t-s} \sim N(0, t-s), \forall 0 < s < t.$
- Continuous paths:

$$P(\omega \in \Omega | X_t(\omega) \text{ is a continuous sample path}) = 1.$$

Brownian motion starts at $x \neq 0$

For real value x , the process defined by $X_t = x + B_t$, for $t \geq 0$ is called Brownian motion started at x .

- $X_0 = x, X_t \sim N(x, t)$
- Independent increments:

$$P(X_t - X_s \leq k | \mathcal{F}_s) = P(X_t - X_s \leq k).$$

- $X_t - X_s \stackrel{d}{=} B_{t-s} \sim N(0, t-s), \forall 0 < s < t.$
- Continuous paths:

$$P(\omega \in \Omega | X_t(\omega) \text{ is a continuous sample path}) = 1.$$

- The probability density function of X_t is given by

$$K_t(x, y) = \frac{1}{\sqrt{2\pi t}} e^{-(y-x)^2/(2t)}.$$

Exercise

- 1 Write a R code to simulate a Brownian motion start at $x = 5$.
- 2 A particle's position is modelled by a Brownian motion start at $x = 1$. Find the probability that its position is at most 3 at time $t = 3$.

Example

Let $(X_t)_{t \geq 0}$ be a Brownian motion started at $x = 3$. Find $P(X_2 > 0)$.

Example

Let $(X_t)_{t \geq 0}$ be a Brownian motion started at $x = 3$. Find $P(X_2 > 0)$.

- $X_t = B_t + 3$, where $(B_t)_{t \geq 0}$ is a standard Brownian motion.

Example

Let $(X_t)_{t \geq 0}$ be a Brownian motion started at $x = 3$. Find $P(X_2 > 0)$.

- $X_t = B_t + 3$, where $(B_t)_{t \geq 0}$ is a standard Brownian motion.
- $P(X_2 > 0) = P(B_t + 3 > 0) = P(B_t > -3) = \int_{-3}^{\infty} \frac{1}{\sqrt{4\pi}} e^{-x^2/4} dx = 0.983.$

Example

Let $(X_t)_{t \geq 0}$ be a Brownian motion started at $x = 3$. Find $P(X_2 > 0)$.

- $X_t = B_t + 3$, where $(B_t)_{t \geq 0}$ is a standard Brownian motion.
- $P(X_2 > 0) = P(B_t + 3 > 0) = P(B_t > -3) = \int_{-3}^{\infty} \frac{1}{\sqrt{4\pi}} e^{-x^2/4} dx = 0.983$.
- In R, type `1-pnorm(-3,0,sqrt(2))` will help us compute the integral.

Conditional distribution of Brownian motion

Compute $E[B_4|B_2]$ and $Var[B_4|B_2]$.

Conditional distribution of Brownian motion

Compute $E[B_4|B_2]$ and $Var[B_4|B_2]$.

- $$E[B_4|B_2] = E[B_4 - B_2 + B_2|B_2] = E[B_4 - B_2|B_2] + E[B_2|B_2] = E[B_4 - B_2] + B_2 = B_2$$

Conditional distribution of Brownian motion

Compute $E[B_4|B_2]$ and $Var[B_4|B_2]$.

- $E[B_4|B_2] = E[B_4 - B_2 + B_2|B_2] = E[B_4 - B_2|B_2] + E[B_2|B_2] = E[B_4 - B_2] + B_2 = B_2$
- $Var[B_4|B_2] = Var[B_4 - B_2 + B_2|B_2] = Var[B_4 - B_2|B_2] + Var[B_2|B_2] = Var[B_4 - B_2] + 0 = 2.$

Conditional distribution of Brownian motion

Compute $E[B_2|B_5]$ and $Var[B_2|B_5]$.

Conditional distribution of Brownian motion

Compute $E[B_2|B_5]$ and $Var[B_2|B_5]$.

- $E[B_2|B_5] = E[B_2|B_5 - B_2 + B_2]$

Conditional distribution of Brownian motion

Compute $E[B_2|B_5]$ and $Var[B_2|B_5]$.

- $E[B_2|B_5] = E[B_2|B_5 - B_2 + B_2]$
- Let $X = B_2$, $U = B_5 - B_2$, $Y = B_5$

Conditional distribution of Brownian motion

Compute $E[B_2|B_5]$ and $Var[B_2|B_5]$.

- $E[B_2|B_5] = E[B_2|B_5 - B_2 + B_2]$
- Let $X = B_2$, $U = B_5 - B_2$, $Y = B_5$
- $f_X(x) = \frac{1}{\sqrt{4\pi}} e^{-x^2/4}$

Conditional distribution of Brownian motion

Compute $E[B_2|B_5]$ and $Var[B_2|B_5]$.

- $E[B_2|B_5] = E[B_2|B_5 - B_2 + B_2]$
- Let $X = B_2$, $U = B_5 - B_2$, $Y = B_5$
- $f_X(x) = \frac{1}{\sqrt{4\pi}} e^{-x^2/4}$
- $f_U(u) = \frac{1}{\sqrt{6\pi}} e^{-u^2/6}$

Conditional distribution of Brownian motion

Compute $E[B_2|B_5]$ and $\text{Var}[B_2|B_5]$.

- $E[B_2|B_5] = E[B_2|B_5 - B_2 + B_2]$
- Let $X = B_2$, $U = B_5 - B_2$, $Y = B_5$

- $f_X(x) = \frac{1}{\sqrt{4\pi}} e^{-x^2/4}$

- $f_U(u) = \frac{1}{\sqrt{6\pi}} e^{-u^2/6}$

- $f_Y(y) = \frac{1}{\sqrt{10\pi}} e^{-y^2/10}$

Conditional distribution of Brownian motion

Compute $E[B_2|B_5]$ and $\text{Var}[B_2|B_5]$.

- $E[B_2|B_5] = E[B_2|B_5 - B_2 + B_2]$
- Let $X = B_2$, $U = B_5 - B_2$, $Y = B_5$

- $f_X(x) = \frac{1}{\sqrt{4\pi}} e^{-x^2/4}$

- $f_U(u) = \frac{1}{\sqrt{6\pi}} e^{-u^2/6}$

- $f_Y(y) = \frac{1}{\sqrt{10\pi}} e^{-y^2/10}$

- $f_{X|Y}(x|y) = \frac{f_{X,Y}(x,y)}{f_Y(y)} = \frac{f_{X,U}(x,y-x)}{f_Y(y)} = \frac{f_X(x)f_U(y-x)}{f_Y(y)} =$

$$\frac{1}{\sqrt{2\pi \frac{6}{5}}} \exp\left(-\frac{(x - \frac{2y}{5})^2}{2\frac{6}{5}}\right)$$

Conditional distribution of Brownian motion

Compute $E[B_2|B_5]$ and $Var[B_2|B_5]$.

Conditional distribution of Brownian motion

Compute $E[B_2|B_5]$ and $Var[B_2|B_5]$.

- The conditional distribution of B_2 given B_5 is normal with mean $2y/5$ and variance $6/5$.

Conditional distribution of Brownian motion

Compute $E[B_2|B_5]$ and $Var[B_2|B_5]$.

- The conditional distribution of B_2 given B_5 is normal with mean $2y/5$ and variance $6/5$.
- $E[B_2|B_5 = y] = 2y/5$ and $Var[B_2|B_5 = y] = 6/5$

Conditional distribution of Brownian motion

Compute $E[B_2|B_5]$ and $Var[B_2|B_5]$.

- The conditional distribution of B_2 given B_5 is normal with mean $2y/5$ and variance $6/5$.
- $E[B_2|B_5 = y] = 2y/5$ and $Var[B_2|B_5 = y] = 6/5$
- $E[B_2|B_5] = 2B_5/5$ and $Var[B_2|B_5] = 6/5$

Stopping time

For a continuous-time stochastic process $(X_t)_{t \geq 0}$ on a filtered probability space $(\Omega, \mathcal{F}, (\mathcal{F}_t)_{t \geq 0}, \mathcal{P})$, a stopping time τ with respect to $(X_t)_{t \geq 0}$ is a non-negative random variable such that the event $\tau \leq t$ depends only on the information available up to time t , but not after time t . In other words, $\{\tau \leq t\} \in \mathcal{F}_t$.

Stopping time

For a continuous-time stochastic process $(X_t)_{t \geq 0}$ on a filtered probability space $(\Omega, \mathcal{F}, (\mathcal{F}_t)_{t \geq 0}, \mathcal{P})$, a stopping time τ with respect to $(X_t)_{t \geq 0}$ is a non-negative random variable such that the event $\tau \leq t$ depends only on the information available up to time t , but not after time t . In other words, $\{\tau \leq t\} \in \mathcal{F}_t$.

- τ is a stopping time with respect to $(\mathcal{F}_t)_{t \geq 0}$ if at any time t , it can be decided whether the event $\tau \leq t$ has already occurred or not, just based on the information contained in \mathcal{F}_t generated by $(X_s)_{0 \leq s \leq t}$.

Stopping time:example

- The day you get married is a stopping time.

Stopping time:example

- The day you get married is a stopping time.
- A stock share price is \$10. The first time the stock price hits \$12 is a stopping time.

Stopping time:example

- The day you get married is a stopping time.
- A stock share price is \$10. The first time the stock price hits \$12 is a stopping time.
- The day you will buy your last car is not a stopping time.

Stopping time: Proposition

Let τ and θ be stopping times with respect to the probability space $(\Omega, \mathcal{F}, (\mathcal{F}_t)_{t \geq 0}, \mathcal{P})$.

- i) Every constant time is a stopping time.
- ii) The minimum $\tau \wedge \theta = \min(\tau, \theta)$ is also a stopping time.
- iii) The maximum $\tau \vee \theta = \max(\tau, \theta)$ is also a stopping time.

Proof

- i) Let a is a constant time. The event $\{a \leq t\}$ is clearly belong to \mathcal{F}_t .
- ii) $\{\tau \wedge \theta > t\} = \{\min(\tau, \theta) > t\} = \{\tau > t\} \cap \{\theta > t\} \in \mathcal{F}_t$. Thus $\{\tau \wedge \theta \leq t\} = \{\tau \wedge \theta > t\}^c \in \mathcal{F}_t$.
- iii) $\{\tau \vee \theta \leq t\} = \{\max(\tau, \theta) \leq t\} = \{\tau \leq t\} \cap \{\theta \leq t\} \in \mathcal{F}_t$.

First passage time

Let $(B_t)_{t \geq 0}$ be a Brownian motion and let m be a real number.

First passage time

Let $(B_t)_{t \geq 0}$ be a Brownian motion and let m be a real number.

- Denote the first time $(B_t)_{t \geq 0}$ hits level m is
 $\tau_m = \inf\{t \geq 0 | B_t = m\}$: the first passage time.

First passage time

Let $(B_t)_{t \geq 0}$ be a Brownian motion and let m be a real number.

- Denote the first time $(B_t)_{t \geq 0}$ hits level m is $\tau_m = \inf\{t \geq 0 | B_t = m\}$: the first passage time.
- If $(B_t)_{t \geq 0}$ never reaches the level m , we set $\tau_m = \infty$.

First passage time

Let $(B_t)_{t \geq 0}$ be a Brownian motion and let m be a real number.

- Denote the first time $(B_t)_{t \geq 0}$ hits level m is $\tau_m = \inf\{t \geq 0 | B_t = m\}$: the first passage time.
- If $(B_t)_{t \geq 0}$ never reaches the level m , we set $\tau_m = \infty$.
- τ_m is a stopping time: $\{\tau_m \leq t\}$ is equivalent to $\max_{0 \leq s \leq t} B_s \leq m$, which is in \mathcal{F}_t .

Strong Markov property

Let $(B_t)_{t \geq 0}$ be a standard Brownian motion and let τ be a stopping time with respect to $(B_t)_{t \geq 0}$. Define $Z(t) = B_{t+\tau} - B_\tau$ then

Strong Markov property

Let $(B_t)_{t \geq 0}$ be a standard Brownian motion and let τ be a stopping time with respect to $(B_t)_{t \geq 0}$. Define $Z(t) = B_{t+\tau} - B_\tau$ then

- $(Z_t)_{t \geq 0}$ is also a standard Brownian motion.

Strong Markov property

Let $(B_t)_{t \geq 0}$ be a standard Brownian motion and let τ be a stopping time with respect to $(B_t)_{t \geq 0}$. Define $Z(t) = B_{t+\tau} - B_\tau$ then

- $(Z_t)_{t \geq 0}$ is also a standard Brownian motion.
- For each $t > 0$, $(Z_s)_{0 \leq s \leq t}$ is independent with $(B_u)_{0 \leq u \leq \tau}$.

Strong Markov property

Let $(B_t)_{t \geq 0}$ be a standard Brownian motion and let τ be a stopping time with respect to $(B_t)_{t \geq 0}$. Define $Z(t) = B_{t+\tau} - B_\tau$ then

- $(Z_t)_{t \geq 0}$ is also a standard Brownian motion.
- For each $t > 0$, $(Z_s)_{0 \leq s \leq t}$ is independent with $(B_u)_{0 \leq u \leq \tau}$.
- $(B_t)_{t \geq \tau}$ is a Brownian motion start with B_τ .

Example

Let $(B_t)_{t \geq 0}$ be a standard Brownian motion and let $\tau_m = \inf\{t \geq 0 | B_t = m\}$. Define $Z(t) = B_{t+\tau_m} - B_{\tau_m}$, with then

Example

Let $(B_t)_{t \geq 0}$ be a standard Brownian motion and let $\tau_m = \inf\{t \geq 0 | B_t = m\}$. Define $Z(t) = B_{t+\tau_m} - B_{\tau_m}$, with then

- $(Z_t)_{t \geq 0}$ is also a standard Brownian motion

Example

Let $(B_t)_{t \geq 0}$ be a standard Brownian motion and let $\tau_m = \inf\{t \geq 0 | B_t = m\}$. Define $Z(t) = B_{t+\tau_m} - B_{\tau_m}$, with then


- $(Z_t)_{t \geq 0}$ is also a standard Brownian motion
- For each $t > 0$, $(Z_s)_{0 \leq s \leq t}$ is independent with $(B_u)_{0 \leq u \leq \tau_m}$

Example

Let $(B_t)_{t \geq 0}$ be a standard Brownian motion and let $\tau_m = \inf\{t \geq 0 | B_t = m\}$. Define $Z(t) = B_{t+\tau_m} - B_{\tau_m}$, with then


- $(Z_t)_{t \geq 0}$ is also a standard Brownian motion
- For each $t > 0$, $(Z_s)_{0 \leq s \leq t}$ is independent with $(B_u)_{0 \leq u \leq \tau_m}$
- $(B_t)_{t \geq \tau_m}$ is a Brownian motion start with m .

Reflection principle of Brownian motion




Reflection principle of Brownian motion

- $P(\tau_m \leq t, B_t \geq m) = P(\tau_m \leq t, B_t \leq m)$




Reflection principle of Brownian motion

- $P(\tau_m \leq t, B_t \geq m) = P(\tau_m \leq t, B_t \leq m)$
- $P(\tau_m \leq t, B_t \leq w) = P(B_t \geq 2m - w).$



Reflection principle of Brownian motion

- $P(\tau_m \leq t, B_t \geq m) = P(\tau_m \leq t, B_t \leq m)$
- $P(\tau_m \leq t, B_t \leq w) = P(B_t \geq 2m - w).$
- $P(\tau_m \leq t, B_t \geq w) = P(B_t \leq 2m - w)$



First passage time distribution

Let $\tau_7 = \inf\{t \geq 0 | W_t = 7\}$. Prove that

First passage time distribution

Let $\tau_7 = \inf\{t \geq 0 | W_t = 7\}$. Prove that

- the random variable τ_7 has cumulative distribution function

$$P(\tau_7 \leq t) = \frac{2}{\sqrt{2\pi}} \int_{7/\sqrt{t}}^{\infty} e^{-y^2/2} dy$$

and the density function:

$$f_{\tau_7}(t) = \frac{d}{dt}P(\tau_7 \leq t) = \frac{7}{\sqrt{2\pi}t^3} e^{-49/(2t)}, t \geq 0.$$

First passage time distribution

Let $\tau_7 = \inf\{t \geq 0 | W_t = 7\}$. Prove that

- the random variable τ_7 has cumulative distribution function

$$P(\tau_7 \leq t) = \frac{2}{\sqrt{2\pi}} \int_{7/\sqrt{t}}^{\infty} e^{-y^2/2} dy$$

and the density function:

$$f_{\tau_7}(t) = \frac{d}{dt}P(\tau_7 \leq t) = \frac{7}{\sqrt{2\pi}t^3} e^{-49/(2t)}, t \geq 0.$$

- $P(\tau_7 < \infty) = 1$.

First passage time distribution: proof

First passage time distribution: proof

- We have $P(\tau_7 \leq t) = P(\tau_7 \leq t, B_t \leq 7) + P(\tau_7 \leq t, B_t \geq 7)$.

First passage time distribution: proof

- We have $P(\tau_7 \leq t) = P(\tau_7 \leq t, B_t \leq 7) + P(\tau_7 \leq t, B_t \geq 7)$.
- $P(\tau_7 \leq t, B_t \geq 7) = P(B_t \geq 7)$

First passage time distribution: proof

- We have $P(\tau_7 \leq t) = P(\tau_7 \leq t, B_t \leq 7) + P(\tau_7 \leq t, B_t \geq 7)$.
- $P(\tau_7 \leq t, B_t \geq 7) = P(B_t \geq 7)$
- $P(\tau_7 \leq t, B_t \leq 7) = P(B_t \geq 7)$.

First passage time distribution: proof

- We have $P(\tau_7 \leq t) = P(\tau_7 \leq t, B_t \leq 7) + P(\tau_7 \leq t, B_t \geq 7)$.
- $P(\tau_7 \leq t, B_t \geq 7) = P(B_t \geq 7)$
- $P(\tau_7 \leq t, B_t \leq 7) = P(B_t \geq 7)$.
- $P(\tau_7 \leq t) = 2P(B_t \geq 7) = 2 \int_7^\infty f_{B_t}(x) dx$

First passage time distribution: proof

- We have $P(\tau_7 \leq t) = P(\tau_7 \leq t, B_t \leq 7) + P(\tau_7 \leq t, B_t \geq 7)$.
- $P(\tau_7 \leq t, B_t \geq 7) = P(B_t \geq 7)$
- $P(\tau_7 \leq t, B_t \leq 7) = P(B_t \geq 7)$.
- $P(\tau_7 \leq t) = 2P(B_t \geq 7) = 2 \int_7^\infty f_{B_t}(x) dx$
- $f_{B_t}(x) = \frac{1}{\sqrt{2\pi t}} e^{-x^2/(2t)}.$

First passage time distribution: proof

- We have $P(\tau_7 \leq t) = P(\tau_7 \leq t, B_t \leq 7) + P(\tau_7 \leq t, B_t \geq 7)$.
- $P(\tau_7 \leq t, B_t \geq 7) = P(B_t \geq 7)$
- $P(\tau_7 \leq t, B_t \leq 7) = P(B_t \geq 7)$.
- $P(\tau_7 \leq t) = 2P(B_t \geq 7) = 2 \int_7^\infty f_{B_t}(x) dx$
- $f_{B_t}(x) = \frac{1}{\sqrt{2\pi t}} e^{-x^2/(2t)}$.
- $P(\tau_7 \leq t) = \frac{2}{\sqrt{2\pi t}} \int_7^\infty e^{-x^2/(2t)} dx =$
 $\frac{2}{\sqrt{2\pi}} \int_{7/\sqrt{t}}^\infty e^{-y^2/2} dy, \text{ with } y = x/\sqrt{t}.$

First passage time distribution

The density function: $f_{\tau_7}(t) = \frac{d}{dt}P(\tau_7 \leq t) = \frac{7}{\sqrt{2\pi}t^3}e^{-\frac{49}{2t}}, t \geq 0.$

$$P(\tau_7 < \infty) = \lim_{t \rightarrow \infty} \frac{2}{\sqrt{2\pi}} \int_{7/\sqrt{t}}^{\infty} e^{-y^2/2} dy = \frac{2}{\sqrt{2\pi}} \int_0^{\infty} e^{-y^2/2} dy = 1$$

First passage time distribution

Let $\tau_{-3} = \inf\{t \geq 0 | W_t = -3\}$. Find the cumulative distribution function and probability density function of τ_{-3} .

Example

A particle moves according to Brownian motion started at $x = 1$. After $t = 3$ hours, the particle is at level 1.5. Find the probability that the particle reaches level 2 sometime in the next hour.

Example

A particle moves according to Brownian motion started at $x = 1$. After $t = 3$ hours, the particle is at level 1.5. Find the probability that the particle reaches level 2 sometime in the next hour.

- After $t = 3$, the particle moves according to Brownian motion started at $x = 1.5$.

Example

A particle moves according to Brownian motion started at $x = 1$. After $t = 3$ hours, the particle is at level 1.5. Find the probability that the particle reaches level 2 sometime in the next hour.

- After $t = 3$, the particle moves according to Brownian motion started at $x = 1.5$.
- The probability of the event that the particle reaches level 2 sometime in the next hour (before $t = 4$) is identical with the probability of the event the standard Brownian motion first hits level $m = 2 - 1.5 = 0.5$ in the time interval $[0, 1]$.

Example

A particle moves according to Brownian motion started at $x = 1$. After $t = 3$ hours, the particle is at level 1.5. Find the probability that the particle reaches level 2 sometime in the next hour.

- After $t = 3$, the particle moves according to Brownian motion started at $x = 1.5$.
- The probability of the event that the particle reaches level 2 sometime in the next hour (before $t = 4$) is identical with the probability of the event the standard Brownian motion first hits level $m = 2 - 1.5 = 0.5$ in the time interval $[0, 1]$.
- The desired probability is

$$P(\tau_{0.5} \leq 1) = \int_0^1 \frac{0.5}{\sqrt{2\pi t^3}} e^{-0.5^2/(2t)} dt = 0.617.$$

Example

A particle moves according to Brownian motion started at $x = 2$. After $t = 2$ hours, the particle is at level 1. Find the probability that the particle reaches level -2 sometime in the next two hours.

First passage time

For all $m \neq 0$, the random variable τ_m has cumulative distribution function

$$P(\tau_m \leq t) = \frac{2}{\sqrt{2\pi}} \int_{|m|/\sqrt{t}}^{\infty} e^{-y^2/2} dy$$

and the density function:

$$f_{\tau_m}(t) = \frac{d}{dt} P(\tau_m \leq t) = \frac{|m|}{t\sqrt{2\pi t}} e^{-m^2/(2t)}, t \geq 0.$$

First passage time

The first passage time distribution has some surprising properties. Consider

$$\begin{aligned} P(\tau_m < \infty) &= \lim_{t \rightarrow \infty} P(\tau_m < t) = \lim_{t \rightarrow \infty} 2 \int_{|m|/\sqrt{t}}^{+\infty} \frac{e^{-x^2/2}}{\sqrt{2\pi}} dx \\ &= 2 \int_0^{+\infty} \frac{e^{-x^2/2}}{\sqrt{2\pi}} dx = 1. \end{aligned}$$

This means the Brownian motion will hit the level m in a finite period of time, with probability 1, for all m , no matter how large the value of m .

Stopped process

Given $(Z_t)_{t \geq 0}$ a stochastic process and τ is a stopping time. The stopped process $(Z_{t \wedge \tau})_{t \geq 0}$ is defined as

$$(Z_{t \wedge \tau})_{t \geq 0} = \begin{cases} Z_t & \text{if } t < \tau \\ Z_\tau & \text{if } t \geq \tau \end{cases} .$$

Optional stopping theorem

Assume that $(M_t)_{t \geq 0}$ is a martingale and τ is a stopping time, with respect to the filtration $(\mathcal{F}_t)_{t \geq 0}$. Then the stopped process $(M_{t \wedge \tau})_{t \geq 0}$ is also a martingale with respect to the filtration $(\mathcal{F}_t)_{t \geq 0}$.

Optional stopping theorem

Assume that $(M_t)_{t \geq 0}$ is a martingale and τ is a stopping time, with respect to the filtration $(\mathcal{F}_t)_{t \geq 0}$. Then the stopped process $(M_{t \wedge \tau})_{t \geq 0}$ is also a martingale with respect to the filtration $(\mathcal{F}_t)_{t \geq 0}$.

- If $P(\tau < \infty) = 1$ then

$$E[M_\tau] = E\left[\lim_{t \rightarrow \infty} M_{\tau \wedge t}\right] = \lim_{t \rightarrow \infty} E[M_{\tau \wedge t}] = E[M_0].$$

Exercise

Let $(W_t)_{t \geq 0}$ be a standard Brownian motion,
 $\tau_{-2} = \inf\{t \geq 0 \mid W_t = -2\}$, $\tau_3 = \inf\{t \geq 0 \mid W_t = 3\}$. Compute the
probability that $\tau_3 < \tau_{-2}$.

Exercise

Let $(W_t)_{t \geq 0}$ be a standard Brownian motion,
 $\tau_{-2} = \inf\{t \geq 0 \mid W_t = -2\}$, $\tau_3 = \inf\{t \geq 0 \mid W_t = 3\}$. Compute the probability that $\tau_3 < \tau_{-2}$.

- Let $\tau = \inf\{t \geq 0 \mid W_t \notin (-2, 3)\}$

Exercise

Let $(W_t)_{t \geq 0}$ be a standard Brownian motion,
 $\tau_{-2} = \inf\{t \geq 0 \mid W_t = -2\}$, $\tau_3 = \inf\{t \geq 0 \mid W_t = 3\}$. Compute the probability that $\tau_3 < \tau_{-2}$.

- Let $\tau = \inf\{t \geq 0 \mid W_t \notin (-2, 3)\}$
-

$$\begin{aligned}\{\tau \leq t\} &= \left\{ \max_{0 \leq u \leq t} W_u \geq 3 \right\} \cup \left\{ \min_{0 \leq u \leq t} W_u \leq -2 \right\} \\ &= \{\tau_3 \leq t\} \cup \{\tau_{-2} \leq t\} \in \mathcal{F}_t\end{aligned}$$

Exercise

Let $(W_t)_{t \geq 0}$ be a standard Brownian motion,
 $\tau_{-2} = \inf\{t \geq 0 \mid W_t = -2\}$, $\tau_3 = \inf\{t \geq 0 \mid W_t = 3\}$. Compute the probability that $\tau_3 < \tau_{-2}$.

- Let $\tau = \inf\{t \geq 0 \mid W_t \notin (-2, 3)\}$



$$\begin{aligned}\{\tau \leq t\} &= \left\{ \max_{0 \leq u \leq t} W_u \geq 3 \right\} \cup \left\{ \min_{0 \leq u \leq t} W_u \leq -2 \right\} \\ &= \{\tau_3 \leq t\} \cup \{\tau_{-2} \leq t\} \in \mathcal{F}_t\end{aligned}$$

- τ is a stopping time.

Exercise

Let $(W_t)_{t \geq 0}$ be a standard Brownian motion,
 $\tau_{-2} = \inf\{t \geq 0 \mid W_t = -2\}$, $\tau_3 = \inf\{t \geq 0 \mid W_t = 3\}$. Compute the probability that $\tau_3 < \tau_{-2}$.

- Let $\tau = \inf\{t \geq 0 \mid W_t \notin (-2, 3)\}$
-

$$\begin{aligned}\{\tau \leq t\} &= \{\max_{0 \leq u \leq t} W_u \geq 3\} \cup \{\min_{0 \leq u \leq t} W_u \leq -2\} \\ &= \{\tau_3 \leq t\} \cup \{\tau_{-2} \leq t\} \in \mathcal{F}_t\end{aligned}$$

- τ is a stopping time.
- $1 \geq P(\tau < \infty) \geq P(\tau_3 < \infty) = 1 \Rightarrow P(\tau < \infty) = 1.$

Exercise

Let $(W_t)_{t \geq 0}$ be a standard Brownian motion,
 $\tau_{-2} = \inf\{t \geq 0 \mid W_t = -2\}$, $\tau_3 = \inf\{t \geq 0 \mid W_t = 3\}$. Compute the probability that $\tau_3 < \tau_{-2}$.

- Let $\tau = \inf\{t \geq 0 \mid W_t \notin (-2, 3)\}$
-

$$\begin{aligned}\{\tau \leq t\} &= \left\{ \max_{0 \leq u \leq t} W_u \geq 3 \right\} \cup \left\{ \min_{0 \leq u \leq t} W_u \leq -2 \right\} \\ &= \{\tau_3 \leq t\} \cup \{\tau_{-2} \leq t\} \in \mathcal{F}_t\end{aligned}$$

- τ is a stopping time.
- $1 \geq P(\tau < \infty) \geq P(\tau_3 < \infty) = 1 \Rightarrow P(\tau < \infty) = 1.$
- $(W_t)_{t \geq 0}$ is a martingale

Exercise

- From the optional stopping theorem, $(W_{\tau \wedge t})_{t \geq 0}$ is a martingale and

$$\begin{aligned} E[W_\tau] &= E[\lim_{t \rightarrow \infty} W_{\tau \wedge t}] = \lim_{t \rightarrow \infty} E[W_{\tau \wedge t}] \\ &= E[W_{\tau \wedge 0}] = E[W_0] = 0 \end{aligned}$$

Exercise

- From the optional stopping theorem, $(W_{\tau \wedge t})_{t \geq 0}$ is a martingale and

$$\begin{aligned} E[W_\tau] &= E[\lim_{t \rightarrow \infty} W_{\tau \wedge t}] = \lim_{t \rightarrow \infty} E[W_{\tau \wedge t}] \\ &= E[W_{\tau \wedge 0}] = E[W_0] = 0 \end{aligned}$$

-

$$\begin{aligned} E[W_\tau] &= 3P(W_\tau = 3) - 2P(W_\tau = -2) \\ &= 3P(\tau_3 < \tau_{-2}) - 2P(\tau_3 > \tau_{-2}) \end{aligned}$$

Exercise

- From the optional stopping theorem, $(W_{\tau \wedge t})_{t \geq 0}$ is a martingale and

$$\begin{aligned} E[W_\tau] &= E[\lim_{t \rightarrow \infty} W_{\tau \wedge t}] = \lim_{t \rightarrow \infty} E[W_{\tau \wedge t}] \\ &= E[W_{\tau \wedge 0}] = E[W_0] = 0 \end{aligned}$$

•

$$\begin{aligned} E[W_\tau] &= 3P(W_\tau = 3) - 2P(W_\tau = -2) \\ &= 3P(\tau_3 < \tau_{-2}) - 2P(\tau_3 > \tau_{-2}) \end{aligned}$$

- $P(\tau_3 < \tau_{-2}) = \frac{2}{5}.$

Exercise

Let $(W_t)_{t \geq 0}$ be a standard Brownian motion,
 $\tau_{-7} = \inf\{t \geq 0 | W_t = -7\}$, $\tau_5 = \inf\{t \geq 0 | W_t = 5\}$. Compute the
probability that $\tau_5 > \tau_{-7}$.

Exercise

Let $(W_t)_{t \geq 0}$ be a Brownian motion. Compute the expectation of the first time $(W_t)_{t \geq 0}$ hits either the level (-2) or 3 .

Exercise

Let $(W_t)_{t \geq 0}$ be a Brownian motion. Compute the expectation of the first time $(W_t)_{t \geq 0}$ hits either the level (-2) or 3 .

- Let $\tau = \inf\{t \geq 0 | W_t \notin (-2, 3)\}$

Exercise

Let $(W_t)_{t \geq 0}$ be a Brownian motion. Compute the expectation of the first time $(W_t)_{t \geq 0}$ hits either the level (-2) or 3 .

- Let $\tau = \inf\{t \geq 0 | W_t \notin (-2, 3)\}$



$$\begin{aligned}\{\tau \leq t\} &= \{\max_{0 \leq u \leq t} W_u \geq 3\} \cup \{\min_{0 \leq u \leq t} W_u \leq -2\} \\ &= \{\tau_3 \leq t\} \cup \{\tau_{-2} \leq t\} \in \mathcal{F}_t\end{aligned}$$

Exercise

Let $(W_t)_{t \geq 0}$ be a Brownian motion. Compute the expectation of the first time $(W_t)_{t \geq 0}$ hits either the level (-2) or 3 .

- Let $\tau = \inf\{t \geq 0 | W_t \notin (-2, 3)\}$



$$\begin{aligned}\{\tau \leq t\} &= \{\max_{0 \leq u \leq t} W_u \geq 3\} \cup \{\min_{0 \leq u \leq t} W_u \leq -2\} \\ &= \{\tau_3 \leq t\} \cup \{\tau_{-2} \leq t\} \in \mathcal{F}_t\end{aligned}$$

- τ is a stopping time.

Exercise

Let $(W_t)_{t \geq 0}$ be a Brownian motion. Compute the expectation of the first time $(W_t)_{t \geq 0}$ hits either the level (-2) or 3 .

- Let $\tau = \inf\{t \geq 0 | W_t \notin (-2, 3)\}$



$$\begin{aligned}\{\tau \leq t\} &= \{\max_{0 \leq u \leq t} W_u \geq 3\} \cup \{\min_{0 \leq u \leq t} W_u \leq -2\} \\ &= \{\tau_3 \leq t\} \cup \{\tau_{-2} \leq t\} \in \mathcal{F}_t\end{aligned}$$

- τ is a stopping time.
- $1 \geq P(\tau < \infty) \geq P(\tau_3 < \infty) = 1 \Rightarrow P(\tau < \infty) = 1.$

Exercise

Let $(W_t)_{t \geq 0}$ be a Brownian motion. Compute the expectation of the first time $(W_t)_{t \geq 0}$ hits either the level (-2) or 3 .

- Let $\tau = \inf\{t \geq 0 | W_t \notin (-2, 3)\}$

-

$$\begin{aligned}\{\tau \leq t\} &= \{\max_{0 \leq u \leq t} W_u \geq 3\} \cup \{\min_{0 \leq u \leq t} W_u \leq -2\} \\ &= \{\tau_3 \leq t\} \cup \{\tau_{-2} \leq t\} \in \mathcal{F}_t\end{aligned}$$

- τ is a stopping time.
- $1 \geq P(\tau < \infty) \geq P(\tau_3 < \infty) = 1 \Rightarrow P(\tau < \infty) = 1.$
- $(W_t)_{t \geq 0}$ is a martingale

Exercise

- From the optional stopping theorem, $(W_{\tau \wedge t})_{t \geq 0}$ is a martingale and

$$\begin{aligned} E[W_\tau] &= E[\lim_{t \rightarrow \infty} W_{\tau \wedge t}] = \lim_{t \rightarrow \infty} E[W_{\tau \wedge t}] \\ &= E[W_{\tau \wedge 0}] = E[W_0] = 0 \end{aligned}$$

Exercise

- From the optional stopping theorem, $(W_{\tau \wedge t})_{t \geq 0}$ is a martingale and

$$\begin{aligned} E[W_\tau] &= E[\lim_{t \rightarrow \infty} W_{\tau \wedge t}] = \lim_{t \rightarrow \infty} E[W_{\tau \wedge t}] \\ &= E[W_{\tau \wedge 0}] = E[W_0] = 0 \end{aligned}$$



$$\begin{aligned} 0 &= E[W_\tau] = 3P(W_\tau = 3) - 2P(W_\tau = -2) \\ &= 3P(W_\tau = 3) - 2[1 - P(W_\tau = 3)] \end{aligned}$$

Exercise

- From the optional stopping theorem, $(W_{\tau \wedge t})_{t \geq 0}$ is a martingale and

$$\begin{aligned} E[W_\tau] &= E[\lim_{t \rightarrow \infty} W_{\tau \wedge t}] = \lim_{t \rightarrow \infty} E[W_{\tau \wedge t}] \\ &= E[W_{\tau \wedge 0}] = E[W_0] = 0 \end{aligned}$$

•

$$\begin{aligned} 0 &= E[W_\tau] = 3P(W_\tau = 3) - 2P(W_\tau = -2) \\ &= 3P(W_\tau = 3) - 2[1 - P(W_\tau = 3)] \end{aligned}$$

- $P(W_\tau = 3) = \frac{2}{5}, P(W_\tau = -2) = \frac{3}{5}.$

Exercise

- $(W_t^2 - t)_{t \geq 0}$ is a martingale

Exercise

- $(W_t^2 - t)_{t \geq 0}$ is a martingale
- From the optional stopping theorem, $(W_{\tau \wedge t}^2 - \tau \wedge t)_{t \geq 0}$ is a martingale and $E[W_\tau^2 - \tau] = E[\lim_{t \rightarrow \infty} W_{\tau \wedge t}^2 - \tau \wedge t] = \lim_{t \rightarrow \infty} E[W_{\tau \wedge t}^2 - \tau \wedge t] = E[W_{\tau \wedge 0}] = E[W_0^2] = 0$.

Exercise

- $(W_t^2 - t)_{t \geq 0}$ is a martingale
- From the optional stopping theorem, $(W_{\tau \wedge t}^2 - \tau \wedge t)_{t \geq 0}$ is a martingale and $E[W_\tau^2 - \tau] = E[\lim_{t \rightarrow \infty} W_{\tau \wedge t}^2 - \tau \wedge t] = \lim_{t \rightarrow \infty} E[W_{\tau \wedge t}^2 - \tau \wedge t] = E[W_{\tau \wedge 0}] = E[W_0^2] = 0$.
-

$$\begin{aligned} E[\tau] &= E[W_\tau^2] = 3^2 P(W_\tau = 3) + (-2)^2 P(W_\tau = -2) \\ &= 9 \frac{2}{5} + 4 \frac{3}{5} = 6. \end{aligned}$$

Maximum of Brownian motion

- Let (B_t) be the standard Brownian motion. Then $(M_t)_{t \geq 0}$, where $M_t = \max_{0 \leq s \leq t} B_s$, forms the maximum process of Brownian motion

Maximum of Brownian motion

- Let (B_t) be the standard Brownian motion. Then $(M_t)_{t \geq 0}$, where $M_t = \max_{0 \leq s \leq t} B_s$, forms the maximum process of Brownian motion
- M_t is the maximum value of Brownian motion on $[0, t]$.

Maximum of Brownian motion

- Let (B_t) be the standard Brownian motion. Then $(M_t)_{t \geq 0}$, where $M_t = \max_{0 \leq s \leq t} B_s$, forms the maximum process of Brownian motion
- M_t is the maximum value of Brownian motion on $[0, t]$.
- $M_t > a$ if and only if $\tau_a < t$.

Maximum of Brownian motion

- Let (B_t) be the standard Brownian motion. Then $(M_t)_{t \geq 0}$, where $M_t = \max_{0 \leq s \leq t} B_s$, forms the maximum process of Brownian motion
- M_t is the maximum value of Brownian motion on $[0, t]$.
- $M_t > a$ if and only if $\tau_a < t$.
- $P(M_t > a) = P(\tau_a < t) = \int_0^t \frac{a}{\sqrt{2\pi s^3}} e^{-a^2/(2s)} ds =$
 $\int_a^{+\infty} \sqrt{\frac{2}{\pi t}} e^{-x^2/(2t)} dx, \quad a^2/s = x^2/t.$

Example

A laboratory instrument takes annual temperature measurements. Measurement errors are assumed to be independent and normally distributed. As precision decreases over time, errors are modeled as standard Brownian motion. For how many years can the lab be guaranteed that there is at least 90% probability that all errors are less than 4 degrees, given that the 95th percentile of the standard normal random distribution is 1.645?

Example

A laboratory instrument takes annual temperature measurements. Measurement errors are assumed to be independent and normally distributed. As precision decreases over time, errors are modeled as standard Brownian motion. For how many years can the lab be guaranteed that there is at least 90% probability that all errors are less than 4 degrees, given that the 95th percentile of the standard normal random distribution is 1.645?

- Find the largest t such that $P(M_t \leq 4) \geq 0.9$.

Example

A laboratory instrument takes annual temperature measurements. Measurement errors are assumed to be independent and normally distributed. As precision decreases over time, errors are modeled as standard Brownian motion. For how many years can the lab be guaranteed that there is at least 90% probability that all errors are less than 4 degrees, given that the 95th percentile of the standard normal random distribution is 1.645?

- Find the largest t such that $P(M_t \leq 4) \geq 0.9$.
- $0.9 \leq P(M_t \leq 4) = 1 - P(M_t > 4) = 1 - 2P(B_t > 4) = 2P(B_t \leq 4) - 1,$

Example

A laboratory instrument takes annual temperature measurements. Measurement errors are assumed to be independent and normally distributed. As precision decreases over time, errors are modeled as standard Brownian motion. For how many years can the lab be guaranteed that there is at least 90% probability that all errors are less than 4 degrees, given that the 95th percentile of the standard normal random distribution is 1.645?

- Find the largest t such that $P(M_t \leq 4) \geq 0.9$.
- $0.9 \leq P(M_t \leq 4) = 1 - P(M_t > 4) = 1 - 2P(B_t > 4) = 2P(B_t \leq 4) - 1$,
- $0.95 \leq P(B_t \leq 4) = P(\sqrt{t}Z \leq 4) = P(Z \leq \frac{4}{\sqrt{t}})$, where $Z \sim N(0, 1)$.

Example

A laboratory instrument takes annual temperature measurements. Measurement errors are assumed to be independent and normally distributed. As precision decreases over time, errors are modeled as standard Brownian motion. For how many years can the lab be guaranteed that there is at least 90% probability that all errors are less than 4 degrees, given that the 95th percentile of the standard normal random distribution is 1.645?

- Find the largest t such that $P(M_t \leq 4) \geq 0.9$.
- $0.9 \leq P(M_t \leq 4) = 1 - P(M_t > 4) = 1 - 2P(B_t > 4) = 2P(B_t \leq 4) - 1$,
- $0.95 \leq P(B_t \leq 4) = P(\sqrt{t}Z \leq 4) = P(Z \leq \frac{4}{\sqrt{t}})$, where $Z \sim N(0, 1)$.
- The 95th percentile of the standard normal random distribution is 1.645. Thus the desired value t should satisfy $4/\sqrt{t} \geq 1.645 \Leftrightarrow t \leq 5.91$ years.

Example

A laboratory instrument takes annual temperature measurements. Measurement errors are assumed to be independent and normally distributed. As precision decreases over time, errors are modeled as standard Brownian motion. For how many years can the lab be guaranteed that there is at least 80% probability that all errors are less than 5 degrees, given that the 90th percentile of the standard normal random distribution is 1.28?

Zeros of Brownian motion

- On any interval $(0, \epsilon)$, no matter how small ϵ , the process crosses the t -axis infinitely many times.

Zeros of Brownian motion

- On any interval $(0, \epsilon)$, no matter how small ϵ , the process crosses the t -axis infinitely many times.
- The times when the process crosses the t -axis are the zeros of Brownian motion.

Zeros of Brownian motion

- On any interval $(0, \epsilon)$, no matter how small ϵ , the process crosses the t -axis infinitely many times.
- The times when the process crosses the t -axis are the zeros of Brownian motion.
- For $0 \leq r < t$, let $z_{r,t}$ be the probability that standard Brownian motion has at least one zero in (r, t) . Then

$$z_{r,t} = \frac{2}{\pi} \arccos\left(\sqrt{\frac{r}{t}}\right).$$

Zeros of Brownian motion

- $z_{r,t} = P(B_s = 0), \text{ for some } s \in (r, t) = \int_{-\infty}^{+\infty} P(B_s = 0 | B_r = x) \frac{1}{\sqrt{2\pi r}} e^{-x^2/(2r)} dx, \text{ for some } s \in (r, t)$

Zeros of Brownian motion

- $z_{r,t} = P(B_s = 0), \text{ for some } s \in (r, t) = \int_{-\infty}^{+\infty} P(B_s = 0 | B_r = x) \frac{1}{\sqrt{2\pi r}} e^{-x^2/(2r)} dx, \text{ for some } s \in (r, t)$
- For $x < 0$, $P(B_s = 0 | B_r = x) = P(M_t \geq 0 | B_r = x) = P(M_t \geq -x | B_r = 0) = P(M_{t-r} > -x | B_0 = 0) = P(M_{t-r} > -x)$.

Zeros of Brownian motion

- $z_{r,t} = P(B_s = 0), \text{ for some } s \in (r, t) = \int_{-\infty}^{+\infty} P(B_s = 0 | B_r = x) \frac{1}{\sqrt{2\pi r}} e^{-x^2/(2r)} dx, \text{ for some } s \in (r, t)$
- For $x < 0$, $P(B_s = 0 | B_r = x) = P(M_t \geq 0 | B_r = x) = P(M_t \geq -x | B_r = 0) = P(M_{t-r} > -x | B_0 = 0) = P(M_{t-r} > -x)$.
- For $x \geq 0$, we consider the reflected Brownian motion $P(B_s = 0 | B_r = x) = P(B_s = 0 | B_r = -x) = P(M_t \geq 0 | B_r = -x) = P(M_t \geq x | B_r = 0) = P(M_{t-r} > x)$. Thus we can express $P(B_s = 0 | B_r = x) = P(M_{t-r} > |x|)$

Zeros of Brownian motion

- $z_{r,t} = P(B_s = 0), \text{ for some } s \in (r, t) = \int_{-\infty}^{+\infty} P(B_s = 0 | B_r = x) \frac{1}{\sqrt{2\pi r}} e^{-x^2/(2r)} dx, \text{ for some } s \in (r, t)$
- For $x < 0$, $P(B_s = 0 | B_r = x) = P(M_t \geq 0 | B_r = x) = P(M_t \geq -x | B_r = 0) = P(M_{t-r} > -x | B_0 = 0) = P(M_{t-r} > -x)$.
- For $x \geq 0$, we consider the reflected Brownian motion $P(B_s = 0 | B_r = x) = P(B_s = 0 | B_r = -x) = P(M_t \geq 0 | B_r = -x) = P(M_t \geq x | B_r = 0) = P(M_{t-r} > x)$. Thus we can express $P(B_s = 0 | B_r = x) = P(M_{t-r} > |x|)$
- $z_{r,t} = \int_{-\infty}^{+\infty} P(M_{t-r} > |x|) \frac{1}{\sqrt{2\pi r}} e^{-x^2/(2r)} dx = \frac{2}{\pi} \arccos(\sqrt{\frac{r}{t}})$.

Zeros of Brownian motion

- $z_{0,\epsilon} = 2/\pi \arccos(0) = 1$. That is $B_t = 0$ for some $0 < t < \epsilon$.

Zeros of Brownian motion

- $z_{0,\epsilon} = 2/\pi \arccos(0) = 1$. That is $B_t = 0$ for some $0 < t < \epsilon$.
- By the strong Markov property, for Brownian motion restarted at t , there is at least one zero in (t, ϵ) , with probability 1.

Zeros of Brownian motion

- $z_{0,\epsilon} = 2/\pi \arccos(0) = 1$. That is $B_t = 0$ for some $0 < t < \epsilon$.
- By the strong Markov property, for Brownian motion restarted at t , there is at least one zero in (t, ϵ) , with probability 1.
- Continuing this way, there are infinitely many zeros in $(0, \epsilon)$.

Brownian motion with drift

- For real value μ and $\sigma > 0$, the process defined by $W_t = \mu t + \sigma B_t$, for $t \geq 0$, is called Brownian motion with drift parameter μ and variance parameter σ^2 .

Brownian motion with drift

- For real value μ and $\sigma > 0$, the process defined by $W_t = \mu t + \sigma B_t$, for $t \geq 0$, is called Brownian motion with drift parameter μ and variance parameter σ^2 .
- For $s, t > 0$, $W_{t+s} - W_t \sim N(\mu s, \sigma^2 s)$.

Brownian motion with drift

- For real value μ and $\sigma > 0$, the process defined by $W_t = \mu t + \sigma B_t$, for $t \geq 0$, is called Brownian motion with drift parameter μ and variance parameter σ^2 .
- For $s, t > 0$, $W_{t+s} - W_t \sim N(\mu s, \sigma^2 s)$.
-

$$\begin{aligned} P(W_t - W_s \leq k | \mathcal{F}_s) &= P(\mu(t-s) + \sigma(B_t - B_s) \leq k | \mathcal{F}_s) \\ &= P(B_t - B_s \leq \frac{k - \mu(t-s)}{\sigma} | \mathcal{F}_s) = P(B_t - B_s \leq \frac{k - \mu(t-s)}{\sigma}) \\ &= P(\mu(t-s) + \sigma(B_t - B_s) \leq k) = P(W_t - W_s \leq k). \end{aligned}$$

Brownian motion with drift

- For real value μ and $\sigma > 0$, the process defined by $W_t = \mu t + \sigma B_t$, for $t \geq 0$, is called Brownian motion with drift parameter μ and variance parameter σ^2 .
- For $s, t > 0$, $W_{t+s} - W_t \sim N(\mu s, \sigma^2 s)$.
-

$$\begin{aligned} P(W_t - W_s \leq k | \mathcal{F}_s) &= P(\mu(t-s) + \sigma(B_t - B_s) \leq k | \mathcal{F}_s) \\ &= P(B_t - B_s \leq \frac{k - \mu(t-s)}{\sigma} | \mathcal{F}_s) = P(B_t - B_s \leq \frac{k - \mu(t-s)}{\sigma}) \\ &= P(\mu(t-s) + \sigma(B_t - B_s) \leq k) = P(W_t - W_s \leq k). \end{aligned}$$

•

$$\begin{aligned} W_t - W_s &= \mu(t-s) + \sigma(B_t - B_s) \\ &\stackrel{d}{=} \mu(t-s) + \sigma B_{t-s} \stackrel{d}{=} W_{t-s}, \quad \forall 0 < s < t. \end{aligned}$$

Model Brownian motion with drift

Suppose we want to model a Brownian motion with drift:

$dW_t = \mu dt + \sigma dB_t$ in a time interval $[0, T]$, where $(B_t)_{t \geq 0}$ be a standard Brownian motion.

Model Brownian motion with drift

Suppose we want to model a Brownian motion with drift:

$dW_t = \mu dt + \sigma dB_t$ in a time interval $[0, T]$, where $(B_t)_{t \geq 0}$ be a standard Brownian motion.

- We divide the interval into n equal subintervals by discrete time points $0 = t_0 < t_1 < \dots < t_n = T$, with the time step $h = t_i - t_{i-1}$.

Model Brownian motion with drift

Suppose we want to model a Brownian motion with drift:

$dW_t = \mu dt + \sigma dB_t$ in a time interval $[0, T]$, where $(B_t)_{t \geq 0}$ be a standard Brownian motion.

- We divide the interval into n equal subintervals by discrete time points $0 = t_0 < t_1 < \dots < t_n = T$, with the time step $h = t_i - t_{i-1}$.
- The value of W_{t_i} is determined by a recursive formula:

$$W_{t_i} = W_{t_{i-1}} + \mu * h + \sigma \sqrt{h} * X,$$

where $X \sim N(0, 1)$.

Model Brownian motion with drift

Suppose we want to model a Brownian motion with drift:

$dW_t = \mu dt + \sigma dB_t$ in a time interval $[0, T]$, where $(B_t)_{t \geq 0}$ be a standard Brownian motion.

- We divide the interval into n equal subintervals by discrete time points $0 = t_0 < t_1 < \dots < t_n = T$, with the time step $h = t_i - t_{i-1}$.
- The value of W_{t_i} is determined by a recursive formula:

$$W_{t_i} = W_{t_{i-1}} + \mu * h + \sigma \sqrt{h} * X,$$

where $X \sim N(0, 1)$.


- All we need to simulate a generalized Brownian motion is to simulate a standard normal random variable, which can be done by using command “`rnorm(1)`” in software R.

R code for simulating Brownian motion with drift

```
BM=function(mu,sigma,T,N) {  
  # mu: the drift term of Brownian motion  
  # sigma: the diffusion term of Brownian motion  
  # T: expiry time  
  # N: number of simulation points  
  h=T/N # the timestep of the simulation  
  X=rep(0, (N+1)) # create space to store values of Brownian  
    motion  
  X[1]=0  
  for(i in 1:N) { X[i+1]=X[i] +mu*h+sigma*sqrt(h)*rnorm(1)}  
  return(X)  
}
```

A sample of standard Brownian motion


```
h <- 1/1000  
t=seq(0,1,by=h)  
plot(t,BM(mu=0.05,sigma=0.3,T=1,N=1000), type="l",  
      col="blue",xlab="t",ylab="B")
```



Plot multiple samples of Brownian motion with drift

```
BMSamplepaths <- function(mu,sigma,T,N,nt)
{h=T/N ## time step
# number of discrete points
#nt: number of samples
t=seq(0,T,by=h) # produce a sequence of time points
X=matrix(rep(0,length(t)*nt), nrow=nt)
## each row of the matrix stores one sample
# #return(X)
for (i in 1:nt) {X[i,]= BM(mu=mu,sigma=sigma,T=T,N=N)}
# # ##Plot
ymax=max(X); ymin=min(X) #bounds for simulated prices
plot(t,X[1,],t='l',main='1000 sample paths of standard
      Brownian motions
      with parameters mu=0.05, sigma=0.3, T =1, N=1000, nt=1000')
}
```

**1000 sample paths of standard Brownian motions
with parameters $\mu=0.05$, $\sigma=0.3$, $T=1$, $N=200$**



Example

Find the probability that Brownian motion with drift parameter $\mu = 0.6$ and variance $\sigma^2 = 0.25$ takes values between 1 and 3 at time $t = 4$.

Example

Find the probability that Brownian motion with drift parameter $\mu = 0.6$ and variance $\sigma^2 = 0.25$ takes values between 1 and 3 at time $t = 4$.

- Brownian motion with drift parameter $\mu = 0.6$ and variance $\sigma^2 = 0.25$ is $W_t = 0.6t + 0.5B_t$.

Example

Find the probability that Brownian motion with drift parameter $\mu = 0.6$ and variance $\sigma^2 = 0.25$ takes values between 1 and 3 at time $t = 4$.

- Brownian motion with drift parameter $\mu = 0.6$ and variance $\sigma^2 = 0.25$ is $W_t = 0.6t + 0.5B_t$.
- $P(1 \leq W_4 \leq 3) = P(1 \leq 0.6 * 4 + 0.5 * B_4 \leq 3) = P(-2.8 \leq B_4 \leq 1.2) = \int_{-2.8}^{1.2} \frac{e^{-x^2/8}}{\sqrt{8\pi}} dx = 0.645$.

Brownian bridge

- Brownian bridge is a Brownian motion starting at x at time t_0 and passing through some point y at time $T > t_0$.

Brownian bridge

- Brownian bridge is a Brownian motion starting at x at time t_0 and passing through some point y at time $T > t_0$.
- $W_{t_0, x}^{T, y}(t) = x + W_{t-t_0} - \frac{t-t_0}{T-t_0}(W_{T-t_0} - y + x).$

Exercise: Simulating Brownian bridge

Write a R code to simulate the Brownian motion

