


1 The worst case

Here should be definition of the group G^


Definition 1.1. Function $size : G \rightarrow \mathbb{N} \cup \{0\}$ is defined recursively:

$$size(w = \pi(w_1, w_2, w_3)) = \begin{cases} 0, & w = e \\ 0, & w = \delta \text{ (reverse node)} \\ |w| + size(w_1) + size(w_2) + size(w_3), & \text{otherwise} \end{cases} \quad (1)$$

Thus, $size(a) = size(b) = size(c) = 1$ by definition.



Similarly, $size(aa) = size(bb) = size(cc) = 2$, $size(ab) = size(ac) = \dots = size(bc) = 4$



Definition 1.2. abc -subset $X \subset \{a, b, c\}^*$ is such subset:

$$X = \{(a^\pi b^\pi c^\pi)^k, (a^\pi b^\pi c^\pi)^k a^\pi, (a^\pi b^\pi c^\pi)^k a^\pi b^\pi \mid k \in \mathbb{N} \cup \{0\}, \pi \in S_3\} \quad (2)$$

where a^π means S_3 group action on set $\{a, b, c\}$;

$(w_1 w_2 w_3)^k$ means repeating k times of the expression in the parenthesis.


Lemma 1.1. $\forall w \in X$ w will have one of the next possible structures:

$$\pi(w_1, w_2, e), \quad \pi(w_1, e, w_2), \quad \pi(e, w_1, w_2),$$

where $w_1, w_2 \in X$, $|w_1|, |w_2| \in \{\lfloor \frac{n}{2} \rfloor, \lceil \frac{n}{2} \rceil\}$, $|w_1| + |w_2| = n$, $n = |w|$

Proof. Follows from definition of multiplication. □

Example:



Therefore, here we have recursive formula for calculating function *size* (1) for elements from X :

$$\text{size}(w \in X) = a(n) = a\left(\left\lfloor \frac{n}{2} \right\rfloor\right) + a\left(\left\lceil \frac{n}{2} \right\rceil\right) + n, \quad a(1) = 1, \quad a(0) = 0, \quad n = |w|$$

Statement 1.1. *Exact form of function $a(n)$:*

$$a(n) = \begin{cases} n(\lfloor \log_2 n + 1 \rfloor) + 2n - 2^{\lfloor \log_2 n \rfloor + 1}, & n > 0 \\ a(0) = 0 \end{cases} \quad (3)$$

Proof.

$$a(n+1) = n+1 + a\left(\left\lfloor \frac{n+1}{2} \right\rfloor\right) + a\left(\left\lceil \frac{n+1}{2} \right\rceil\right) = n+1 + a\left(\left\lfloor \frac{n}{2} \right\rfloor + 1\right) + a\left(\left\lceil \frac{n}{2} \right\rceil\right)$$

$$\text{Let } b(n) := a(n+1) - a(n) = 1 + a\left(\left\lfloor \frac{n}{2} \right\rfloor + 1\right) - a\left(\left\lfloor \frac{n}{2} \right\rfloor\right) \text{ - auxiliary recursion.}$$

$$b(n) = b\left(\left\lfloor \frac{n}{2} \right\rfloor\right) + 1, \quad b(2) = 4 \quad \Rightarrow \quad b(n) = \lfloor \log_2 n \rfloor + 3$$

$$\begin{aligned} a(n+1) &= b(n) + b(n-1) + \dots + b(1) = \sum_{1 \leq k \leq n} (\lfloor \log_2 k + 3 \rfloor) \quad \Rightarrow \quad a(n) = 2(n-1) + \sum_{1 \leq k \leq n} (\lfloor \log_2 k + 1 \rfloor) = \\ &= \left[\begin{array}{l} \text{amount of bits in all the} \\ \text{numbers from 1 to } N-1 \end{array} \right] = 2(n-1) + \sum_{i=1}^{\lfloor \log_2 n \rfloor} (n - 2^i) = n(\lfloor \log_2 n + 1 \rfloor) + 2n - 2^{\lfloor \log_2 n \rfloor + 1} \end{aligned}$$

□

Theorem 1.2. *Elements from abc-subset X have maximum size, or in other words*

$$a(n) = \max(\text{size}(w) \mid w \in \{a, b, c\}^n)$$

Proof. Induction on n .

1. Base: check examples of size calculation.

2. Induction step:

Let $\forall k < n \quad a(k) = \max(\text{size}(w) \mid w \in \{a, b, c\}^k)$. Hereafter we consider any $w \in \{a, b, c\}^n$, $w = \pi(w_1, w_2, w_3)$, $w_1, w_2, w_3 \in \{a, b, c\}^* \cup \{e\}$. Now we need to show that $\text{size}(w) \leq a(n)$

Let $x_1 := |w_1|$, $x_2 := |w_2|$, $x_3 := |w_3|$. If w has maximum size than due to induction hypothesis

$$\text{size}(w) \leq a(x_1) + a(x_2) + a(x_3) + n \quad (4)$$

(we don't know whether any w_1, w_2, w_3 from X could appear in w , so we use \leq symbol). Let's investigate how big this functional could be.

Consider function $f(x) = x \log_2 x$ instead of $a(x)$. We can do it because both $f(x)$ and $a(x)$ are monotonically increasing on $x \geq 1$. Thus, they are maximizing in similar way.

Now we need to solve optimization problem

$$\begin{aligned} &\max(f(x_1) + f(x_2) + f(x_3) \mid x_1 > 1, x_2 > 1, x_3 > 1, x_1 + x_2 + x_3 = n) \\ F &= f(x_1) + f(x_2) + f(x_3) = x_1 \log_2 x_1 + x_2 \log_2 x_2 + x_3 \log_2 x_3 = \left[x_3 = n - x_1 - x_2 \right] = \\ &= x_1 \log_2 x_1 + x_2 \log_2 x_2 + (n - x_1 - x_2) \log_2 (n - x_1 - x_2) = \\ &= x_1 \log_2 x_1 + x_2 \log_2 x_2 + n \log_2 (n - x_1 - x_2) - x_1 \log_2 (n - x_1 - x_2) - x_2 \log_2 (n - x_1 - x_2) \\ \frac{\partial F}{\partial x_1} &= \log_2 x_1 + \frac{1}{\ln 2} - \frac{n}{(n - x_1 - x_2) \ln 2} - \log_2 (n - x_1 - x_2) + \frac{x_1}{(n - x_1 - x_2) \ln 2} + \frac{x_2}{(n - x_1 - x_2) \ln 2} \end{aligned}$$

$$\frac{\partial F}{\partial x_2} = \log_2 x_2 + \frac{1}{\ln 2} - \frac{n}{(n - x_1 - x_2) \ln 2} - \log_2(n - x_1 - x_2) + \frac{x_1}{(n - x_1 - x_2) \ln 2} + \frac{x_2}{(n - x_1 - x_2) \ln 2}$$

If both $\frac{\partial F}{\partial x_1}$ and $\frac{\partial F}{\partial x_2}$ equal to zero then $\log_2 x_1 = \log_2 x_2 \Rightarrow x_1$ should be equal to x_2 .

Let's substitute $x = x_1 = x_2$ to F and differentiate it:

$$\frac{dF}{dx} = 2 \log_2 x + \frac{2}{\ln 2} - \frac{2n}{(n - 2x) \ln 2} - 2 \log_2(n - 2x) + \frac{4x}{(n - 2x) \ln 2} = 0$$

Hence, $x = \frac{n}{3}$ - the single solution (Wolfram). Substituting it to F we can find out that $x_1 = x_2 = x_3 = \frac{n}{3}$ - global minimum. Thus, F (therefore right part of (4)) reaches its maximum value on the bounds of its domain.

Now we are allowed to assume that $\exists i w_i = e$, because we know that otherwise

$$size(w) \leq a(x_1) + a(x_2) + a(x_3) + n \leq a\left(\left\lfloor \frac{n}{2} \right\rfloor\right) + a\left(\left\lceil \frac{n}{2} \right\rceil\right) + n$$

It's easy to check that there are only 2 possible types of w with one or more trivial element below: familiar to us $w \in X$ or such $w = z_1 z_2 z_3 \dots z_n$ where there are at least one i such that $z_i = z_{i+1}$, $\Rightarrow w$ could be reduced $\Rightarrow size(w) < a(n)$ due to definition.

□

2 Generative function

$$\begin{aligned} T(z) &= \sum_w size(w) z^{|w|} = \sum_{w_1} \sum_{w_2} \sum_{w_3} (s(w_1) + s(w_2) + s(w_3) + |w_1| + |w_2| + |w_3|) z^{|w_1| + |w_2| + |w_3|} = \\ &= \sum_{w_1} \sum_{w_2} z^{|w_1| + |w_2|} \left((s(w_1) + s(w_2)) \sum_{w_3} z^{|w_3|} + \sum_{w_3} s(w_3) z^{|w_3|} + (|w_1| + |w_2|) \sum_{w_3} z^{|w_3|} + \sum_{w_3} |w_3| z^{|w_3|} \right) = \\ &= \left[\sum_w z^{|w|} = F(z), \quad \sum_w |w| z^{|w|} = z \sum_w |w| z^{|w|-1} = z F'(z) \right] = \\ &= \sum_{w_1} z^{|w_1|} \sum_{w_2} z^{|w_2|} ((s(w_1) + s(w_2) + |w_1| + |w_2|) F(z) + T(z) + z F'(z)) = \\ &= \sum_{w_1} z^{|w_1|} \left((T(z) + z F'(z)) \sum_{w_2} z^{|w_2|} + F(z) \left((s(w_1) + |w_1|) \sum_{w_2} z^{|w_2|} + \sum_{w_2} s(w_2) z^{|w_2|} + \sum_{w_2} |w_2| z^{|w_2|} \right) \right) = \\ &= \sum_{w_1} z^{|w_1|} [(T(z) + z F'(z)) F(z) + F(z) ((s(w_1) + |w_1|) F(z) + T(z) + z F'(z))] = \\ &= \sum_{w_1} z^{|w_1|} [T(z) F(z) + z F(z) F'(z) + F^2(z) (s(w_1) + |w_1|) + F(z) T(z) + z F(z) F'(z)] = \\ &= \sum_{w_1} z^{|w_1|} [2T(z) F(z) + 2z F(z) F'(z) + F^2(z) (s(w_1) + |w_1|)] = \\ &= 2T(z) F^2(z) + 2z F^2(z) F'(z) + F^2(z) T(z) + z F^2(z) F'(z) = 3T(z) F^2(z) + 3z F^2(z) F'(z) \\ &\Rightarrow T(z) = \frac{3z F^2(z) F'(z)}{1 - 3F^2(z)}, \quad F(z) = \sum_w z^{|w|} = \sum_{n=1}^{\infty} x_n z^n \end{aligned}$$

1. $x_n = 3^n$

$$F(z) = \sum_{n=1}^{\infty} 3^n z^n = \sum_{n=1}^{\infty} (3z)^n = \frac{1}{1 - 3z} \Rightarrow T(z) = \frac{3z \frac{1}{(1 - 3z)^2} \frac{3}{(1 - 3z)^2}}{1 - 3 \frac{1}{(1 - 3z)^2}} = \frac{9z}{81z^4 - 108z^3 + 27z^2 + 6z - 2} = \frac{P(z)}{Q(z)}$$

Roots of $Q(z)$: $\frac{1}{3}, \frac{1}{3}, \frac{1}{3} - \frac{1}{\sqrt{3}}, \frac{1}{3} + \frac{1}{\sqrt{3}}$

The root $\frac{1}{3} - \frac{1}{\sqrt{3}}$ has the least absolute value (~ 0.24). Thus, by theorem of asymptotic growth of the rational generative functions' coefficients

$$[z^n]T(z) \sim C \left(\frac{3\sqrt{3}}{\sqrt{3}-3} \right)^n \Rightarrow \frac{[z^n]T(z)}{3^n} \sim C \left(\frac{\sqrt{3}}{\sqrt{3}-3} \right)^n$$

2. $x_n = 3 * 2^{n-1}$

$$\begin{aligned} F(z) &= \sum_{n=1}^{\infty} 3 * 2^{n-1} z^n = \frac{3}{2} \sum_{n=1}^{\infty} (2z)^n = \frac{3}{2-4z}, \Rightarrow T(z) = \frac{3z \left(\frac{3}{2-4z} \right)^2 \frac{d}{dz} \left(\frac{3}{2-4z} \right)}{1 - 3 \left(\frac{3}{2-4z} \right)^2} = \\ &= \frac{81z}{64z^4 - 128z^3 - 12z^2 + 76z - 23} = \frac{P(z)}{Q(z)} \end{aligned}$$

Roots of $Q(z)$: $\frac{1}{2}, \frac{1}{2}, \frac{1}{2} - \frac{3\sqrt{3}}{4}, \frac{1}{2} + \frac{3\sqrt{3}}{4}$

The root $\frac{1}{2}$ of multiplicity 2 has the least absolute value. Thus, by theorem of asymptotic growth of the rational generative functions' coefficients

$$\begin{aligned} [z^n]T(z) &\sim C(2)^n n, \quad C = 2 \frac{(-2)^2 * P(1/2)}{Q''(1/2)} = \frac{8 * 81/2}{24(32 * 1/4 - 32 * 1/2 - 1)} = -\frac{81}{42} \\ \Rightarrow \frac{[z^n]T(z)}{3 * 2^{n-1}} &\sim \frac{-81 * 2^n * n}{42 * 3 * 2^{n-1}} = -\frac{9}{7}n \end{aligned}$$