

# Introduction to equations with matrices

Math 352 Differential Equations

April 14, 2014

# The machinery of linear algebra

- ▶ Fundamental algebraic construction of linear algebra: *linear combinations*.
- ▶ *Vector spaces* are places where it makes sense to form linear combos.
- ▶ Coefficients can be real, complex, ...
- ▶ The objects being combined are often called vectors, even if they are not vectors like  $\langle 2, -3, 4, 12 \rangle$  or  $2\vec{i} - 6\vec{j}$ .
- ▶ Suppose that  $y_1, y_2, y_3, \dots, y_n$  are differentiable and that  $a_1, \dots, a_n$  are real numbers (scalars).
- ▶ Then  $a_1y_1 + a_2y_2 + \dots + a_ny_n$  is also differentiable.
- ▶ So, the set of differentiable functions forms what is called a *real vector space*.

# Linear systems

A system of linear (algebraic) equations is a system like this:

$$a_{11}x_1 + a_{12}x_2 + a_{13}x_3 + a_{14}x_4 = b_1$$

$$a_{21}x_1 + a_{22}x_2 + a_{23}x_3 + a_{24}x_4 = b_2$$

$$a_{31}x_1 + a_{32}x_2 + a_{33}x_3 + a_{34}x_4 = b_3$$

For example, initial value problems give rise to such systems, as did undetermined coefficients:

$$\begin{aligned} c_1 + c_2 &= 2 \\ -4c_1 + 2c_2 &= -10 \end{aligned}$$

# Matrix form of a system

$$\begin{pmatrix} a_{11} & a_{12} & a_{13} & a_{14} \\ a_{21} & a_{22} & a_{23} & a_{24} \\ a_{31} & a_{32} & a_{33} & a_{34} \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{pmatrix} = \begin{pmatrix} b_1 \\ b_2 \\ b_3 \end{pmatrix}$$

It's cleaner and allows us to focus on the arithmetic.

$$\begin{pmatrix} 1 & 1 \\ -4 & 2 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} = \begin{pmatrix} 2 \\ -10 \end{pmatrix}$$

# Solutions of matrix equations

- ▶ The vector of  $x_i$  is the variable of the matrix equation.
- ▶ A *solution* of the matrix equation is a choice of  $x_i$  that makes all the individual linear equations true.

Two matrix equations are *equivalent* if they have exactly the same set of solutions.

- ▶ Matrices are solved by transforming them into equivalent equations whose solutions are obvious.

# An obvious matrix

Lots of other matrix equations' solutions are obvious, but this is what I really meant:

$$\begin{pmatrix} \color{red}{1} & 2 & \color{blue}{0} & \color{blue}{0} & 2/3 \\ \color{blue}{0} & 0 & \color{red}{1} & \color{blue}{0} & -1 \\ \color{blue}{0} & 0 & \color{blue}{0} & \color{red}{1} & -10 \\ \color{blue}{0} & 0 & \color{blue}{0} & \color{blue}{0} & 0 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \\ x_5 \end{pmatrix} = \begin{pmatrix} b_1 \\ b_2 \\ b_3 \\ b_4 \end{pmatrix}$$

# Reduced echelon form

We say a matrix  $A$  is in reduced echelon form if:

- ▶ the first nonzero entry of each row is a 1
  - ▶ such an entry is called a *pivot* or *leading 1*
- ▶ each pivot is the only nonzero entry in its column
- ▶ each of the pivots after the first one appears to the right of the previous pivot
- ▶ each row without a pivot follows all rows with a pivot

Such a matrix certainly has an obvious solution set, and:

- ▶ Every matrix is equivalent to exactly one matrix in reduced echelon form.

# How do we find a reduced echelon equivalent?

- ▶ The same way we solve the equations to which the matrix equation corresponds: by adding and subtracting the rows from one another.
- ▶ We'll need to keep track of the RHS too, so add it as the last column of the matrix. We'll manipulate this matrix:

$$\begin{pmatrix} a_{11} & a_{12} & \cdots & a_{1n} & b_1 \\ a_{21} & a_{22} & \cdots & a_{2n} & b_2 \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ a_{m1} & a_{m2} & \cdots & a_{mn} & b_m \end{pmatrix}$$



# Gaussian elimination

- ▶ There are three families of operations on matrices that preserve solution sets.
  - ▶ If you do any of these operations to a matrix, the resulting matrix is equivalent.
- ▶ Swapping rows
  - ▶ this doesn't even change the equations, just their order
- ▶ Multiplying a row by a nonzero number
  - ▶ this changes the equations, but not the solution set
- ▶ Adding a nonzero multiple of a row to another row
  - ▶ also doesn't change the solution set.

Using these operations, it's possible to transform each matrix into its unique reduced echelon equivalent.

# In practice: upper-triangular

If you are row-reducing a matrix by hand, reduced echelon form is overkill a lot of the time.

- ▶ Reduce to an upper-triangular matrix
- ▶ All nonzero entries on or above the main diagonal
  - ▶ that is, the upper-left-to-lower-right diagonal
  - ▶ with slope  $-1$

# Entering vectors in Sage

- ▶ Initialize a vector value with `vector()`
- ▶ `u = vector(QQ, [1, 3/2, -1])`
- ▶ Use `QQ` to display fractions instead of decimals
- ▶ `u`
  - ▶ `(1, -3/2, 1)`
- ▶ `v = vector(RR, [1, 3/2, -1]); v`
  - ▶ `(1.0000000000000000, 1.5000000000000000, -1.0000000000000000)`

# Entering matrices

- ▶ Either enter a matrix as a vector of its rows
  - ▶ `A = matrix(QQ, [[1, 2], [3, 4], [5, 6]])`
- ▶ or as a list with specification of number of rows
  - ▶ `A = matrix(QQ, 2, [1,2,3,4,5,6])`
- ▶ Obtain reduced echelon form with `rref`:
  - ▶ `B = A.rref()`

# Example

▶ `A = matrix(QQ, 2, [1, 1, 2, -4, 2, -10]); show(A)`

▶ 
$$\begin{pmatrix} 1 & 1 & 2 \\ -4 & 2 & -10 \end{pmatrix}$$

▶ `B = A.rref(); show(B)`

▶ 
$$\begin{pmatrix} 1 & 0 & 7/3 \\ 0 & 1 & -1/3 \end{pmatrix}$$

# Row operations

- ▶ Want to use Sage to check your work in performing row operations?
  - ▶ The matrix methods below may be useful.
  - ▶ These methods are *destructive*: they change the entries of the matrix on which they're called.
- ▶ `A.rescale_row(i,a)`
  - ▶ multiply row  $i$  by  $a$
- ▶ `A.add_multiple_of_row(i,j,a)`
  - ▶ add  $a$  times row  $j$  to row  $i$
- ▶ `A.swap_rows(i,j)`
  - ▶ swap rows  $i$  and  $j$

# Sage lab assignment

Use Sage to solve the systems of linear equations.

$$\begin{array}{rclcl} 3x & + & 3y & + & 12z & = & 6 \\ x & + & y & + & 4z & = & 2 \\ 2x & + & 5y & + & 20z & = & 10 \\ -x & + & 2y & + & 8z & = & 4 \end{array} \qquad \begin{array}{rclcl} 2x & + & 10y & + & 2z & = & 6 \\ x & + & 5y & + & 2z & = & 6 \\ x & + & 5y & + & z & = & 3 \\ -3x & - & 15y & + & 3z & = & -9 \end{array}$$

$$\begin{array}{rclcl} 2x & + & y & - & z & + & 2w & = & -6 \\ 3x & + & 4y & & & + & w & = & 1 \\ x & + & 5y & + & 2z & + & 6w & = & -3 \\ 5x & + & 2y & - & z & - & w & = & 3 \end{array}$$