

Mathematics 352
Variation of Parameters

1. Use variation of parameters to find the general solution of the equation

$$y'' - y' - 2y = e^{3t}.$$

Solution: The characteristic polynomial is $(r - 2)(r + 1)$, so $y_1 = e^{2t}$ and $y_2 = e^{-t}$ are a fundamental set of solutions to the associated homogeneous equation. We compute $W(y_1, y_2) = (e^{2t})(-e^{-t}) - (2e^{2t})(e^{-t}) = -3e^t$. The variation of parameters formulas yield

$$\begin{aligned} u_1 &= - \int \frac{e^{-t} e^{3t}}{-3e^t} dt = \int \frac{e^t}{3} dt = \frac{e^t}{3}, \\ u_2 &= \int \frac{e^{2t} e^{3t}}{-3e^t} dt = -\frac{1}{3} \int e^{4t} dt = -\frac{1}{12} e^{4t}. \end{aligned}$$

Therefore a particular solution is

$$u_1 y_1 + u_2 y_2 = \frac{1}{3} e^{3t} - \frac{1}{12} e^{3t} = \frac{1}{4} e^{3t}.$$

2. Use variation of parameters to find the general solution of the equation

$$y'' - 4y' + 5y = e^{-4x}.$$

3. Verify that $y_1 = t^2$ and $y_2 = t^{-1}$ are solutions of the equation

$$t^2 y'' - 2y = 0.$$

4. Use variation of parameters to find the general solution of the equation

$$t^2 y'' - 2y = 3t^2 - 1.$$

You can assume $t > 0$ wherever necessary.

Solution: As above we let $y_1 = t^2$ and $y_2 = t^{-1}$. Since their Wronskian is

$$t^2(-t^{-2}) - 2t(t^{-1}) = -1 - 2 = -3,$$

they are an independent set of solutions to the associated homogeneous equation. The variation of parameters equations give

$$\begin{aligned} u_1 &= - \int \frac{t^{-1}(3t^2 - 1)}{-3} dt = \frac{1}{3} \int 3t - t^{-1} dt = \frac{1}{3} \left(\frac{3}{2} t^2 - \ln|t| \right) = \frac{1}{2} t^2 - \frac{1}{3} \ln|t|, \\ u_2 &= \int \frac{t^2(3t^2 - 1)}{-3} dt = \frac{1}{3} \int 3t^4 - t^2 dt = \frac{1}{3} \left(\frac{3}{5} t^5 - \frac{1}{3} t^3 \right) = \frac{1}{5} t^5 - \frac{1}{9} t^3. \end{aligned}$$