

Cross products; classifying planes

Math 251 Calculus 3

September 20, 2013

Distributing and the cyclic relation

Remember: $\hat{i} \times \hat{j} = \hat{k}$, $\hat{j} \times \hat{k} = \hat{i}$, and $\hat{k} \times \hat{i} = \hat{j}$. Plus, scalars operate as expected. So,

$$\begin{aligned}
 (\hat{i} + \hat{j} + \hat{k}) \times (2\hat{i} + 3\hat{j} + 4\hat{k}) &= \hat{i} \times (2\hat{i} + 3\hat{j} + 4\hat{k}) + \hat{j} \times (2\hat{i} + 3\hat{j} + 4\hat{k}) \\
 &\quad + \hat{k} \times (2\hat{i} + 3\hat{j} + 4\hat{k}) \\
 &= 2\hat{i} \times \hat{i} + 3\hat{i} \times \hat{j} + 4\hat{i} \times \hat{k} \\
 &\quad + 2\hat{j} \times \hat{i} + 3\hat{j} \times \hat{j} + 4\hat{j} \times \hat{k} \\
 &\quad + 2\hat{k} \times \hat{i} + 3\hat{k} \times \hat{j} + 4\hat{k} \times \hat{k} \\
 &= (3 - 2)(\hat{i} \times \hat{j}) + (2 - 4)(\hat{k} \times \hat{i}) + (4 - 3)(\hat{j} \times \hat{k}) \\
 &= \hat{k} - 2\hat{j} + \hat{i} = \langle 1, -2, 1 \rangle.
 \end{aligned}$$

Warm-up, II

Find the cross products:

- ▶ $\langle 2, 1, 3 \rangle \times \langle 4, 2, 1 \rangle$
- ▶ $\langle -1, 0, 1 \rangle \times \langle 2, -1, 5 \rangle$

Some setup for classifying planes

Recall:

- ▶ Each ordered pair of points in \mathbf{R}^3 determines a vector. How is this vector determined? How do you get its entries?

New fact/definition:

Some setup for classifying planes

Recall:

- ▶ Each ordered pair of points in \mathbf{R}^3 determines a vector. How is this vector determined? How do you get its entries?
 - ▶ Subtract the tail from the head.

New fact/definition:

Some setup for classifying planes

Recall:

- ▶ Each ordered pair of points in \mathbf{R}^3 determines a vector. How is this vector determined? How do you get its entries?
 - ▶ Subtract the tail from the head.

New fact/definition:

- ▶ If \vec{v} is a vector in \mathbf{R}^3 and \mathbf{P} is a plane, we say that \vec{v} is *normal* to \mathbf{P} if, for each vector \vec{w} contained in \mathbf{P} , we have $\vec{v} \cdot \vec{w} = 0$.

Some setup for classifying planes

Recall:

- ▶ Each ordered pair of points in \mathbf{R}^3 determines a vector. How is this vector determined? How do you get its entries?
 - ▶ Subtract the tail from the head.

New fact/definition:

- ▶ If \vec{v} is a vector in \mathbf{R}^3 and \mathbf{P} is a plane, we say that \vec{v} is *normal* to \mathbf{P} if, for each vector \vec{w} contained in \mathbf{P} , we have $\vec{v} \cdot \vec{w} = 0$.
- ▶ A vector is *contained* in \mathbf{P} if both its head and its tail (and hence, all the point on the vector's “body”) are in \mathbf{P} .

Warm-up for Workshop 03

- Choose a pair of orthogonal vectors and draw them in standard position. Your vectors must not be multiples of \hat{i} or \hat{j} , but I would advise you to choose vectors with z -entry 0—then, you can get away with drawing \mathbf{R}^3 as a plane viewed along the positive z -axis. *Hint.* Use the dot product to make sure your vectors really are perpendicular.

Warm-up for Workshop 03

- ▶ Choose a pair of orthogonal vectors and draw them in standard position. Your vectors must not be multiples of \hat{i} or \hat{j} , but I would advise you to choose vectors with z -entry 0—then, you can get away with drawing \mathbf{R}^3 as a plane viewed along the positive z -axis. *Hint.* Use the dot product to make sure your vectors really are perpendicular.
- ▶ Because your vectors are orthogonal, there is a rectangle based on these vectors (draw two more sides). Use the cross product–sine formula to verify that the area of the rectangle is equal to the length of the cross product of your vectors.

General parallelogram

- ▶ Start a new picture, and draw a new pair of vectors in standard position. Make sure that the angle between them is *not* a multiple of $\pi/2 = \tau/4$. Use the dot product–cosine formula to do this.

General parallelogram

- ▶ Start a new picture, and draw a new pair of vectors in standard position. Make sure that the angle between them is *not* a multiple of $\pi/2 = \tau/4$. Use the dot product–cosine formula to do this.
- ▶ Having ensured your angle is not a multiple of $\pi/2 = \tau/4$, you know that your new pair of vectors forms a nonrectangular parallelogram. Use trigonometry to express the area of this parallelogram in terms of the vector lengths.

General parallelogram

- ▶ Start a new picture, and draw a new pair of vectors in standard position. Make sure that the angle between them is *not* a multiple of $\pi/2 = \tau/4$. Use the dot product–cosine formula to do this.
- ▶ Having ensured your angle is not a multiple of $\pi/2 = \tau/4$, you know that your new pair of vectors forms a nonrectangular parallelogram. Use trigonometry to express the area of this parallelogram in terms of the vector lengths.
- ▶ Check that the length of the cross product yields the area of the parallelogram in this case also.

Wrap-up

Together with the orientation and complementarity properties, the cross product–sine formula *uniquely determines the cross product*. This means that, for any pair of vectors \vec{v} and \vec{w} , there is only one vector \vec{u} satisfying all three. No matter how it is obtained, it must be equal to the cross product $\vec{v} \times \vec{w}$.

- Use the cross product–sine formula and the orientation property of the cross product to *derive* the rule $\hat{j} \times \hat{i} = -\hat{k}$.