

Food web rewiring in a changing world

Timothy J. Bartley^{1,2,6*}, Kevin S. McCann^{2,6}, Carling Bieg², Kevin Cazelles², Monica Granados^{2,3}, Matthew M. Guzzo², Andrew S. MacDougall², Tyler D. Tunney^{4,5} and Bailey C. McMeans¹

Climate change is asymmetrically altering environmental conditions in space, from local to global scales, creating novel heterogeneity. Here, we argue that this novel heterogeneity will drive mobile generalist consumer species to rapidly respond through their behaviour in ways that broadly and predictably reorganize — or rewire — food webs. We use existing theory and data from diverse ecosystems to show that the rapid behavioural responses of generalists to climate change rewire food webs in two distinct and critical ways. First, mobile generalist species are redistributing into systems where they were previously absent and foraging on new prey, resulting in topological rewiring — a change in the patterning of food webs due to the addition or loss of connections. Second, mobile generalist species, which navigate between habitats and ecosystems to forage, will shift their relative use of differentially altered habitats and ecosystems, causing interaction strength rewiring — changes that reroute energy and carbon flows through existing food web connections and alter the food web's interaction strengths. We then show that many species with shared traits can exhibit unified aggregate behavioural responses to climate change, which may allow us to understand the rewiring of whole food webs. We end by arguing that generalists' responses present a powerful and underutilized approach to understanding and predicting the consequences of climate change and may serve as much-needed early warning signals for monitoring the looming impacts of global climate change on entire ecosystems.

Following the poleward shift in species distributions observed with climate warming, Blanchard¹ quipped that the resultant massive compositional changes to the Arctic marine food web² are akin to nature 'rewiring' itself. This northern range expansion is dominated by species that are relatively large, highly mobile and foraging generalists, and so increases the Arctic marine food web's complexity². This influx of generalists thus fundamentally alters the structure of the Arctic marine food web, making it both more connected and less modular, and also perhaps making interaction strengths weaker on average. This one example highlights a potentially global phenomenon: ongoing climate change will continue to fundamentally restructure — that is, 'rewire' — ecosystems. And yet, the ways in which food webs will rewire remain nebulous^{2,3}.

The rewiring of Earth's food webs with climate change should be first detectable as rapid behavioural responses that are most prominently exhibited by mobile generalist species (see Box 1 for key definitions)^{4–6}. For example, Kortsch et al.² found that as waters warm, the fish species advancing north in the marine Barents Sea ecosystem tend to be high-trophic-level consumers that are mobile and forage on a large range of resources. Similarly, many of the terrestrial species dispersing poleward in response to warming are capable of rapid, long-range movement^{7,8}. Species that possess this set of key correlated traits (high mobility in terms of movement or dispersal and flexibility in both foraging and habitat use) comprise what are commonly referred to as generalists² (Box 1). Generalist species are often capable of responding to resource and environmental variation through their behaviour, linking various habitats and ecosystems by transcending the boundaries between them traditionally deemed by ecologists. By linking otherwise distinct habitats and ecosystems, these species structure food webs in space^{2,9,10}. The movement of generalist species clearly has potentially significant implications for material and energy flow within and across ecosystems^{5,6,11,12}, but behavioural responses have been largely

overlooked by ecologists studying how food webs are rewiring with climate change.

In what follows, we argue that the responses of mobile generalist species rewire food webs in a changing world. We first show that climate change has asymmetrical impacts in space from global to local scales, producing novel heterogeneity in environmental conditions worldwide to which species are poised to respond (Fig. 1). We then discuss emerging studies that show this novel heterogeneity drives generalist species to respond rapidly and predictably to novel conditions through their behaviour¹¹. These generalists' responses alter food web in two related but qualitatively distinct ways: by changing food web topology (that is, topological rewiring *sensu* Blanchard¹) and by changing the strengths of existing interactions (that is, interaction strength rewiring). We illustrate rewiring driven by climate warming using two detailed example food webs, one aquatic (Box 2) and one terrestrial (Box 3), each made up of ectothermic organisms that are highly sensitive to changes in temperature¹³. We then illustrate how whole groups of species with shared traits can exhibit unified, aggregate community behavioural responses to climate change that could rewire entire food webs. We end by arguing that these results enticingly suggest that rapid behavioural responses of generalists to climate change represent a powerful tool in monitoring nature's responses to environmental change and can act as a potent addition to the early warning signals (EWSs) toolbox.

Climate change creates novel heterogeneity across scales

While many aspects of global change are thought to homogenize ecosystems and landscapes^{14–16}, climate change impacts are also expected to vary spatially, transforming environmental heterogeneity in subtle but significant ways across scales (Fig. 1)¹⁷. The impacts of climate change are well known to be asymmetrical on large spatial scales (that is, one region or hemisphere is impacted more than another). On the global scale, rates of warming are

¹Department of Biology, University of Toronto Mississauga, Mississauga, Ontario, Canada. ²Department of Integrative Biology, University of Guelph, Guelph, Ontario, Canada. ³Wildlife Conservation Society Canada, Thunder Bay, Ontario, Canada. ⁴Center for Limnology, University of Wisconsin-Madison, Madison, WI, USA. ⁵Gulf Fisheries Centre, Fisheries and Oceans Canada, Moncton, New Brunswick, Canada. ⁶These authors contributed equally: Timothy J. Bartley, Kevin S. McCann. *e-mail: timothy.bartley@gmail.com

Box 1 | Key terms and concepts relating to the rewiring of the world's food webs with climate change

Rewiring	Fundamental changes in the structure of a food web that alter the pathways of nutrients and/or energy in an ecosystem. Rewiring can occur from changes in either food web topology (topological rewiring) or interaction strength (interaction strength rewiring).
Topological rewiring	Changes in the topology of a local food web (that is, who eats whom), which often result from novel species introductions and/or the loss of species.
Interaction strength rewiring	Changes in the strengths of interactions in a local food web (that is, the magnitude of the effect on the energy flow from one species to another), which often result from changes in the consumptive demand of a consumer associated with changes in the consumer's behaviour.
Novel heterogeneity	Unprecedented, increased distinctiveness in the relative environmental conditions of habitats that arises from the asymmetrical impacts of climate change. These asymmetrical impacts of climate change occur on various scales, from within ecosystems to between entire hemispheres.
Behavioural response	Changes in behaviour by a species or suite of species, such as dispersal, movement, habitat use and foraging actions, that occur in reaction to changes in environmental conditions.
Mobile generalist species	Species that tend to possess high mobility (in terms of movement or dispersal) and flexibility in both foraging and habitat use. As a result, these species tend to have a large dietary breadth and link macrohabitats through their foraging actions, allowing them to respond behaviourally to environmental variation in space and time.
Structural EWSs	Properties in food web structure (topology and interaction strengths) that suggest imminent and drastic changes in ecosystems, such as major shifts in stability or regime.

asymmetric between hemispheres, with warming of land and oceans in the Northern Hemisphere outpacing the Southern Hemisphere^{18,19} (Fig. 1a). Other global scale properties are being asymmetrically altered too, including precipitation²⁰, El Niño and La Niña frequencies²¹, and ice extent²². Climate models also predict asymmetry in responses within hemispheres, with polar and temperate ecosystems being more strongly impacted than tropical and equatorial systems^{18,23}. Regional effects of climate change include greater temperature increases on Earth's land masses than in the oceans²⁴ (Fig. 1b). Taken together, these asymmetrical impacts are leaving a complex large-scale footprint of climate change.

Asymmetrical climate change impacts are also expected on smaller scales, both within and across ecosystems, although they are not as well studied as global scale responses^{18,21,23,25}. These smaller scale asymmetrical impacts are likely to arise because ecosystems and the various habitats that comprise them have different physical and abiotic properties that differentially filter the impacts of climate change¹¹, comparable to the factors that create climate refugia²⁶. The result is asymmetrical impacts of climate change in different habitats and ecosystems (for example, one habitat or ecosystem, say, warms more or more rapidly than another), creating a small-scale heterogenous mosaic in space. For example, while the surface waters of most lakes worldwide are warming, warming rates vary between lakes depending on local properties such as surface area, shape and depth²⁷. Within lakes, thermal stratification produces asymmetric warming between shallow nearshore and deep offshore macrohabitats because surface waters that warm under direct contact with the air do not mix with deeper waters, which remain cold even in the summer (Fig. 1c)^{22–24}. Similarly, in terrestrial systems, climate change is predicted to impact mountainous regions asymmetrically via feedbacks between latitude and elevation that depend on local topography, elevation, slope and treelines^{28–30}. On even smaller scales, global changes such as warming or increased precipitation interact with nutrient conditions to increase the local heterogeneity in limiting factors in terrestrial ecosystems^{28–31}. These asymmetric climate change impacts from micro- to macroscales will produce a novel, transformed heterogenous palette of fine- and coarse-grained habitats and ecosystems.

Rewiring through behavioural responses

Novel heterogeneity stemming from climate change ought to elicit rapid behavioural responses by mobile generalist species. These responses could be due to either the direct physiological

consequences of these new environmental conditions or the resultant changes in resource availability or distribution in space. Importantly, mobile high-trophic-level generalist species can navigate across the landscape, moving between habitats, ecosystems and even hemispheres, structuring food webs in space^{10,31–33}. The responses of these species should therefore rewire food webs in two key ways (Box 1). First, generalist species will expand into systems where they were previously rare or absent, resulting in topological rewiring — the addition or loss of food web connections or whole food web pathways. Topological rewiring is analogous to adding and removing 'wires' in an electrical network. Second, generalist species will shift how they move and forage across multiple habitats and ecosystems, causing interaction strength rewiring — changes in the consumption rates of existing food web connections, thus rerouting energy flows through existing pathways. Interaction strength rewiring is akin to altering the 'load' on the wires in an electrical network. Because the mobile generalist species that transcend the spatial boundaries between habitats and ecosystems link their energy and nutrient dynamics, their behavioural responses promise to fundamentally shift the interplay between sub-webs or food web compartments in different habitats and ecosystems. And when groups of generalist species with shared traits respond in concert to the altered spatial mosaic created by climate change, they have the potential to fundamentally reorganize the structures of whole ecosystems and biomes.

Topological rewiring. On the global scale, numerous studies have demonstrated the poleward movement of many species in various ecosystems worldwide, altering community structure and potentially restructuring local food webs^{2,3,7,8,34–36}. Similar patterns exist across altitudinal gradients, as exemplified by upslope shifts in the distribution of bees, butterflies and birds^{37–39}. Importantly, these shifts tend to be dominated by generalist species, which are often more mobile and adapted to deal with spatial variation in resources and conditions^{2,7,8,35,36}. These latitudinal and altitudinal advances of generalists strongly imply that local food webs should be experiencing species introductions and so undergoing topological rewiring as they gain new connections^{1,2,40}. Local food webs may also lose connections because of the loss of species^{36,40,41}. Notably, though, species ranges are expanding poleward and upslope faster than they are contracting, and the rate of species invasions appears to be outpacing the rates of local declines^{7,42}. Taken together these changes suggest that, on average, we expect the reshuffling of species to skew local diversity towards a more generalized set of species^{3,43–45}.

This skew towards generalists can fundamentally alter the topology of local food webs, increasing connectance and reducing modularity of the rewired food webs. These topological changes have potentially dramatic implications for stability and the maintenance of biodiversity^{46–48}, and topological rewiring from range expansion of tropical species has indeed been associated with dramatic phase shifts in temperate marine food webs⁴⁹.

One clear example of topological rewiring comes from Kortsch et al.², who examined how climate change will impact the food web of the Barents Sea, which borders on the Arctic Ocean. They found that the boreal fish moving poleward into the Barents Sea tended to be omnivorous generalist fish species. The addition of these generalists into the food web increased connectance and simultaneously reduced modularity by linking previously disparate modules corresponding to pelagic and benthic macrohabitats. Kortsch et al.² anticipate that the outcome of this topological rewiring in the Barents Sea food web will be altered patterns in carbon flow within this food web.

Interaction strength rewiring. Importantly, the asymmetrical impacts of climate change across scales are driving mobile generalist species to respond in ways other than redistributing across the globe and driving topological rewiring. Mobile generalist species may simultaneously alter the strengths of trophic interactions within food webs^{50–52}. The distribution of interaction strengths, which is determined by the foraging actions of consumers, is a key part of food web structure that is well known to underlie the maintenance of diversity in, and the stability of, complex communities^{46,48,52–55}. Research on food web rewiring has focused on topological changes in food web structure, with less emphasis on how interaction strengths will be altered by climate change. Some research has shown expected changes in average interaction strength with warming⁵⁶. However, altered heterogeneity in space from climate change ought to change other aspects of interaction strength, such as strengthening some interactions and weakening others, or rearranging the distribution of interaction strengths.

Many generalists forage across the landscape, coupling spatially distinct habitats and ecosystems⁵⁷. Interaction strength rewiring can occur when these spatially distinct habitats and ecosystems are asymmetrically altered by climate change, driving generalists to change their relative use of resources in space. For example, asymmetrical warming between habitats may have physiological consequences for a generalist. The warming may increase metabolic demand, driving the generalist to increase its food consumption and thereby increasing the flow of energy through the existing food web pathways. However, if the metabolic consequences are too costly in the warmer habitat, a generalist may change its behaviour by decreasing use of food sources in the warmer habitat and increasing use of cooler habitats. This response would simultaneously reduce consumption on the existing food web pathway derived from the warmer habitat and increase consumption on the food web pathway from the cooler habitat. In this way, the amount of energy flow along existing warmed-up pathways (wires) is diminished, and the energy flow along other, less impacted pathways may be unchanged or increased. This change in mobile generalist consumers' behaviour also results in the decoupling of adjacent habitats or ecosystems. Given that some migratory species, such as some whales and seabirds, couple the Northern and Southern Hemispheres and adjust their behaviour to changes in climatic conditions³⁸, interaction strength rewiring also has the potential to play out on vast spatial scales.

Interaction strength rewiring has been documented in diverse ecosystems across the globe. Among the best-studied examples of interaction strength rewiring are boreal lake ecosystems in Canada that have lake trout (*Salvelinus namaycush*) as a top predator. The cold-water-adapted lake trout respond to the differential warming of the nearshore macrohabitat by retreating to the cooler offshore




Fig. 1 | The asymmetrical impacts of climate change create novel heterogeneity, from local to global spatial scales. a, Global temperature data from 1880 to 2017 indicate temperatures in the Northern Hemisphere are increasing more rapidly than in the Southern Hemisphere^{103,104}. **b**, The ratio of land/sea warming rates from many climate change models shows that land is warming faster than seas²⁴. **c**, Owing to thermal stratification in lakes, indicated by this vertical temperature profile, the nearshore (littoral) areas and surface waters of lakes are warming faster than deep and offshore (pelagic) areas. **d**, Temperature increases vertically farther from the soil surface to the top of grasses in grassland ecosystems¹⁰⁵.


habitat and reducing their reliance on nearshore food resources (for details, see Box 2). Similarly, other studies^{4,59–61} have shown interaction strength rewiring in grassland invertebrate food webs. When experimental warming shifts the vertical temperature gradient in grasslands, the sit-and-wait predatory spider *Pisaurina mira* moves down towards the relatively cool soil microhabitat, introducing novel intraguild predation causing extirpation of the active predatory spider *Phidippus rimator*, reducing the feeding time of the grasshopper *Melanoplus femurrubrum*, and indirectly

Box 2 | The rewiring of lake food webs with climate change

One well-documented example of rewiring comes from the north-temperate lake ecosystems that have cold-water lake trout (*Salvelinus namaycush*) as a top predator. Lake trout are highly mobile foragers that prefer cold water and can actively move and feed between shallow nearshore (littoral) and deep offshore (pelagic) macrohabitats³¹ (panel a of the figure in Box 2). The extent of littoral foraging by lake trout is governed by the thermal accessibility of the nearshore macrohabitat because lake trout experience a cost when making forays from their deep pelagic refuge into warmer littoral areas^{31,32,100,107–110}. However, these two macrohabitats that lake trout couple will be differentially impacted by climate change, which is expected to warm the nearshore and surface macrohabitat faster than the deep offshore macrohabitat^{111,112}. This differential warming of the nearshore zone forces lake trout to respond by decoupling from the nearshore, with lake trout moving into deeper colder waters and relying less heavily on nearshore resources (panels a and b of the figure in Box 2). The shift in habitat use by lake trout is revealed by long-term telemetry, which shows increases in the depth of detection as the nearshore macrohabitat warms (panel c of the figure in Box 2). This is paralleled with a shift in lake trout diet that is apparent in stable carbon isotope signatures and stomach content analysis, which both show reduced reliance on nearshore food resources with increasing summer air temperature (panel b of the figure in Box 2)^{32,108}. The behavioural response of lake trout to the differentially warmed littoral habitat thus represents rewiring of carbon flow through the whole lake ecosystem.

This rewiring of lake ecosystems has important consequences for both lake trout and whole lake ecosystems. Difficulty in garnering prey from the nearshore reduces both growth and condition factor in lake trout³¹, suggesting a potential loss of stability (in that lake trout persistence is threatened) from reduced access to the littoral carbon pathway. Reduced nearshore foraging changes the life history traits of lake trout and reduces density in a way that may erode their top-down effects^{100,106}. Temperate lake ecosystems are also highly seasonal environments, and lake trout show important seasonal shifts in behaviour and habitat use^{31,110}. Climate change is altering various abiotic factors in lakes and affects some seasons more than others, suggesting that climate change will alter the seasonal flexes in lake food web structure. Such climate change impacts that are asymmetrical in time may also drive food web rewiring. Importantly, other species in boreal lake ecosystems, including the planktivorous cisco (*Coregonus artedii*) and the piscivorous walleye (*Sander vitreus*), both similarly display paired behavioural and dietary shifts away from the nearshore in

increasingly warm lakes, showing that rewiring occurs at multiple places in lake food webs¹⁰⁶. Such behavioural and dietary shifts may be a common response of thermally sensitive species in these lakes, producing unified aggregate behavioural responses to climate change (see 'Aggregate rewiring' and Fig. 3). **a,b** were adapted from ref. ³², PNAS; **c** was adapted from ref. ³¹, PNAS.



altering the biomass of grasses and herbs (for details, see Box 3). Additionally, Yurkowski et al.⁶² show how the northward advance of a species due to climate change can cause interaction strength rewiring in Arctic marine ecosystems. As capelin (*Mallotus villosus*) move northward into Cumberland Sound, Nunavut, Canada, both beluga whales (*Delphinapterus leucas*) and Greenland halibut (*Reinhardtius hippoglossoides*) increase their foraging on forage fish, changing the summertime relationship between belugas and halibut from a primarily predator–prey interaction to a primarily competitive interaction (Fig. 2a)⁶². A larger scale example of across ecosystem interaction strength rewiring takes place at the sea–land interface. During periods of reduced sea ice, polar bears (*Ursus maritimus*) spend more time on land, spatially isolated from their preferred prey of ringed seals (*Pusa hispida*; see Fig. 2b)^{63,64}. At this time, the bears predate more on nesting seabirds and their eggs, altering the strengths of their interactions with these resources^{63,65}. This foraging switch is believed to be insufficient for polar bears to


maintain their condition, which is expected to negatively impact their populations⁶⁶. Interaction strength rewiring may even occur on the largest spatial scales because migratory seabirds and whales, which couple hemispheres, now have to migrate farther poleward with climate change⁶⁷.

Spatially asymmetrical climate change may also drive interaction strength rewiring in more complex ways, such as by producing phenological shifts. One intriguing example is that of Kodiak brown bears (*Ursus arctos middendorffi*), which feed on both terrestrial red elderberry (*Sambucus racemosa*) and aquatic sockeye salmon (*Oncorhynchus nerka*)⁶⁸ (Fig. 2c). While the productivity of these two resources was previously staggered in time, climate impacts pushed the elderberry to bloom earlier and now peak in synchrony with the relatively unaffected salmon spawns. This temporal synchronization effectively decoupled a connection between terrestrial and aquatic habitats that was mediated by bears⁶⁸. As climate change research has tended to initially cling to temperature changes

Box 3 | The rewiring of grassland food webs with climate change

Temperate grassland food webs are expected to rewire in the face of climate change^{4,59–61}. Grasslands have naturally occurring vertical temperature profiles corresponding to distance from the ground, creating distinct microhabitats near the soil surface and at the top of the grass canopy (Fig. 1d). Under ambient temperatures, the two spider predators are spatially separated, with the sit-and-wait predator *Phidippus mira* near the tops of the grasses and the active predator *Phidippus rimator* lower down near the soil (panel a of the figure in Box 3)¹⁰⁵. Experimental warming shifts the entire temperature gradient^{4,59–61}, driving the sit-and-wait predator *P. mira* to respond by moving down closer to the shaded thermal refugium near the soil surface as greater heights become too warm^{4,33,105,113,114} (panel b of the figure in Box 3). When *P. mira* responds by changing its microhabitat use and moving down, it spatially overlaps with the sit-and-wait predatory spider, *P. rimator*, driving intraguild predation that knocked out the *P. rimator* in Barton and Schmitz's¹⁰⁵ experimental microcosms. The behavioural response of *P. mira* impacted the foraging behaviour of herbivorous grasshoppers, *Melanoplus femurrubrum*. As *P. mira* moved down and farther away from *M. femurrubrum* with increasing temperature, *M. femurrubrum* showed increasing feeding time (panel c of the

figure in Box 3)⁶¹. As a result, changes in top-down effects from predatory spiders drove indirect effects on herbaceous plant biomass, with the biomass of grasses and herbs in experimentally warmed mesocosms significantly altered when compared to control mesocosms (panel d of the figure in Box 3)¹⁰⁵. This indirect effect is critical given that the direct effects of warming on plant biomass are less than the indirect effects of top-down control by spiders⁶⁰. These studies also hint at the consequences of this grassland food web rewiring, with the loss of *P. rimator* suggesting possible impacts on stability and the change in plant biomass suggesting possible changes to ecosystem function. Barton and co-authors have also looked at how factors such as wind and precipitation can rewire terrestrial food webs^{115,116}, implying food web rewiring occurs with other aspects of climate change. Intriguingly, Barton and Schmitz¹¹⁷ show that daytime and night-time warming have opposite effects on spider activity, producing opposite trophic cascades. These results suggest that climate change asymmetries in time may also drive food web rewiring. Taken together, these studies make a unique case study of how asymmetrical impacts of climate change rewire grassland food webs. a,b,d were adapted from ref. ¹⁰⁵, Wiley; c was adapted from ref. ⁶¹, Royal Society.



(as noted by VanDerWal et al.⁶⁹), many of our examples of rewiring focus on the impacts of asymmetrical warming. However, climate change is multifaceted, with many dimensions of climate change

(for example, precipitation) expected to be asymmetric in ways that similarly elicit rapid behavioural responses in mobile generalists and thus broadly rewiring food webs^{70,71}.




Fig. 2 | Three examples of food web rewiring with climate change from diverse ecosystems. a, Rewiring of the Arctic marine food web in Cumberland Sound, Nunavut, Canada. As capelin (*Mallotus villosus*) move northward into Arctic marine ecosystems, both beluga whales (*Delphinapterus leucas*) and Greenland halibut (*Reinhardtius hippoglossoides*) increase their foraging on forage fish. These responses change the summertime relationship between belugas and halibut from a primarily predator–prey interaction to a primarily competitive interaction⁶². **b**, Rewiring of the food web across the Arctic land–sea interface. During periods of reduced sea ice, polar bears (*Ursus maritimus*) spend more time on land, spatially isolated from their preferred prey of ringed seals (*Pusa hispida*). While on land, the bears predate more on nesting seabirds and their eggs and less on ringed seals, altering the strengths of their interactions with these resources^{63–65}. **c**, Rewiring of the food webs of coastal Pacific North America. Kodiak brown bears (*Ursus arctos middendorffi*) feed on both terrestrial red elderberry (*Sambucus racemosa*) and on sockeye salmon (*Oncorhynchus nerka*). While these two resources were previously staggered in time, climate impacts pushed the elderberry to bloom earlier and now in synchrony with salmon, effectively forcing the decoupling of terrestrial and aquatic habitat that was mediated by bears⁶⁸. **a** was adapted from ref. ⁶², Royal Society; **c** was adapted from ref. ⁶⁸, PNAS.

Aggregate rewiring. The bulk of research on how climate change impacts food webs has focused on one or a small number of species, with less focus on how climate change may reorganize whole food webs⁷². Yet, the rewiring of food webs is probably not limited to a single generalist species response; entire suites of species within a food web may respond en masse, especially if they share key traits that drive their responses. To test this idea, we expand on the previous research showing how lake trout responses to climate change rewire lake food webs (described in Box 2). Lake trout responses result from reduced accessibility of the differentially warmed nearshore macrohabitat in lakes. Lake trout are one of many cold-water-adapted fish that inhabit these lakes, and fish are generally relatively mobile⁷³. Thus, reduced thermal accessibility of the nearshore macrohabitat may drive many of these species to exhibit similar behavioural responses to that of lake trout, generating a unified response of the entire cold-water thermal guild (Fig. 3a). As expected, extensive spatial catch-per-unit-effort data from 721 lakes in Ontario, Canada⁷⁴ across a natural temperature gradient show that the cold-water guild is on average caught in deeper water in warmer lakes (Fig. 3b), indicating an aggregate behavioural response towards increased offshore habitat use. In addition,

most of the 13 cold-water species, which span several trophic levels, individually were on average caught in deeper water in warmer lakes (Fig. 3c). Because such shifts towards offshore habitat use are associated with decreased nearshore foraging^{31,32}, these unified behavioural results strongly imply significant rewiring throughout lake food webs, with major consequences for carbon flow in these ecosystems. Curiously, Dulvy et al.⁷⁵ document similar aggregate behavioural responses in the North Sea. As the bottom temperature of shallow shelf seas warmed from 1980 to 2004, the whole bottom-dwelling fish assemblage comprising 28 species moved into deeper waters⁷⁵. The deepening of fish assemblages in these ecosystems exemplifies two powerful case studies for how rapid behavioural responses to climate change are rewiring interaction strengths on the whole food web scale. If groups of species with shared traits are widespread, these unified, aggregate responses may be common with climate change. Aggregate behavioural responses would allow us to scale from understanding how single species rewire food webs to understanding how whole food webs rewire with climate change. In combination with the aggregate range shifts documented for some groups of species^{75,76}, aggregate responses may be vital to understanding food web rewiring with climate change.




Fig. 3 | The aggregate rewiring of food webs through the unified behavioural responses of entire suites of species. **a**, The aggregate behavioural response of cold-water fish to move into deeper, offshore waters with climate warming, which suggests the rewiring of boreal shield lake food webs. **b**, The residual average \log_{10} depth of capture for 13 cold-water fish species increases across a gradient of increasing average recent air temperature based on spatial catch-per-unit-effort data from 721 lakes in Ontario, Canada, indicating that cold-water species were on average caught in deeper water in warmer lakes¹⁰⁶ (see Supplementary Information). **c**, The slope coefficient (with standard error) for regression models of the residual average \log_{10} depth of capture across a spatial gradient of average recent air temperature for each of 13 cold-water species, showing that many species contribute to the unified behavioural response of these species to increased temperature¹⁰⁶ (see Supplementary Information). Figure adapted from ref. ¹⁰⁶, published under a Creative Commons licence (<https://creativecommons.org/licenses/by/2.5/ca/>).

Stability and structural EWSs

Here, we have argued that the asymmetrical impacts of climate change ought to broadly and predictably rewire food webs in terms of both topology and interaction strength. The impacts of climate change are often strongly linked with simultaneous changes in other forms of human perturbation (for example, species invasions) that may also drive rewiring^{77,78}. These broadly imposed human impacts are allowing generalist species to redistribute around the globe, functionally homogenizing biodiversity, overwhelming more specialized species⁷⁹ and rewiring food webs^{40,42,80}. As food web structure and stability are inextricably linked⁸¹, both topological rewiring and interaction strength rewiring have the potential to drastically alter stability. Despite the notion that heterogeneity largely plays a stabilizing role in ecosystems^{82–86}, the novel heterogeneity and behavioural responses we discuss here may not always act as a stabilizing force in newly rewired ecosystems⁸⁷. The topological changes in Arctic marine food webs documented by Kortsch et al.² — increased connectance and reduced modularity — tend to be destabilizing because they synchronize whole food web responses to perturbations^{46,53,54,88,89}. Yet, an influx of generalists may promote stability by weakening average interaction strengths⁹⁰ and allowing adaptive responses that mute variation⁹. Thus, in the interim, the exact ramifications of rewiring on stability may appear nebulous; however, the ultimate consequences of rewiring for stability are less ambiguous. With continuing asymmetrical climate change, some habitats and ecosystems are likely to become completely inaccessible for the mobile generalists that couple them, leading to extensive decoupling that is well known to be destabilizing⁹¹. In addition, altered climatic conditions are likely to make ecosystems and habitats inhospitable for some species and impact key ecological and metabolic rates that are strongly linked to stability, making extinctions prevalent and inevitable and racking up extinction debt in many ecosystems⁹². Biodiversity loss on this scale is strongly linked to a loss in stability.

Given that generalist species responses critically influence stability, ecologists may be able to use these responses as ‘structural’ EWSs to climate change impacts on ecosystems. EWSs have been successfully applied to forecast changes in diverse systems, from stock markets to ecosystems. Current ecological EWSs, which are largely based on time series of population abundance, have some

significant challenges because empirical time series are often too short to decipher key signatures of a looming loss in stability, such as critical slowing down^{93–95}. This aspect of EWSs is especially concerning for long-lived, higher-trophic-level organisms with population dynamic signatures such as cycles and generation times that span multiple decades⁹⁶. As a result, researchers have recently called for additional methods to be added to the EWS toolbox, including concomitant changes in spatial patterning within an ecosystem⁹⁷. Our arguments here suggest that monitoring the behaviour and foraging of high-trophic-level generalists can help detect key structural changes in food webs that indicate the imminent collapse of one or more species. Importantly, such behavioural assays using generalist species — the same species whose times series ought to be difficult to track — would expose pending collapse on much shorter timescales and with much less intensive sampling efforts than time-series-based approaches. For example, Guzzo et al.³¹ use data collected over a period of 11 years to show that lake trout, which can live for decades, exhibit rapid shifts in foraging and behaviour to warming. Given the rapid rate and large scale of environmental change worldwide, new structural indicators of looming change will be imperative to maintain the diversity and functioning of the biological systems on which we rely for critical ecosystem services^{4,98,99}. With further research, we can harness generalists’ responses to predict functional outcomes of climate change on the world’s ecosystems.

Ecologists are already documenting rapid behavioural responses to changing environmental conditions using a variety of tools that could serve as structural indicators of major changes to ecosystems. Increasingly common tools for diet analysis include biotracers such as stable isotopes, fatty acids and DNA barcoding, which can readily track changes in foraging behaviour¹⁰⁰. Yurkowski et al.⁶² (Fig. 2a) provide an excellent example of using stable isotopes to monitor ecosystem changes before and after the northward advance of a species with climate change. Theory suggests that the shift from a primarily predator–prey interaction to a primarily competitive interaction such as that documented by Yurkowski et al.⁶² generally corresponds to a significant reduction in stability, perhaps foreshadowing major changes to this ecosystem. Another example of dietary monitoring comes from seabirds, which are known to couple across enormous

spatial scales⁵⁸. Seabirds that feed on multiple prey items (for example, sardines and anchovies) have shown foraging shifts away from sardines that precede fishery surveys, which eventually indicated significant sardine population decline⁵⁸. Diet-based tools could be particularly powerful when paired with traditional methods of detecting changes in animal movement with climate change, such as telemetry^{31,101}. Perhaps these tools will reveal that generalists show increased behavioural variation prior to the looming major loss in system stability (for example, loss of a key resource), analogous to the increased population abundance variance predicted by classical EWS theory. Because behavioural changes such as movement are often accompanied by physiological changes, we might also expect a suite of organismal responses (for example, growth, age at maturity, activity levels) in concert with climate change. Generalists' dietary, behavioural and physiological responses together are likely to help address the unequivocal need for tools to detect looming collapses in ecosystems¹⁰² and provide an across-scale integrated approach to biomonitoring the relentless impacts of climate change¹⁰⁰.

Data availability

The data used to produce Figs. 1 and 3 can be found online at <https://doi.org/10.5281/zenodo.1158733>.

Received: 1 March 2018; Accepted: 28 November 2018;

Published online: 11 February 2019

References

- Blanchard, J. L. A. A rewired food web. *Nature* **527**, 7–8 (2015).
- Kortsch, S., Primicerio, R., Fossheim, M., Dolgov, A. V. & Aschan, M. Climate change alters the structure of Arctic marine food webs due to poleward shifts of boreal generalists. *Proc. R. Soc. B* **282**, 20151546 (2015).
- Kortsch, S. et al. Climate-driven regime shifts in Arctic marine benthos. *Proc. Natl Acad. Sci. USA* **109**, 14052–14057 (2012).
- Harmon, J. P. & Barton, B. T. On their best behavior: how animal behavior can help determine the combined effects of species interactions and climate change. *Ann. NY Acad. Sci.* **1297**, 139–147 (2013).
- Knowlton, J. L. & Graham, C. H. Using behavioral landscape ecology to predict species' responses to land-use and climate change. *Biol. Conserv.* **143**, 1342–1354 (2010).
- Kearney, M., Shine, R. & Porter, W. P. The potential for behavioral thermoregulation to buffer 'cold-blooded' animals against climate warming. *Proc. Natl Acad. Sci. USA* **106**, 3835–3840 (2009).
- Walther, G.-R. et al. Ecological responses to recent climate change. *Nature* **416**, 389–395 (2002).
- Parmesan, C. & Yohe, G. A globally coherent fingerprint of climate change impacts across natural systems. *Nature* **421**, 37–42 (2003).
- McCann, K. S. & Rooney, N. The more food webs change, the more they stay the same. *Phil. Trans. R. Soc. B* **364**, 1789–1801 (2009).
- Rooney, N., McCann, K. S. & Moore, J. C. A landscape theory for food web architecture. *Ecol. Lett.* **11**, 867–881 (2008).
- Schindler, D. E. Warmer climate squeezes aquatic predators out of their preferred habitat. *Proc. Natl Acad. Sci. USA* **114**, 9764–9765 (2017).
- Schmitz, O. J., Beckerman, A. P. & O'Brien, K. M. Behaviorally mediated trophic cascades: effects of predation risk on food web interactions. *Ecology* **78**, 1388–1399 (1997).
- Huey, R. B. & Kingsolver, J. G. Evolution of thermal sensitivity of ectotherm performance. *Trends Ecol. Evol.* **4**, 131–135 (1989).
- Allen, C. & Holling, C. S. *Discontinuities in Ecosystems and Other Complex Systems* (Columbia Univ. Press, New York, 2008).
- Tilman, D., Cassman, K. G., Matson, P. A., Naylor, R. & Polasky, S. Agricultural sustainability and intensive production practices. *Nature* **418**, 671–677 (2002).
- Butler, S. J., Vickery, J. A. & Norris, K. Farmland biodiversity and the footprint of agriculture. *Science* **315**, 381–384 (2007).
- Field, C. B., Barros, V., Stocker, T. F. & Dahe, Q. (eds) *Managing the Risks of Extreme Events and Disasters to Advance Climate Change Adaptation: Special Report of the Intergovernmental Panel on Climate Change* (Cambridge Univ. Press, Cambridge, UK, 2012).
- Flato, G. M. & Boer, G. J. Warming asymmetry in climate change simulations. *Geophys. Res. Lett.* **28**, 195–198 (2001).
- Xu, Y. & Ramanathan, V. Latitudinally asymmetric response of global surface temperature: implications for regional climate change. *Geophys. Res. Lett.* **39**, L13706 (2012).
- Chou, C., Tu, J.-Y. Y. & Tan, P.-H. H. Asymmetry of tropical precipitation change under global warming. *Geophys. Res. Lett.* **34**, L17708 (2007).
- Trenberth, K. E. & Hoar, T. J. El Niño and climate change. *Geophys. Res. Lett.* **24**, 3057–3060 (1997).
- Marshall, J. et al. The ocean's role in polar climate change: asymmetric Arctic and Antarctic responses to greenhouse gas and ozone forcing. *Phil. Trans. R. Soc. A* **372**, 20130040 (2014).
- Stouffer, R. J., Manabe, S. & Bryan, K. Interhemispheric asymmetry in climate response to a gradual increase of atmospheric CO₂. *Nature* **342**, 660–662 (1989).
- Sutton, R. T., Dong, B. & Gregory, J. M. Land/sea warming ratio in response to climate change: IPCC AR4 model results and comparison with observations. *Geophys. Res. Lett.* **34**, L02701 (2007).
- Karl, T. R. et al. Global warming: evidence for asymmetric diurnal temperature change. *Geophys. Res. Lett.* **18**, 2253–2256 (1991).
- Morelli, T. L. et al. Managing climate change refugia for climate adaptation. *PLoS ONE* **11**, e0159909 (2016).
- O'Reilly, C. M. et al. Rapid and highly variable warming of lake surface waters around the globe. *Geophys. Res. Lett.* **42**, 10773–10781 (2015).
- Kaarlejärvi, E., Eskelinen, A. & Olofsson, J. Herbivores rescue diversity in warming tundra by modulating trait-dependent species losses and gains. *Nat. Commun.* **8**, 419 (2017).
- Körner, C. *Alpine Treelines: Functional Ecology of the Global High Elevation Tree Limits*. (Springer, Basel, 2012).
- Crimmins, S. M., Dobrowski, S. Z., Greenberg, J. A., Abatzoglou, J. T. & Mynsberge, A. R. Changes in climatic water balance drive downhill shifts in plant species' optimum elevations. *Science* **331**, 324–327 (2011).
- Guzzo, M. M., Blanchfield, P. J. & Rennie, M. D. Behavioral responses to annual temperature variation alter the dominant energy pathway, growth, and condition of a cold-water predator. *Proc. Natl. Acad. Sci. USA* **114**, 9912–9917 (2017).
- Tunney, T. D., McCann, K. S., Lester, N. P. & Shuter, B. J. Effects of differential habitat warming on complex communities. *Proc. Natl Acad. Sci. USA* **111**, 8077–8082 (2014).
- Schmitz, O. J. & Barton, B. T. Climate change effects on behavioral and physiological ecology of predator–prey interactions: implications for conservation biological control. *Biol. Control* **75**, 87–96 (2014).
- Alofs, K. M., Jackson, D. A. & Lester, N. P. Ontario freshwater fishes demonstrate differing range-boundary shifts in a warming climate. *Divers. Distrib.* **20**, 123–136 (2014).
- Parmesan, C. et al. Poleward shifts in geographical ranges of butterfly species associated with regional warming. *Nature* **399**, 579–583 (1999).
- Menendez, R. et al. Species richness changes lag behind climate change. *Proc. R. Soc. B* **273**, 1465–1470 (2006).
- Wilson, R. J., Gutiérrez, D., Gutiérrez, J. & Monserrat, V. J. An elevational shift in butterfly species richness and composition accompanying recent climate change. *Glob. Change Biol.* **13**, 1873–1887 (2007).
- Kerr, J. T. et al. Climate change impacts on bumblebees across continents. *Science* **349**, 177–180 (2015).
- Lurgi, M., López, B. C. & Montoya, J. M. Climate change impacts on body size and food web structure on mountain ecosystems. *Phil. Trans. R. Soc. B* **367**, 3050–3057 (2012).
- Tilyanakis, J. M. & Morris, R. J. Ecological networks across environmental gradients. *Annu. Rev. Ecol. Syst.* **48**, 25–48 (2017).
- Travis, J. M. J. Climate change and habitat destruction: a deadly anthropogenic cocktail. *Proc. R. Soc. B* **270**, 467–473 (2003).
- Sax, D. F., Gaines, S. D. & Brown, J. H. Species invasions exceed extinctions on islands worldwide: a comparative study of plants and birds. *Am. Nat.* **160**, 766–783 (2002).
- Pires, M. M. Rewilding ecological communities and rewiring ecological networks. *Perspect. Ecol. Conserv.* **15**, 257–265 (2017).
- CaraDonna, P. J. et al. Interaction rewiring and the rapid turnover of plant–pollinator networks. *Ecol. Lett.* **20**, 385–394 (2017).
- Lu, X. et al. Drought rewires the cores of food webs. *Nat. Clim. Change* **6**, 875–878 (2016).
- May, R. M. *Stability and Complexity in Model Ecosystems* (Princeton Univ. Press, Princeton, NJ, USA, 1973).
- Yodzis, P. The connectance of real ecosystems. *Nature* **284**, 544–545 (1980).
- Pimm, S. The complexity and stability of ecosystems. *Nature* **307**, 321–326 (1984).
- Vergés, A. et al. The tropicalization of temperate marine ecosystems: climate-mediated changes in herbivory and community phase shifts. *Proc. R. Soc. B* **281**, 20140846 (2014).
- McCann, K. S. The diversity–stability debate. *Nature* **405**, 228–233 (2000).
- McCann, K., Hastings, A. & Huxel, G. R. Weak trophic interactions and the balance of nature. *Nature* **395**, 794–798 (1998).
- Allesina, S. & Tang, S. Stability criteria for complex ecosystems. *Nature* **483**, 205–208 (2012).

53. McCann, K. S. *Food Webs* (Princeton Univ. Press, Princeton, NJ, USA, 2011).
54. Neutel, A.-M. et al. Stability in real food webs: weak links in long loops. *Science* **296**, 1120–1123 (2002).
55. Yodzis, P. The stability of real ecosystems. *Nature* **289**, 674–676 (1981).
56. Lurgi, M., Lopez, B. C. & Montoya, J. M. Novel communities from climate change. *Phil. Trans. R. Soc. B* **367**, 2913–2922 (2012).
57. Humphries, M. M., Umbanhowar, J. & McCann, K. S. Bioenergetic prediction of climate change impacts on northern mammals. *Integr. Comp. Biol.* **44**, 152–162 (2004).
58. Velarde, E., Ezcurra, E. & Anderson, D. W. Seabird diets provide early warning of sardine fishery declines in the Gulf of California. *Sci. Rep.* **3**, 1332 (2013).
59. Cherry, M. J. & Barton, B. T. Effects of wind on predator-prey interactions. *Food Webs* **13**, 92–97 (2017).
60. Barton, B. T., Beckerman, A. P. & Schmitz, O. J. Climate warming strengthens indirect interactions in an old-field food web. *Ecology* **90**, 2346–2351 (2009).
61. Barton, B. T. Local adaptation to temperature conserves top-down control in a grassland food web. *Proc. R. Soc. B* **278**, 3102–3107 (2011).
62. Yurkowski, D. J. et al. Temporal shifts in intraguild predation pressure between beluga whales and Greenland halibut in a changing Arctic. *Biol. Lett.* **13**, 20170433 (2017).
63. Prop, J. et al. Climate change and the increasing impact of polar bears on bird populations. *Front. Ecol. Evol.* **3**, 33 (2015).
64. Hamilton, C. D., Kovacs, K. M., Ims, R. A., Aars, J. & Lydersen, C. An Arctic predator-prey system in flux: climate change impacts on coastal space use by polar bears and ringed seals. *J. Anim. Ecol.* **86**, 1054–1064 (2017).
65. Smith, P. A., Elliott, K. H., Gaston, A. J. & Gilchrist, H. G. Has early ice clearance increased predation on breeding birds by polar bears? *Polar Biol.* **33**, 1149–1153 (2010).
66. Dey, C. J. et al. Increasing nest predation will be insufficient to maintain polar bear body condition in the face of sea ice loss. *Glob. Change Biol.* **23**, 1821–1831 (2017).
67. Robinson, R. A. et al. Travelling through a warming world: climate change and migratory species. *Endanger. Species Res.* **7**, 87–99 (2009).
68. Deacy, W. W. et al. Phenological synchronization disrupts trophic interactions between Kodiak brown bears and salmon. *Proc. Natl Acad. Sci. USA* **114**, 10432–10437 (2017).
69. VanDerWal, J. et al. Focus on poleward shifts in species' distribution underestimates the fingerprint of climate change. *Nat. Clim. Change* **3**, 239–243 (2013).
70. Chen, I.-C., Hill, J. K., Ohlemüller, R., Roy, D. B. & Thomas, C. D. Rapid range shifts of species associated with high levels of climate warming. *Science* **333**, 1024–1026 (2011).
71. Siepielski, A. M. et al. Precipitation drives global variation in natural selection. *Science* **355**, 959–962 (2017).
72. Gilman, S. E., Urban, M. C., Tewksbury, J., Gilchrist, G. W. & Holt, R. D. A framework for community interactions under climate change. *Trends Ecol. Evol.* **25**, 325–331 (2010).
73. Peters, R. H. *The Ecological Implications of Body Size* (Cambridge Studies in Ecology, Cambridge Univ. Press, Cambridge, UK, 1983).
74. Sandstrom, S., Rawson, M. & Lester, N. *Manual of Instructions for Broad-Scale Fish Community Monitoring Using North American (NAI) and Ontario Small Mesh (ON2) Gillnets* (Queen's Printer for Ontario, 2013).
75. Dulvy, N. K. et al. Climate change and deepening of the North Sea fish assemblage: a biotic indicator of warming seas. *J. Appl. Ecol.* **45**, 1029–1039 (2008).
76. Sunday, J. M. et al. Species traits and climate velocity explain geographic range shifts in an ocean-warming hotspot. *Ecol. Lett.* **18**, 944–953 (2015).
77. Ling, S. D. et al. Global regime shift dynamics of catastrophic sea urchin overgrazing. *Phil. Trans. R. Soc. B* **370**, 20130269 (2014).
78. Overpeck, J. et al. Arctic environment change of the last four centuries. *Science* **278**, 1251–1257 (1997).
79. Clavel, J., Julliard, R. & Devictor, V. Worldwide decline of specialist species: toward a global functional homogenization? *Front. Ecol. Environ.* **9**, 222–228 (2011).
80. Vander Zanden, M. J., Casselman, J. M. & Rasmussen, J. B. Stable isotope evidence for the food web consequences of species invasions in lakes. *Nature* **401**, 464–467 (1999).
81. Rooney, N. & McCann, K. S. Integrating food web diversity, structure and stability. *Trends Ecol. Evol.* **27**, 40–45 (2012).
82. Cassman, K. G. et al. Ecological intensification of cereal production systems: yield potential, soil quality, and precision agriculture. *Proc. Natl Acad. Sci. USA* **96**, 5952–5959 (1999).
83. Fedoroff, N. V. et al. Radically rethinking agriculture for the 21st century. *Science* **327**, 833–834 (2010).
84. Tilman, D. et al. Forecasting agriculturally driven global environmental change. *Science* **292**, 281–284 (2001).
85. Eldredge, N. *Life in the Balance: Humanity and the Biodiversity Crisis* (Princeton Univ. Press, 2000).
86. Piao, S. et al. The impacts of climate change on water resources and agriculture in China. *Nature* **467**, 43–51 (2010).
87. Gilljam, D., Curtsdotter, A. & Ebenman, B. Adaptive rewiring aggravates the effects of species loss in ecosystems. *Nat. Commun.* **6**, 8412 (2015).
88. Pimm, S. L. & Lawton, J. H. Are food webs divided into compartments? *J. Anim. Ecol.* **49**, 879–898 (1980).
89. Moore, J. C. & Hunt, W. H. Resource compartmentation and the stability of real ecosystems. *Nature* **333**, 261–263 (1988).
90. Wootton, K. L. & Stouffer, D. B. Many weak interactions and few strong: food-web feasibility depends on the combination of the strength of species' interactions and their correct arrangement. *Theor. Ecol.* **9**, 185–195 (2016).
91. McCann, K. S., Rasmussen, J. B. & Umbanhowar, J. The dynamics of spatially coupled food webs. *Ecol. Lett.* **8**, 513–523 (2005).
92. Gilbert, B. et al. A bioenergetic framework for the temperature dependence of trophic interactions. *Ecol. Lett.* **17**, 902–914 (2014).
93. Scheffer, M. et al. Early-warning signals for critical transitions. *Nature* **461**, 53–59 (2009).
94. Boettiger, C., Ross, N. & Hastings, A. Early warning signals: the charted and uncharted territories. *Theor. Ecol.* **6**, 255–264 (2013).
95. Drake, J. M. & Griffen, B. D. Early warning signals of extinction in deteriorating environments. *Nature* **467**, 456–459 (2010).
96. Yodzis, P. & Innes, S. Body size and consumer-resource dynamics. *Am. Nat.* **139**, 1151–1175 (1992).
97. Kéfi, S. et al. Early warning signals of ecological transitions: methods for spatial patterns. *PLoS ONE* **9**, 10–13 (2014).
98. Oliver, T. H. et al. Biodiversity and resilience of ecosystem functions. *Trends Ecol. Evol.* **30**, 673–684 (2015).
99. Loreau, M. et al. Biodiversity and ecosystem functioning: current knowledge and future challenges. *Science* **294**, 804–808 (2001).
100. McMeans, B. C. et al. The adaptive capacity of lake food webs: from individuals to ecosystems. *Ecol. Monogr.* **86**, 4–19 (2016).
101. Aspillaga, E. et al. Thermal stratification drives movement of a coastal apex predator. *Sci. Rep.* **7**, 526 (2017).
102. Sato, C. F. & Lindenmayer, D. B. Meeting the global ecosystem collapse challenge. *Conserv. Lett.* **11**, e12348 (2017).
103. *GISS Surface Temperature Analysis (GISTEMP)* (NASA Goddard Institute for Space Studies, 2018); <https://data.giss.nasa.gov/gistemp/>
104. Hansen, J. et al. Global surface temperature change. *Rev. Geophys.* **48**, RG4004 (2010).
105. Barton, B. T. & Schmitz, O. J. Experimental warming transforms multiple predator effects in a grassland food web. *Ecol. Lett.* **12**, 1317–1325 (2009).
106. Bartley, T. J. *Flexible Food Web Structure in a Variable World*. PhD thesis, University of Guelph (2017).
107. Dolson, R., McCann, K., Rooney, N. & Ridgway, M. Lake morphometry predicts the degree of habitat coupling by a mobile predator. *Oikos* **118**, 1230–1238 (2009).
108. Tunney, T. D., McCann, K. S., Lester, N. P. & Shuter, B. J. Food web expansion and contraction in response to changing environmental conditions. *Nat. Commun.* **3**, 1105 (2012).
109. Morbey, Y. E., Couture, P., Busby, P. & Shuter, B. J. Physiological correlates of seasonal growth patterns in lake trout *Salvelinus namaycush*. *J. Fish Biol.* **77**, 2298–2314 (2010).
110. Guzzo, M. M. & Blanchfield, P. J. Climate change alters the quantity and phenology of habitat for lake trout (*Salvelinus namaycush*) in small Boreal Shield lakes. *Can. J. Fish. Aquat. Sci.* **74**, 871–884 (2017).
111. Keller, W. Implications of climate warming for Boreal Shield lakes: a review and synthesis. *Environ. Rev.* **15**, 99–112 (2007).
112. Adrian, R. et al. Lakes as sentinels of climate change. *Limnol. Oceanogr.* **54**, 2283–2297 (2009).
113. Barton, B. T. & Ives, A. R. Direct and indirect effects of warming on aphids, their predators, and ant mutualists. *Ecology* **95**, 1479–1484 (2014).
114. Barton, B. T. & Ives, A. R. Species interactions and a chain of indirect effects driven by reduced precipitation. *Ecology* **95**, 486–494 (2014).
115. Penczykowski, R. M., Connolly, B. M. & Barton, B. T. Winter is changing: trophic interactions under altered snow regimes. *Food Webs* **13**, 80–91 (2017).
116. Barton, B. T. Reduced wind strengthens top-down control of an insect herbivore. *Ecology* **95**, 2375–2381 (2014).
117. Barton, B. T. & Schmitz, O. J. Opposite effects of daytime and nighttime warming on top-down control of plant diversity. *Ecology* **99**, 13–20 (2018).

Acknowledgements

This project was in part funded by the University of Guelph's Canada First Research Excellence Fund project 'Food from Thought' awarded to K.S.M. and A.S.M. and a Discovery Grant from the National Science and Engineering Research Council of Canada

awarded to K.S.M. and A.S.M. T.J.B. was supported by a Canada Graduate Scholarship from the National Science and Engineering Research Council of Canada. We would like to thank the Ontario Ministry of Natural Resources and Forestry (OMNRF) and their Broad-Scale Fisheries Monitoring Program.

Author contributions

T.J.B. and K.S.M. conceived the concept for and contributed equally to this paper. All authors contributed to the development of the ideas and to the writing and editing of the text, led by T.J.B. and K.S.M. T.J.B. and M.G. prepared the figures using data from T.J.B., T.D.T. and M.M.G., as well as other sources. T.J.B. and K.S.M. led the final draft preparation and submission stages with comments from all authors being received prior to submission.

Competing interests

The authors declare no competing interests.

Additional information

Reprints and permissions information is available at www.nature.com/reprints.

Correspondence should be addressed to T.J.B.

Publisher's note: Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

© Springer Nature Limited 2019