

On the sensitivity of food webs to multiple disturbances

David Beauchesne^{1,2,}, Kevin Cazelles³, Philippe Archambault², Dominique Gravel⁴*

¹Institut des sciences de la mer, Université du Québec à Rimouski, Rimouski, QC, Canada

²Québec Océan, Département de biologie, Université Laval, Québec, QC, Canada

³Department of Integrative Biology, University Of Guelph, Guelph, Ontario, Canada N1G 2W1

⁴Département de biologie, Université de Sherbrooke, Sherbrooke, QC, Canada

Correspondence:

David Beauchesne

david.beauchesne@uqar.ca

1 Introduction

1.1 Context

- **Global changes, multiple stressors and food webs**

- Global changes are resulting in increasingly intricate environmental stress exposure regimes (Bowler et al., 2019; Côté et al., 2016; Halpern et al., 2015). These can in turn induce complex and unpredictable environmental effects that propagate through entire ecological communities by way of species interactions (Bascompte, 2009; Montoya et al., 2009).

- **Uncertainty associated with multiple stressors**

- Largest uncertainty in predicting environmental effects is the potential for complex driver interactions (Côté et al., 2016; Darling and Côté, 2008).
- Stressors can combine non-additively and result in effects that are greater (*i.e.* synergistic) or lower (*i.e.* antagonistic) than the sum of individual effects (Côté et al., 2016; Crain et al., 2008; Darling and Côté, 2008)
- Net effects of multiple stressors can be additive (*i.e.* joint effect equal to the sum of individual effects), synergistic (joint effect superior to the sum of individual effects), antagonistic (joint effect inferior to the sum of individual effects) or dominant (joint effect equal to an individual effect) (e.g. Crain et al., 2008; Côté et al., 2016; Darling and Côté, 2008).
- This is of particular significance for management, because we have mostly been operating under the assumption that stressors are mostly additive and thus can be managed independently.
- Maybe cite (Hodgson et al., 2019) (cite it for certain in the article).
- most research on driver effects in marine environments remains overwhelmingly focused on single driver assessments (O’Brien et al., 2019).

- **Limits of *in situ* and experimental approaches to study multiple disturbances**

- The number of stressors and of their potential interactions limits the insights we can glean into the effects of multiple disturbances *in situ* and in experimental settings such as mesocosm (Côté et al., 2016).
- In food webs, this is compounded by the myriad of possible interactions
- between species

- **Limited insights from null model testing**

- The knowledge we do have has mostly been gleaned from null model testing providing little insights into ecological mechanisms underlying non-additive stressor effects (De Laender, 2018).

- **Shift towards ecological modelling to better understand mechanisms**

- (De Laender, 2018; Schäfer and Piggott, 2018; Thompson et al., 2018a)

- **Some insights on the effects of disturbances on ecological communities from theoretical ecology**
 - Importance of interactions and web complexity
 - Indirect effects [Wootton (1993); Yodzis (2000); Wootton (2002); Montoya et al. (2009); ogorman2009]; see intro séminaire 1
 - Different types of interactions will lead to different overall effects (???).
 - * Different sensitivity for species involved in different types of interactions (segway to motifs)
 - How direct and indirect effects combine to affect food web sensitivity to disturbances
- **Theoretical has thus far mostly been concerned mostly with single disturbances and resistance of communities to extinctions**
 - Little insights into how different pathways of effect influences food webs, let alone pathways of multiple effects.
 - Little on the role of species and their interactions in propagating or buffering against disturbances

1.2 Objectives

1. Objective
2. Pathways of effect
3. Archetypes
4. Motifs
5. Terminology

- **State the objective**

- Here, we investigate the role of species in mediating the effects of multiple disturbances on food webs. We also seek to answer questions of particular significance for management: 1) should species interactions be considered in impact assessments, and 2) should the effects of stressors be evaluated separately or in combination?
- OR
- Here, we focus on evaluating the structural role of species and their trophic interactions in mediating food web sensitivity to multiple stressors. We also seek to answer questions of particular significance for management:
 - 1) should species interactions be considered in impact assessments, and
 - 2) should the effects of stressors be evaluated separately or in combination?

- **How we do this**

- To do so, we focus on how trophic position and interaction types affect species sensitivity to disturbances and the likelihood of species acting as buffer against or amplifiers of the effects of multiple disturbances.

91 • **Transition to motifs**

92 – ...?

93 • **Motif description and use**

- 94 – A food web can be decomposed into a set of smaller n -species subgraphs called
95 motifs (Milo et al., 2004; Stouffer et al., 2007). For example, there are 13 distinct
96 3-species motifs composed of 30 unique positions (Stouffer et al., 2007, 2012).
97 Motifs are the backbone of food webs and their study has unearthed valuable
98 insights on community dynamics such as [...].
- 99 – Their type and frequency has been linked to food web stability and persistence
100 (???)
- 101 – Like many disturbance studies in theoretical ecology, however, these have focused
102 less on the dynamics of population abundances and more on the resistance of food
103 webs to extinctions.
- 104 – Motifs have been used to investigate the persistence of food web to species extinc-
105 tions (Stouffer and Bascompte, 2010) and the benefit associated to each species
106 in food web persistence (Stouffer et al., 2012).
- 107 – *The ecological role of a species in a network is a direct rest if its interactions with*
108 *other species (Luczkovich 2003; Olesen 2007; Allesina 2009) (in Stouffer 2012)*
- 109 – *The number and types of motifs that make up a food web are known to directly*
110 *affect the web's stability and persistence (Neutel 2002; Kondoh 2008; Allesina*
111 *2008; García-Domingo 2008; Stouffer 2010; in Stouffer 2012)*

112 • **Our focus**

- 113 – Here we focus on the most abundant types of interactions: omnivory, tri-trophic
114 food chain, exploitative competition and apparent competition (Camacho et al.,
115 2007; Stouffer and Bascompte, 2010)
- 116 – Two additional motifs, *i.e.* partially connected and disconnected were also consid-
117 ered in order to evaluate whether interactions in food webs are truly more likely
118 to be characterized by non-linear effects.
- 119 – To study the effects of multiple stressors in a trophic context, we rather focus on
120 the resulting disturbances following different pathways of effect rather than on
121 the disturbances themselves. This means that we will not investigate the effects
122 of multiple stressors applied to a single species in the food web. This precludes us
123 from investigating the sensitivity of species to each individual stressor. Rather,
124 we investigate the effects of disturbances to multiple species simultaneously. But
125 see Thompson et al. (2018b) and Thompson et al. (2018a) for a description of
126 a modelling approach incorporating multiple sources of stress in a food web
127 model.
- 128 – We are interested in pathways that affect the trophic dynamic of food webs, *i.e.*
129 those that target population growth (*i.e.* birth and death rates) and the rates at
130 which species interact (*i.e.* attack and conversion rates).

131 • **Pathways of effect**

- Studying the effects of multiple disturbances means that we will be focusing on disturbances affecting multiple species, referred to as pathways of multiple effects ($D_{i,j}$, $D_{i,k}$, $D_{j,k}$, and $D_{i,j,k}$).
- In a food web context, we will define linear and non-linear effect as a function of whether these pathways of multiple effects result in additive ($D_{i,j} = D_i + D_j$), synergistic ($D_{i,j} \gg D_i + D_j$), antagonistic ($D_{i,j} \ll D_i + D_j$) or dominant ($D_{i,j} = D_i \vee D_j$) effects.

• **Terminology for types of pathways and position profile**

- Species can occupy different roles in these pathways of multiple effects. Investigating species profile (e.g.* Stouffer et al., 2012) could thus inform us on the role played by individual species in buffering against or amplifying the effects of multiple disturbances.
- We define 4 key roles in species propagating or buffering against multiple disturbances:

Role	Description	Mathematically	Example
Weak entry point	Species directly or indirectly more affected than expected	Synergistic effect on focal species	S_i in $\Delta i D_{i,j} \gg \Delta i D_i + \Delta i D_j$
Biotic multiplier	Species propagating disturbances	Peripheral species involved in synergistic effect	S_j in $\Delta i D_{i,j} \gg \Delta i D_i + \Delta i D_j$
Biotic sink	Species directly or indirectly affected less than expected	Antagonistic effect on focal species	S_i in $\Delta i D_{i,j} \ll \Delta i D_i + \Delta i D_j$
Biotic buffer	Species blocking the propagation of disturbances	Peripheral species involved in antagonistic effect	S_j in $\Delta i D_{i,j} \ll \Delta i D_i + \Delta i D_j$
Biotic dominant	Species whose individual effect dictates the response of other species	Focal or peripheral species driving dominance effect	S_j in $\Delta i D_{i,j} = \Delta i D_j$




Figure 1: Conceptualize the effects of multiple disturbances on food webs

2 Methodology

2.1 Models

The dynamics of the four most abundant 3-species motifs (*i.e.* tri-trophic food chain, omnivory, exploitative and apparent competition) in empirical food webs (Stouffer and Bascompte, 2010) were modeled using Lotka-Volterra equation systems (Table S1). Two additional motifs were included to serve as controls to test the importance of considering species interactions when evaluating environmental effects, *i.e.* a partially connected motifs with a disconnected resource and a predator and a prey interaction, and a fully disconnected motif with three independent species modelled as resources.

Resources were modeled using logistic growth equations of the form $\frac{dX_i}{dt} = X_i(r_i - \alpha_{ii}X_i - \sum \alpha_{ij}X_j)$, where X are species, i is the resource, j are the consumers, r_i is the intrinsic resource growth rate, α_{ii} is the density-dependent effect of the resource on itself and α_{ij} is the rate at which consumer j affects resource i , *i.e.* the attack rate.

Consumers and predators were modeled with a Type I functional response of the form $\frac{dX_j}{dt} = X_j(-m_j + \sum e_{ij}\alpha_{ij}X_i - \alpha_{jk}X_k)$, where m is the mortality rate and e is the rate at which resource biomass is transformed into consumer biomass, *i.e.* the conversion rate. e is a scaling parameter of the attack rate which cannot exceed 1, since 1 unit of resource biomass cannot be transformed into more than 1 unit of consumer biomass.

Models were solved at equilibrium since we wished to study the dynamics of the motifs in the context of persistent food webs, which is closer to reality. As no equilibrium exists for the exploitative competition motif with Lotka-Volterra models of these forms, competitive parameters of the form $\alpha_{jj}\alpha_{jk}X_jX_k - \alpha_{jj}X_j^2$ were included in the consumer models to constrain their growth and obtain equilibrium solutions. Sage (???) was used to solve all equation systems at equilibrium. All other analyses were performed using R (???). All code and data used to perform the simulations and analyses are available at [...].

Initial model parameter values for intrinsic growth (r) and resource density-dependence (α_{ii}) parameters were fixed to 1 and 0.001, respectively, to bound all solutions. Competitive parameters for the exploitative competition motif were also fixed at 0.001 since those parameters were not to be investigated in our analyses. Conversion rates (e) were fixed to 0.5. Finally, a total of 100 sets of mortality (m) and attack rates (α_{ij}) were evaluated using a simulated annealing algorithm optimizing for consumer abundance.

2.2 Disturbances

Parameters selected to simulate disturbances were those related to population growth (r and m) and interaction rates (e and α_{ij}). Disturbances were simulated by applying a 1% variation on all combinations of selected model parameters to simulate all possible unique pathways of univariate and multivariate effects. Parameter variation were only applied as a simulation of stress, hence mortality rates were increased, while all other parameters were decreased by 1%.

184 Parameter combinations resulted in between 7 (disconnected) and 511 (omnivory possible
185 pathways of effect, for a total of 930 unique pathways of effects (*to check, I may be multiplying*
186 *the number of unique pathways for exploitative and apparent competition*).

187 The sensitivity to disturbances was measured as percent change between the analytical abun-
188 dance after parameter variation and the initial abundance, for each species and each pathway
189 of effect. The amplification potential of pathways of multiple effects was measured as percent
190 difference between the analytical abundance of the multivariate disturbance model and the
191 sum of the univariate disturbance models. A value of 0 equals a null or additive effect, a
192 value below 0 is a dominant or antagonistic effect, and a value over 0 is a synergistic effect.




Figure 2: Disturbances

3 Results

3.1 Univariate disturbances

3.2 Multivariate disturbances

3.3 Additivity

4 Supplementary Material

4.1 Models

Motifs	Equation systems	Initial parameters values
Tri-trophic food chain	$\begin{aligned}\frac{dX_i}{dt} &= X_i(r_i - \alpha_{ii}X_i - \alpha_{ij}X_j) \\ \frac{dX_j}{dt} &= X_j(e_{ij}\alpha_{ij}X_i - \alpha_{jk}X_k - m_j) \\ \frac{dX_k}{dt} &= X_k(e_{jk}\alpha_{jk}X_j - m_k)\end{aligned}$	$\begin{aligned}r_i &= 1 \\ \alpha_{ii} &= 0.001 \\ \alpha_{ij}, \alpha_{jk} &\in [0.0001, 0.01] \\ e_{ij}, e_{jk} &= 0.5 \\ m_j, m_k &\in [0.01, 0.5]\end{aligned}$
Omnivory	$\begin{aligned}\frac{dX_i}{dt} &= X_i(r_i - \alpha_{ii} - \alpha_{ij}X_j - \alpha_{ik}X_k) \\ \frac{dX_j}{dt} &= X_j(e_{ij}\alpha_{ij}X_i - \alpha_{jk}X_k - m_j) \\ \frac{dX_k}{dt} &= X_k(e_{ik}\alpha_{ik}X_i + e_{jk}\alpha_{jk}X_j - m_k)\end{aligned}$	$\begin{aligned}r_i &= 1 \\ \alpha_{ii} &= 0.001 \\ \alpha_{ij}, \alpha_{ik}, \alpha_{jk} &\in [0.0001, 0.01] \\ e_{ij}, e_{ik}, e_{jk} &= 0.5 \\ m_j, m_k &\in [0.01, 0.5]\end{aligned}$
Exploitative competition	$\begin{aligned}\frac{dX_i}{dt} &= X_i(r_i - \alpha_{ii} - \alpha_{ij}X_j - \alpha_{ik}X_k) \\ \frac{dX_j}{dt} &= X_j(e_{ij}\alpha_{ij}X_i - \alpha_{jj}\alpha_{jk}X_k - \alpha_{jj}X_j - m_j) \\ \frac{dX_k}{dt} &= X_k(e_{ik}\alpha_{ik}X_i - \alpha_{kk}\alpha_{kj}X_j - \alpha_{kk}X_k - m_k)\end{aligned}$	$\begin{aligned}r_i &= 1 \\ \alpha_{ii}, \alpha_{jj}, \alpha_{kk}, \alpha_{jk}, \alpha_{kj} &= 0.001 \\ \alpha_{ij}, \alpha_{ik} &\in [0.0001, 0.01] \\ e_{ij}, e_{ik} &= 0.5 \\ m_j, m_k &\in [0.01, 0.5]\end{aligned}$
Apparent competition	$\begin{aligned}\frac{dX_i}{dt} &= X_i(r_i - \alpha_{ii}X_i - \alpha_{ik}X_k) \\ \frac{dX_j}{dt} &= X_j(r_j - \alpha_{jj}X_j - \alpha_{jk}X_k) \\ \frac{dX_k}{dt} &= X_k(e_{ik}\alpha_{ik}X_i + e_{jk}\alpha_{jk}X_j - m_k)\end{aligned}$	$\begin{aligned}r_i, r_j &= 1 \\ \alpha_{ii}, \alpha_{jj} &= 0.001 \\ \alpha_{ik}, \alpha_{jk} &\in [0.0001, 0.01] \\ e_{ik}, e_{jk} &= 0.5 \\ m_k &\in [0.01, 0.5]\end{aligned}$
Partially disconnected	$\begin{aligned}\frac{dX_i}{dt} &= X_i(r_i - \alpha_{ii}X_i - \alpha_{ik}X_k) \\ \frac{dX_j}{dt} &= X_j(r_j - \alpha_{jj}X_j) \\ \frac{dX_k}{dt} &= X_k(e_{ik}\alpha_{ik}X_i - m_k)\end{aligned}$	$\begin{aligned}r_i, r_j &= 1 \\ \alpha_{ii}, \alpha_{jj} &= 0.001 \\ \alpha_{ik} &\in [0.0001, 0.01] \\ e_{ik} &= 0.5 \\ m_k &\in [0.01, 0.5]\end{aligned}$
Disconnected	$\begin{aligned}\frac{dX_i}{dt} &= X_i(r_i - \alpha_{ii}X_i) \\ \frac{dX_j}{dt} &= X_j(r_j - \alpha_{jj}X_j) \\ \frac{dX_k}{dt} &= X_k(r_k - \alpha_{kk}X_k)\end{aligned}$	$\begin{aligned}r_i, r_j, r_k &= 1 \\ \alpha_{ii}, \alpha_{jj}, \alpha_{kk} &= 0.001\end{aligned}$

Table 2: Systems of Lotka-Volterra equations used to model the effect of multiple disturbances




Figure 3: Positions

5 Next points

- Non-linear effects in motifs
- Species contribution to non-linear effects
- Species profiles (frequency of times occupying roles that contribute to non-linear effects; see Stouffer et al. (2012))
- Graphs to present these results
- Methods

6 Notes - to explore

- With synergies, one should expect higher rates of community collapse, i.e. loss of species to extinction, because the effects increase more rapidly. Maybe look at Stouffer to better dissect their result on community persistence as a function of motifs. Perhaps they also explain which are the motif positions that tend to go extinct more rapidly?

7 Interesting points

- Effect limit (Schäfer and Piggott, 2018): maximum effect size for a response (*e.g.* 100% mortality, zero growth or reproduction)

8 Literature to cite - or at least look at!

- Adams (2005)
- Brown et al. (2013)
- Brown et al. (2014)
- Christensen et al. (2006)
- Crain et al. (2008)
- Darling et al. (2013)
- Folt et al. (1999)
- Galic et al. (2018) *
- Jackson et al. (2016)
- Kath et al. (2018)
- Lange et al. (2018)
- Piggott et al. (2015)
- Schäfer and Piggott (2018) *
- Segner et al. (2014)
- Thompson et al. (2018a)
- Thompson et al. (2018b)
- Vinebrooke et al. (2004)

9 References

- Adams, S. M. (2005). Assessing cause and effect of multiple stressors on marine systems. *Marine Pollution Bulletin* 51, 649–657.
- Bascompte, J. (2009). Disentangling the Web of Life. *Science* 325, 416–419. doi:10.1126/science.1170749.
- Bowler, D., Bjorkmann, A., Dornelas, M., Myers-Smith, I., Navarro, L., Niamir, A., et al. (2019). The geography of the Anthropocene differs between the land and the sea. *bioRxiv*. doi:10.1101/432880.
- Brown, C. J., Saunders, M. I., Possingham, H. P., and Richardson, A. J. (2013). Managing for interactions between local and global stressors of ecosystems. *PloS one* 8, e65765.
- Brown, C. J., Saunders, M. I., Possingham, H. P., and Richardson, A. J. (2014). Interactions between global and local stressors of ecosystems determine management effectiveness in cumulative impact mapping. *Diversity and Distributions* 20, 538–546. doi:10.1111/ddi.12159.
- Camacho, J., Stouffer, D. B., and Amaral, L. A. N. (2007). Quantitative analysis of the local structure of food webs. *Journal of Theoretical Biology* 246, 260–268. doi:10.1016/j.jtbi.2006.12.036.
- Christensen, M. R., Graham, M. D., Vinebrooke, R. D., Findlay, D. L., Paterson, M. J., and Turner, M. A. (2006). Multiple anthropogenic stressors cause ecological surprises in boreal lakes. *Global Change Biology* 12, 2316–2322.
- Côté, I. M., Darling, E. S., and Brown, C. J. (2016). Interactions among ecosystem stressors and their importance in conservation. *Proceedings of the Royal Society B: Biological Sciences* 283, 20152592. doi:10.1098/rspb.2015.2592.
- Crain, C. M., Kroeker, K., and Halpern, B. S. (2008). Interactive and cumulative effects of multiple human stressors in marine systems. *Ecology Letters* 11, 1304–1315. doi:10.1111/j.1461-0248.2008.01253.x.
- Darling, E. S., and Côté, I. M. (2008). Quantifying the evidence for ecological synergies. *Ecology Letters* 11, 1278–1286. doi:10.1111/j.1461-0248.2008.01243.x.
- Darling, E. S., McClanahan, T. R., and Côté, I. M. (2013). Life histories predict coral community disassembly under multiple stressors. *Global Change Biology* 19, 1930–1940. doi:10.1111/gcb.12191.
- De Laender, F. (2018). Community-and ecosystem-level effects of multiple environmental change drivers: Beyond null model testing. *Global change biology* 24, 5021–5030.
- Folt, C. L., Chen, C. Y., Moore, M. V., and Burnaford, J. (1999). Synergism and antagonism among multiple stressors. *Limnology and oceanography* 44, 864–877.
- Galic, N., Sullivan, L. L., Grimm, V., and Forbes, V. E. (2018). When things don’t add up: Quantifying impacts of multiple stressors from individual metabolism to ecosystem processing. *Ecology Letters* 21, 568–577. doi:10.1111/ele.12923.

- Halpern, B. S., Frazier, M., Potapenko, J., Casey, K. S., Koenig, K., Longo, C., et al. (2015). Spatial and temporal changes in cumulative human impacts on the world's ocean. *Nature Communications* 6. doi:10.1038/ncomms8615.
- Hodgson, E. E., Halpern, B. S., and Essington, T. E. (2019). Moving Beyond Silos in Cumulative Effects Assessment. *Frontiers in Ecology and Evolution* 7. doi:10.3389/fevo.2019.00211.
- Jackson, M. C., Loewen, C. J. G., Vinebrooke, R. D., and Chimimba, C. T. (2016). Net effects of multiple stressors in freshwater ecosystems: A meta-analysis. *Global Change Biology* 22, 180–189. doi:10.1111/gcb.13028.
- Kath, J., Thomson, J. R., Thompson, R. M., Kefford, B. J., Dyer, F. J., and Mac Nally, R. (2018). Interactions among stressors may be weak: Implications for management of freshwater macroinvertebrate communities. *Diversity and Distributions* 24, 939–950.
- Lange, K., Bruder, A., Matthaei, C. D., Brodersen, J., and Paterson, R. A. (2018). Multiple-stressor effects on freshwater fish: Importance of taxonomy and life stage. *Fish and Fisheries* 19, 974–983.
- Milo, R., Itzkovitz, S., Kashtan, N., Levitt, R., Shen-Orr, S., Ayzenshtat, I., et al. (2004). Superfamilies of Evolved and Designed Networks. *Science* 303, 1538–1542. doi:10.1126/science.1089167.
- Montoya, J., Woodward, G., Emmerson, M. C., and Solé, R. V. (2009). Press perturbations and indirect effects in real food webs. *Ecology* 90, 2426–2433. doi:10.1890/08-0657.1.
- O'Brien, A. L., Dafforn, K. A., Chariton, A. A., Johnston, E. L., and Mayer-Pinto, M. (2019). After decades of stressor research in urban estuarine ecosystems the focus is still on single stressors: A systematic literature review and meta-analysis. *Science of The Total Environment*. doi:10.1016/j.scitotenv.2019.02.131.
- Piggott, J. J., Townsend, C. R., and Matthaei, C. D. (2015). Reconceptualizing synergism and antagonism among multiple stressors. *Ecology and Evolution* 5, 1538–1547. doi:10.1002/ece3.1465.
- Schäfer, R. B., and Piggott, J. J. (2018). Advancing understanding and prediction in multiple stressor research through a mechanistic basis for null models. *Global Change Biology* 24, 1817–1826. doi:10.1111/gcb.14073.
- Segner, H., Schmitt-Jansen, M., and Sabater, S. (2014). *Assessing the impact of multiple stressors on aquatic biota: The receptor's side matters*. ACS Publications.
- Stouffer, D. B., and Bascompte, J. (2010). Understanding food-web persistence from local to global scales. *Ecology Letters* 13, 154–161. doi:10.1111/j.1461-0248.2009.01407.x.
- Stouffer, D. B., Camacho, J., Jiang, W., and Nunes Amaral, L. A. (2007). Evidence for the existence of a robust pattern of prey selection in food webs. *Proceedings of the Royal Society B: Biological Sciences* 274, 1931–1940. doi:10.1098/rspb.2007.0571.
- Stouffer, D. B., Sales-Pardo, M., Sirer, M. I., and Bascompte, J. (2012). Evolutionary Conservation of Species' Roles in Food Webs. *Science* 335, 1489–1492. doi:10.1126/science.1216556.

309 Thompson, P. L., MacLennan, M. M., and Vinebrooke, R. D. (2018a). An improved null
 310 model for assessing the net effects of multiple stressors on communities. *Global change biology*
 311 24, 517–525.

312 Thompson, P. L., MacLennan, M. M., and Vinebrooke, R. D. (2018b). Species interactions
 313 cause non-additive effects of multiple environmental stressors on communities. *Ecosphere* 9,
 314 e02518.

315 Vinebrooke, R. D., Cottingham, K. L., Scheffer, J. N., Marten, Dodson, S. I., Maberly,
 316 S. C., and Sommer, U. (2004). Impacts of multiple stressors on biodiversity and ecosys-
 317 tem functioning: The role of species co-tolerance. *Oikos* 104, 451–457. doi:10.1111/j.0030-
 318 1299.2004.13255.x.

319 Wootton, J. T. (1993). Indirect Effects and Habitat Use in an Intertidal Community:
 320 Interaction Chains and Interaction Modifications. *The American Naturalist* 141, 71–89.
 321 doi:10.1086/285461.

322 Wootton, J. T. (2002). Indirect effects in complex ecosystems: Recent progress and future
 323 challenges. *Journal of Sea Research* 48, 157–172. doi:10.1016/S1385-1101(02)00149-1.

324 Yodzis, P. (2000). Diffuse Effects in Food Webs. *Ecology* 81, 261–266. doi:10.1890/0012-
 325 9658(2000)081[0261:DEIFW]2.0.CO;2.