


available at www.sciencedirect.comjournal homepage: www.elsevier.com/locate/ecolmodel

The food web structure and ecosystem properties of a filter-feeding carps dominated deep reservoir ecosystem

Qi-Gen Liu^{a,b}, Yong Chen^c, Jia-Le Li^a, Li-Qiao Chen^{b,*}

^a The Key Laboratory of Aquatic Genetic Resources and Aquacultural Ecology (AGRA), The Ministry of Agriculture, Shanghai Fisheries University, Shanghai 200090, PR China

^b College of Life Science, East China Normal University, Shanghai 200062, PR China

^c School of Marine Sciences, University of Maine, Orono, ME 04469, USA

ARTICLE INFO

Article history:

Received 4 January 2006

Received in revised form

23 November 2006

Accepted 27 November 2006

Published on line 8 January 2007

Keywords:

Ecopath model

Hypophthalmichthys molitrix

Aristichthys nobilis

Stocking

Ecosystem properties

Trophic level

Trophic flow

Throughput

Lake Qiandaohu

ABSTRACT

An Ecopath model was constructed to describe the ecosystem of Lake Qiandaohu, a stock-enhanced large deep Chinese reservoir with silver carp (*Hypophthalmichthys molitrix*) and bighead carp (*Aristichthys nobilis*) dominated in its pelagic community. The food web structure and ecosystem property of the reservoir were analyzed and evaluated. The results showed that there were seven trophic levels (TLs) in the system, with the trophic flows primarily occurring through the first four TLs. The food web structure of this ecosystem was characterized with a bulged intermediate trophic level, which was contrary to the wasp-waist food web structure occurred in most natural aquatic ecosystems. The corresponding trophic flow pattern showing by transfer efficiencies (TEs) between TLs indicated that the trophic flows primarily went through from TL I to II with a high TE (of over 50%) and through a flow loop or short cut between detritus and TL II but greatly reduced from TL II to III with a lowest TE of 2.5% due to the bulged biomass at TL II. The trophic flow loop greatly increased the throughput recycled, which, together with high connectance index (CI), system omnivory index (SOI), Finn's cyclic index (FCI) and Finn's mean path length (FML), might be beneficial to the maintaining of ecosystem stability. Finally, ecosystem property indices showed that this reservoir had a high value of P_p/R and P_p/B , indicating this ecosystem of short history was immature, but highly productive. This silver carp and bighead carp dominated deep reservoir ecosystem had both the characteristics of high productivity of an immature ecosystem and the feature of high stability of a mature ecosystem.

© 2006 Elsevier B.V. All rights reserved.

1. Introduction

Food web structure and interactions have a decisive role in determining the dynamics of an ecosystem, and are of interests of many ecological studies (e.g., Halfon et al., 1996; Elser et al., 1998; Kitchell et al., 2000; Aoki and Mizushima, 2001; Angelini and Agostinho, 2005; Gamito and Erzini, 2005). Lakes and reservoirs in China are particularly in need of the

study of the food webs for the following two reasons: (1) the food webs of these ecosystems are usually characterized with the dominance of the two artificially stocked filter-feeding planktivorous Chinese carps, silver carp (*Hypophthalmichthys molitrix*) and bighead carp (*Aristichthys nobilis*), and their ecological effects on aquatic ecosystems have been controversial and contradictive (e.g., Januszko, 1974; Kajak et al., 1975; Opuszynski, 1979; Smith, 1985, 1989; Spataru and Gophen,

* Corresponding author. Tel.: +86 21 62233637.

E-mail addresses: qgliu@shfu.edu.cn (Q.-G. Liu), lqchen@bio.ecnu.edu.cn (L.-Q. Chen).
0304-3800/\$ – see front matter © 2006 Elsevier B.V. All rights reserved.
doi:10.1016/j.ecolmodel.2006.11.028