



British Ecological Society

---

On the Fauna of a Brown Flux of an Elm Tree, *Ulmus procera* Salisb.

Author(s): I. Robinson

Source: *Journal of Animal Ecology*, Vol. 22, No. 1 (May, 1953), pp. 149-153

Published by: [British Ecological Society](#)

Stable URL: <http://www.jstor.org/stable/1697>

Accessed: 31/03/2011 01:20

---

Your use of the JSTOR archive indicates your acceptance of JSTOR's Terms and Conditions of Use, available at <http://www.jstor.org/page/info/about/policies/terms.jsp>. JSTOR's Terms and Conditions of Use provides, in part, that unless you have obtained prior permission, you may not download an entire issue of a journal or multiple copies of articles, and you may use content in the JSTOR archive only for your personal, non-commercial use.

Please contact the publisher regarding any further use of this work. Publisher contact information may be obtained at <http://www.jstor.org/action/showPublisher?publisherCode=briteco>.

Each copy of any part of a JSTOR transmission must contain the same copyright notice that appears on the screen or printed page of such transmission.

JSTOR is a not-for-profit service that helps scholars, researchers, and students discover, use, and build upon a wide range of content in a trusted digital archive. We use information technology and tools to increase productivity and facilitate new forms of scholarship. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).



*British Ecological Society* is collaborating with JSTOR to digitize, preserve and extend access to *Journal of Animal Ecology*.

<http://www.jstor.org>

# ON THE FAUNA OF A BROWN FLUX OF AN ELM TREE, *ULMUS PROCERA* SALISB.

BY I. ROBINSON

*Department of Zoology, University of Sheffield*

(With 2 Figures in the Text)

The following observations were made on an elm tree brown flux in Sheffield over a period of about two years. During this time the flux had a continuous existence and carried a fauna which formed a balanced association. The purpose of the investigations was to throw light on the ingredients of this association, its seasonal fluctuations and the interdependence of its members.

I take pleasure in thanking Prof. L. E. S. Eastham for suggesting the problem and for help during the investigation. It is pleasant also to thank Dr H. Oldroyd, Dr F. van Emden and Dr P. Freeman of the British Museum (Natural History) for identifying the flux-visiting Diptera.

## THE FLUX

Records of brown fluxes and their faunas have previously been made by Ludwig (1888); Massee (1897); Ogilvie (1924) and Keilin (1921, 1927). The flux here under consideration owes its existence to an out-pouring of sap from the wound of a branch scar at the height of about 6 ft. above ground-level. The sap oozes continually, though less copiously in winter than in spring and summer. In so doing, it forms a stream over the tree bark varying in width from 9 in. to 1 ft. In this position the substance of the flux develops as a spongy mass which holds the sap emerging from the tree and thus provides a stable micro-aquatic environment to which certain organisms are attracted and in which they live and often breed.

In spring and early summer the flux is dark brown and slimy. In winter, on the other hand, the lower parts are dry and light stone-coloured—features in agreement with the findings of Ogilvie (1924). The pH of the flux material was 8.7 throughout the year. Ogilvie (1924) records a pH of a similar elm flux as 9.0–9.5. Standard tests for starches, sugars, fats and proteins yielded negative results though a polysaccharide, not starch and possibly cellulose, was indicated.

The animals inhabiting the brown flux form a characteristic fauna which is constantly undergoing change throughout the year. The members of this fauna are:

(a) The mite *Hericia hericia* Kramer (Tyroglyphidae).

(b) The larvae of the following Diptera: *Dasyhelea obscura* Winnertz (Ceratopogonidae), *Aulacigaster leucopeza* (Meigen) (Aulacigasteridae), *Brachyopa insensilis* Collin (Syrphidae), two undetermined species of *Systemus* (Dolichopodidae), one undetermined species of *Phaonia* (Anthomyidae).

(c) The Staphylinid beetle *Thamiarea hospita* Maerk.