

A QCNN for Quantum State Preparation

Carnegie Vacation Scholarship

David Amorim

Week 5
(29/07/2024 - 02/08/2024)

Table of Contents

① Preliminaries

② Improving the Loss Function

③ Next Steps

Aims for the Week

The following aims were set at the last meeting (29/07/2024):

Improve Loss Function

Work on an improved version of WILL. Incorporate some phase extraction metrics (e.g. χ , ϵ) into the loss function.

Investigate Phase Extraction

Study the relationship between mismatch and the extracted phase, i.e. study the operator $\tilde{Q}^\dagger(\hat{I} \otimes \hat{R})\tilde{Q}$.

Mitigate Barren Plateaus

Work on strategies to mitigate barren plateaus, e.g. implement layer-by-layer training.

THIS WEEK INCLUDE EXAMPLES WITH HIGH m !!
KEEP WORKING ON CHIL!! MAYBE RENAME TO QRQ??_v DOES
NOT WORK !!! LLLL

Table of Contents

① Preliminaries

② Improving the Loss Function

③ Next Steps

WILL Revisited

- As discussed at the meeting on 29/07, the definition of **WILL** (weighted L_p loss) was amended to:

$$\text{WILL}_{p,q} = \left(\sum_k \left| x_k - y_k \right|^p + |x_k| \left| [k]_m - \Psi([k]_n) \right|^q \right)^{1/p}, \quad (1)$$

where the changes to the previous definition are highlighted

- Testing this for different Ψ (with $L = 6$, $m = 3$ and 600 epochs) yielded the following optimal values for p , q :

$\Psi(f)$	p	q
$\sim f$	0.25	0.5
$\sim f^2$	1	1.5
Ψ_{H23}	0.75	2

Table 1: Optimal identified p , q values for WILL

Comparing SAM, WIM, and WILL

	SAM	WIM	WILL
μ	3.4e-2	6.0e-2	4.5e-1
σ	1.4e-1	1.1e-1	4.7e-1
ϵ	1.9e-2	9.2e-2	2.6e-1
χ	3.2e-2	5.1e-2	3.7e-1
Ω	4.46	3.19	0.76

Table 2: Comparing loss function metrics for $\Psi(f) \sim f$ ($L = 6$, $m = 3$, 600 epochs)




Figure 1: Comparing extracted phase functions for $\Psi(f) \sim f$ ($L = 6$, $m = 3$, 600 epochs)

Comparing SAM, WIM, and WILL

	SAM	WIM	WILL
μ	1.9e-1	2.3e-1	6.6e-1
σ	1.2e-1	1.0e-1	4.1e-1
ϵ	2.2e-1	4.2e-1	2.8e-2
χ	1.9e-1	2.0e-1	6.1e-1
Ω	1.39	1.05	0.57

Table 3: Comparing loss function metrics for $\Psi(f) \sim f^2$ ($L = 6$, $m = 3$, 600 epochs)




Figure 2: Comparing extracted phase functions for $\Psi(f) \sim f^2$ ($L = 6$, $m = 3$, 600 epochs)

Comparing SAM, WIM, and WILL

	SAM	WIM	WILL
μ	6.8e-2	8.4e-2	7.6e-2
σ	1.8e-1	1.2e-1	2.6e-1
ϵ	4.5e-2	1.8e-1	7.3e-3
χ	7.4e-2	1.0e-1	6.2e-2
Ω	2.75	2.07	2.48

Table 4: Comparing loss function metrics for Ψ_{H23} ($L = 6$, $m = 3$, 600 epochs)




Figure 3: Comparing extracted phase functions for Ψ_{H23} ($L = 6$, $m = 3$, 600 epochs)

Table of Contents

① Preliminaries

② Improving the Loss Function

③ Next Steps

Next Steps

- ...