

UNIMODALITY IDEAS

AARON LANDESMAN

1. DIRECTIONS TO MOVE

- (1) Look at generalising p_i^r for general r .
- (2) Generalizing to q analog of cyclic group.
- (3) Try relating p_i, q_i .
- (4) Coding which groups G we have $p_i = q_i$.
- (5) When are $p_i = q_i$.
- (6) Try to compute q_i .
- (7) Look at simple groups, and maybe solvable groups, try quotienting by normal subgroups?
- (8) Are there any ways to combine G_1, G_2 where G_i are groups with $p_i = q_i$.
- (9) Are there some characterisations of groups with q_i, p_i .
- (10) How to use sage, what can we do with groups?

2. CYCLIC GROUP EDGES, THIS SECTION IS WRONG.

Remark 2.0.1. All subscripts will be taken $(\text{mod } n)$.

Theorem 2.0.2. The statistic p_i as Zijian defined are unimodal for the necklace poset.

Proof.

□

Lemma 2.0.3. $q_i = \binom{n-1}{i-1}$.

Proof.

□

Lemma 2.0.4. The difference $p_i - q_i$ is the number of pairs $(x, y), x \leq y$, such that there exists σ so that $\sigma x \leq y$, but there does not exist g with $gx = \sigma x, gy = y$.

Proof.

□

Definition 2.0.5. We call (x, y) a special pair if there exists σ so that $\sigma x \leq y$, but there does not exist g with $gx = \sigma x, gy = y$.

Lemma 2.0.6. $p_i - q_i$ is the number of orbits of special pairs

Proof.

□

Remark 2.0.7. We are restricting to the cyclic group C_n , and so we are assuming that all elements are generated by the permutation $c = (12 \cdots n)$. We now wish to bound the number of orbits of special pairs.

Remark 2.0.8. To compute the number of orbits of special pairs, we can always assume $x = \{t_1, t_2, \dots, t_{i-1}\}_<$ and $y = \{t_1, t_2, \dots, t_i\}_<$ by simply composing with an element of C_n in order to make the missing t_i the biggest element of the set. We may as well assume $t_1 = 1$.

Lemma 2.0.9. *Suppose (x, y) is a special pair, as in ?? with $\sigma x \leq y$. Then, σ is the permutation sending t_1 to t_2 or t_1 to t_i .*

Proof. Suppose otherwise, that it sent t_1 to t_k for $k \neq 2, i$. Then, since the relative ordering of the elements are preserved, we must have t_l is sent to t_{l+k-1} . Since we can only act by elements of C_n , this gives us that the values $t_{l+1} - t_l = t_{k+l} - t_{k+l-1}$ for all $l \in [i]$. However, this means that $\sigma y = y$, (THAT IS THE TRICKIEST PART TO SEE) so x, y is not a special pair. \square

Lemma 2.0.10. *Any special pair (x, y) must have $y = \{t_1, t_1 + a, t_1 + 2a, \dots, t_1 + (i-1)a\}$ for some $a < n$.*

Proof. By ?? we must have that σ sends t_1 to t_2 or t_i . Let's assume it sends t_1 to t_2 , the other case is similar. However, this means we must send t_l to t_{l+1} , which means $t_l - t_{l-1} = t_{l+1} - t_l$, which means that y is of the claimed form. \square

Lemma 2.0.11. *The number of orbits of special pairs is between $n-1$ and $\frac{n}{2}-1$.*

Proof. By the above, there are at most n orbits (as determined by the value of a) which can have special pairs. However, it is clear that $a = i$ and $a = n - i$ lie in the same orbit. Therefore, we have $\frac{n}{2}$ identifications, which tells us $p_i - q_i \geq \frac{n}{2} - 1$. On the other hand, in the worst possible case, we have all n orbits are equivalent, which implies $p_i - q_i \leq n$. \square

Lemma 2.0.12. *The p_i statistics are unimodal.*

Proof. By ?? we know the differences of the q_i on the nose it is obvious that $q_i - q_{i-1} \geq \frac{n}{2}$. However, by ?? we have $n-1 \geq (q_i - p_i) \geq \frac{n}{2} - 1$. Therefore, $p_i - p_{i-1} \geq \frac{n}{2} - \frac{n}{2} \geq 0$. \square

3. IDEAS FROM MONDAY JUNE 30

Let $S_{x,y} = \{\sigma | \sigma x \leq y\}$. I think $q_i = \sum_{G(x,y)} \frac{|S_{x,y}|}{|Stab(x) \cup Stab(y)|}$

Now, let us think of σ as a representative of $S_{x,y}/(Stab_x \cup Stab_y)$. Note, the condition that σ does not stabilize x means we need the all the vertices of y to be present in $x \cup \sigma x$.

Possible idea, we might still have to always send $i_1 \mapsto i_n$? Not so sure about how to make this precise though.

Lemma 3.0.13. *$Stab(x) \neq e \implies Stab(y) = e$, and $Stab(x) \neq e \implies Stab(y) = e$,*

Proof. If $c^k \in Stab(x)$, $c^k \neq e$, where c is the cyclic generator, and k is minimal. We know that every elements in x spaced every k apart. So, we can treat it as a necklace of length k . We just have to show that after adding a single bead it cannot have any symmetries. However, if we could do this in a relatively prime fashion, we would obtain many additional equalities between differences of places in x , and we could not change just one of those differences due to cyclic symmetry of order k , since we would not get an equality in one of the n/k parts, but we would get the equality in all the others. \square

Lemma 3.0.14. *In order for $|\frac{|S_{x,y}|}{|Stab(x) \cup Stab(y)|}| > 1$, we must have $Stab(x) \cup Stab(y) = e$.*

Proof. This is similar to the previous lemma, in either case, we get equalities between differences of the one with a stabilizer, which end up contradicting the one additional point. \square

Lemma 3.0.15. *In order to have $|\frac{|S_{x,y}|}{|Stab(x) \cup Stab(y)|}| > 1$, x must be of the form of a cyclic element union 1 dude, and y must be of the form of the cyclic element union 2 dudes, and the rotation must take x to y .*

Proof. \square

Lemma 3.0.16. *Now we just need to count the number of cyclic generator dudes. Clearly at most n , since the number of divisors are less than n . Therefore, since each contributes at most n , we only have to bound successive differences of binomial coefficients by n^2 .*

Proof. For each prime divisor, we can obtain one such cyclic generator dude. First, there are the ones which all have the same difference, these seem to be most common. There are exactly $n/2-1$ of these? The other counts will depend on the orbit sizes. \square

4. PROOF OF CYCLIC GROUP EDGE NUMBERS IS UNIMODAL

Lemma 4.0.17. *The number $q_i - p_i$ is in bijection with the number of orbits with representative y satisfying the following properties.*

- (1) *There is some $k|n$, possibly with $k = n$, such that there are $\lfloor \frac{i}{k} \rfloor$ distinct sets of the form $\{a + k, \dots, a + nk\}$, taken $(\text{mod } n)$*
- (2) *The remaining $s = i - \lfloor \frac{i}{k} \rfloor$, with $s \neq 1, s \neq k$. elements all have the same value $(\text{mod } \frac{1}{n}k)$ and there exists $a \in [n]$ and $r < \frac{n}{2}$ such that the remaining elements all lie in the set $\{a + r, a + 2r, \dots, a + sr\}$.*

Proof. Split by the cases of $k = n$, in which case we have a full cycle, and this is clearly the only possibility, or otherwise $k \neq n$, in which case some set of elements must be fixed, and then we will have such a “tail cycle” as in the second property \square

Definition 4.0.18. *We call the non full cycle, as described in the second property a tail cycle.*

Remark 4.0.19. *The goal of this section is to bound $q_{i-1} < p_i \leq q_i$. Note we trivially know $p_i \leq q_i$, Additionally, this bound will trivially show the p_i are unimodal (they are obviously symmetric by taking complements.) Therefore, to accomplish our goal, it suffices to show $q_i - p_i > q_i - q_{i-1} = \binom{n-2}{i-1}$, since we know $q_i = \binom{n-i}{i-1}$.*

Notation 4.0.20. *Denote the number of edges from $(B_n/G)_i$ to $(B_n/G)_{i-1}$ by $p(n, i)$. Similarly, define $q(n, i)$ as the q_i for B_n .*

Lemma 4.0.21.

$$p_i \leq \sum_{k=\frac{n}{p}, p|k, p \text{ prime}} p(k, \lceil \frac{i}{k} \rceil) \lfloor \frac{p}{2} \rfloor$$

Proof. Note that by ??, we only have to count the number of G -orbits satisfying the condition of ??. Clearly, for every $t|n$, we can write t as a factor of some $k = \frac{n}{p}$. Therefore, for any cycle size t , it can be decomposed as a disjoint union of $\frac{k}{t}$ cycles

of size k . Therefore, summing the number of equivalence classes corresponding to elements of $p_i - q_i$, overcounts the number of equivalence classes.

Now, I claim the number of equivalence classes composed of cycles of length k is $p(k, \lceil \frac{i}{k} \rceil) \lfloor \frac{k}{2} \rfloor$. To see this, we describe a simple bijection. We know that the tail cycle has between 2 and $k-1$ elements. Now, restrict to only the first k elements of the cycle. Each representative has a set of $\lfloor \frac{i}{k} \rfloor$ full cycles, and 1 partial cycle. For each such element, we correspond all $\lceil \frac{i}{k} \rceil$ to an element $y_k \in B_k$, and the elements of $\lfloor \frac{i}{k} \rfloor$ with an element x_k of B_k . Clearly $x < y$. This uniquely characterizes the locations of the full cycles $(\text{mod } k)$, and the location of the partial cycle. Hence, there are $p(k, \lceil \frac{i}{k} \rceil)$. The reason we have to add in a factor of $\lfloor \frac{k}{2} \rfloor$ is simply to tell us the difference between two adjacent element of the cycle in x_k . It can be any multiples of k from k to $k \cdot p$, but of course a difference of $k \cdot i$ is equivalent to $k \cdot p - i$, and these are the only two equivalent cycles. Hence, this proves the bound. \square

Theorem 4.0.22. *We can bound $p_i \leq q_i - q_{i-1}$*

Proof. Using the above relation, we can bound $p_i \leq \sum_{k=\frac{n}{p}, p|k, p \text{ prime}} p(k, \lceil \frac{i}{k} \rceil) \lfloor \frac{k}{2} \rfloor$.

We know $p(n, i) \leq q(n, i)$, and so using this bound, we have $p_i \leq \sum_{k=\frac{n}{p}, p|k, p \text{ prime}} q(k, \lceil \frac{i}{k} \rceil) \lfloor \frac{k}{2} \rfloor$,

and then since we know $q(k, \lceil \frac{i}{k} \rceil) = \binom{k-1}{\lceil \frac{i}{k} \rceil - 1}$, we can write $p_i \leq \sum_{k=\frac{n}{p}, p|k, p \text{ prime}} \binom{k-1}{\lceil \frac{i}{k} \rceil - 1} \lfloor \frac{k}{2} \rfloor$.

Now, there are clearly at most $\log n$ terms in this sum, since there are fewer than $\log n$ distinct prime factors of n . And therefore, for all $k|n, k > 3$, we just have to bound $\log n \cdot \binom{k-1}{\lceil \frac{i}{k} \rceil - 1} \lfloor \frac{k}{2} \rfloor \cdot \log n < \binom{n-2}{i-1}$, which is quite obvious, at least for $n > 5$ or so. \square