

Vectors in Space

OpenStax Calculus Vol. 3

2.1 Vectors in the Plane

A vector in a plane is represented by a directed line segment, its endpoints are called the initial and terminal points respectively. An arrow from the initial point to the terminal point indicates the direction of the vector and the length of the line segment represents its magnitude. The notation $||\vec{v}||$ or $||\mathbf{v}||$ is used to denote the magnitude of the vector \vec{v} (or \mathbf{v}). The zero vector is a vector with the same initial and terminal point denoted $\vec{0}$ or $\mathbf{0}$.

Vectors with the same magnitude and direction are considered equivalent vectors and are treated as equal even if they have different initial points. Two vectors are parallel if they have the same or opposite directions. Vectors are defined by magnitude and direction regardless of the location of the initial point.


Combining Vectors

Scalars are quantities that have only a magnitude and no direction. Multiplying a vector by a scalar changes the vector's magnitude; this is called scalar multiplication. Changing the magnitude of a vector does not indicate a change in its direction.

Definition of Scalar Multiplication


The product $k\vec{v}$ of a vector \vec{v} and scalar k is a vector with a magnitude that is $|k| \cdot ||\vec{v}||$ and a direction that is equal to the direction of \vec{v} if $k > 0$ and opposite the direction of \vec{v} if $k < 0$. If $k = 0$ or $\vec{v} = \vec{0} \implies k\vec{v} = \vec{0}$

Because each vector may have its own direction, the process for adding vectors is different from adding scalars. The most common graphical method for adding two vectors is to place the initial point of the second vector at the terminal point of the first. The sum of the vectors \vec{v} and \vec{w} is the vector with an initial point that coincides with the initial point of \vec{v} and a terminal point that coincides with the terminal point of \vec{w} .



For $\vec{u} = \vec{v} + \vec{w}$, the initial point of \vec{u} is the initial point of \vec{v} and the terminal point is the terminal point of \vec{w} . These three vectors form a triangle, it follows that the length of any one side is less than the sum of the lengths of the remaining sides, therefore: $||\vec{u}|| \leq ||\vec{v}|| + ||\vec{w}||$

For vector subtraction, $\vec{v} - \vec{w}$ is defined as $\vec{v} + (-\vec{w}) = \vec{v} + (-1)\vec{w}$. The vector $\vec{v} - \vec{w}$ is called the vector difference and is depicted graphically by drawing a vector from the terminal point of \vec{w} to the terminal point of \vec{v} .



Vector Components

A vector with initial point at the origin is called a standard-position vector. Because the initial point of any vector in standard position is (0,0), it can be described by the coordinates of its terminal point.

Definition

The vector with initial point (0,0) and terminal point (x,y) can be written in component form as

$$\vec{v} = \langle x, y \rangle$$

The scalars x and y are called the components of \vec{v}

For a vector not already in standard form its component form can be determined algebraically by subtracting the x and y values of the initial point from those of the terminal point.

Let \vec{v} be a vector with initial point (x_i, y_i) and terminal point (x_t, y_t) . Then \vec{v} can be expressed in component form as


$$\vec{v} = \langle x_t - x_i, y_t - y_i \rangle$$

To find the magnitude of a vector, calculate the distance between its initial and terminal points. The magnitude of vector $\vec{v} = \langle x, y \rangle$ is denoted $||\vec{v}||$ and can be calculated using the formula

$$||\vec{v}|| = \sqrt{x^2 + y^2}$$

Based on this, it is clear that for any vector \vec{v} , $||\vec{v}|| \geq 0$, and $||\vec{v}|| = 0$ if and only if $\vec{v} = \vec{0}$.

The magnitude of a vector can also be derived using Pythagorean theorem.



Expressing vectors in component form allows scalar multiplication and vector addition to be performed algebraically.

Let $\vec{v} = \langle x_1, y_1 \rangle$ and $\vec{w} = \langle x_2, y_2 \rangle$ be vectors, and let k be a scalar.

Scalar Multiplication: $k\vec{v} = \langle kx_1, ky_1 \rangle$

Vector Addition: $\vec{v} + \vec{w} = \langle x_1, y_1 \rangle + \langle x_2, y_2 \rangle = \langle x_1 + x_2, y_1 + y_2 \rangle$

Properties of Vector Operations

Let \vec{u}, \vec{v} , and \vec{w} be vectors in a plane. Let r and s be scalars.

i.	$\vec{u} + \vec{v} = \vec{v} + \vec{u}$	Commutative Property
ii.	$(\vec{u} + \vec{v}) + \vec{w} = \vec{u} + (\vec{v} + \vec{w})$	Associative Property
iii.	$\vec{u} + \vec{0} = \vec{u}$	Additive Identity Property
iv.	$\vec{u} + (-\vec{u}) = \vec{0}$	Additive Inverse Property
v.	$r(s\vec{u}) = (rs)\vec{u}$	Associativity of Scalar Multiplication
vi.	$(r + s)\vec{u} = r\vec{u} + s\vec{u}$	Distributive Property
vii.	$r(\vec{u} + \vec{v}) = r\vec{u} + r\vec{v}$	Distributive Property
viii.	$1\vec{u} = \vec{u}$	Identity Property
ix.	$0\vec{u} = \vec{0}$	Zero Property

Proof of Commutative & Distributive Properties

Let $\vec{u} = \langle x_1, y_1 \rangle$ and $\vec{v} = \langle x_2, y_2 \rangle$

Commutative Property

Apply the commutative property for \mathbb{R}

$$\vec{u} + \vec{v} = \langle x_1 + x_2, y_1 + y_2 \rangle = \langle x_2 + x_1, y_2 + y_1 \rangle = \vec{v} + \vec{u}$$

Distributive Property

Apply the distributive property for \mathbb{R}

$$\begin{aligned}
 r(\vec{u} + \vec{v}) &= r \cdot \langle x_1 + x_2, y_1 + y_2 \rangle \\
 &= \langle r(x_1 + x_2), r(y_1 + y_2) \rangle \\
 &= \langle rx_1 + rx_2, ry_1 + ry_2 \rangle \\
 &= \langle rx_1, ry_1 \rangle + \langle rx_2, ry_2 \rangle \\
 &= r\vec{u} + r\vec{v}
 \end{aligned}$$