

(a)
RMS over sites of $\bar{\phi}_i - \mathcal{M}_\star(\bar{h}_i)$
(b)
Off-diagonal RMS of $C^{\check{\phi}}_{ij}(\tau{=}0) - \bar{C}^{\check{\phi}}_{ij}(\tau{=}0)$
measured
single-site
TAP-solved
(c)
Relative RMS error
(d)
Off-diagonal RMS of $C^{\check{\Delta}}_{ij}(\tau{=}0)$, floor subtracted
Sigmoid
Sigmoid, biases
Hebbian, $k = -2$
Hebbian, $k = +2$
Lotka–Volterra
$N^{-1/2}$
N^{-1}
sampling floor
$C^{\check{\phi}}_{ij}(\tau{=}0)$
-0.51 ± 0.01
-0.97 ± 0.02
-1.00 ± 0.06
-1.02 ± 0.04
-0.46 ± 0.01
-0.50 ± 0.05
-0.50 ± 0.03
-1.07 ± 0.02
-0.49 ± 0.01
-0.93 ± 0.03
-0.90 ± 0.04
-0.92 ± 0.04
-0.42 ± 0.03
-0.43 ± 0.03
-0.45 ± 0.04
-1.05 ± 0.03
-0.51 ± 0.01
-0.96 ± 0.02
-0.94 ± 0.02
-0.94 ± 0.03
-0.47 ± 0.01
-0.44 ± 0.02
-0.47 ± 0.02
-1.03 ± 0.02
-0.52 ± 0.01
-0.91 ± 0.02
-0.97 ± 0.02
-0.98 ± 0.03
-0.44 ± 0.02
-0.47 ± 0.02
-0.48 ± 0.03
-1.06 ± 0.03
-0.52 ± 0.02
-0.94 ± 0.05
-0.90 ± 0.05
-0.90 ± 0.04
-0.51 ± 0.03
-0.44 ± 0.02
-0.43 ± 0.02
-1.11 ± 0.05
N