Remark 3.15. By [19, Remark 2.28], if a definable function between two sets in Γ is a vrv-contraction then it is piecewise $\mathbb{Z}$-affine (with constant terms of the form $\mathrm{vrv}(t)$, where $t \in \mathrm{RV}$ is definable, that is, $t \in \mathrm{RV}(\mathbb{S})$). Moreover, a definable bijection between two objects of $\Gamma[k]$ is a $\Gamma[k]$-morphism if and only if it is definably a piecewise $\mathrm{GL}_k(\mathbb{Z})$-transformation. The "if" direction is clear. For the "only if" direction, see [8, Lemma 10.1] or [19, Lemma 2.29].

Referee R050 [Clarification] – Piecewise integral-affine behavior.
Original source line 1289. The first sentence needs "piecewise $\mathbb{Z}$-affine," not globally "$\mathbb{Z}$-linear." For example, $\gamma \mapsto |\gamma|$ is the vrv-contraction of the definable map $t \mapsto t$ for $\mathrm{vrv}(t) \geq 0$, $t \mapsto t^{-1}$ otherwise. Its two integral-linear formulas differ. Retain the specified restriction on the translation constants.

Definition 3.16 (Γ-categories with volume forms)**.** An object of the category $\mu\Gamma[k]$ is a definable pair (I, ω), where $I \in \Gamma[k]$ and $\omega : I \longrightarrow \Gamma$ is a function. A $\mu\Gamma[k]$-morphism between two objects (I, ω), (J, σ) is a $\Gamma[k]$-morphism $F : I \longrightarrow J$ such that for all $\alpha \in I$,

$$\omega(\alpha) = \sigma(F(\alpha)) + \mathrm{Jcb}_\Gamma F(\alpha).$$

The category $\mu\Gamma^{\mathrm{fin}}[k]$ is the obvious full subcategory of $\mu\Gamma[k]$.

Notation 3.17. For each $(U, f, \omega) \in \mu\mathrm{RV}[k]$, write $\omega_f : U \longrightarrow \Gamma$ for the function given by $u \longmapsto \Sigma(\mathrm{vrv} \circ f)(u) + \omega(u)$. Similarly, for $(I, \sigma) \in \mu\Gamma[k]$, write $\sigma_I : I \longrightarrow \Gamma$ for the function given by $\gamma \longmapsto \Sigma\gamma + \sigma(\gamma)$.

There is a natural map $\Gamma[*] \longrightarrow \mathrm{RV}[*]$ given by $I \longmapsto (I^\sharp, \mathrm{id})$ (see Notation 2.7). This map induces a commutative diagram in the category of graded semirings:

$$\begin{array}{ccc} \mathbf{K}_+\,\Gamma^{\mathrm{fin}}[*] & \rightarrowtail & \mathbf{K}_+\,\mathrm{RES}[*] \\ \downarrow & & \downarrow \\ \mathbf{K}_+\,\Gamma[*] & \rightarrowtail & \mathbf{K}_+\,\mathrm{RV}[*] \end{array}$$

where all the arrows are monomorphisms. The map

$$\mathbf{K}_+\,\mathrm{RES}[*] \times \mathbf{K}_+\,\Gamma[*] \longrightarrow \mathbf{K}_+\,\mathrm{RV}[*]$$

determined by the assignment

$$([(U, f)], [I]) \longmapsto [(U \times I^\sharp, f \times \mathrm{id})] \tag{3.3}$$

is well-defined and is clearly $\mathbf{K}_+\,\Gamma^{\mathrm{fin}}[*]$-bilinear. Hence it induces a $\mathbf{K}_+\,\Gamma^{\mathrm{fin}}[*]$-linear map

$$\Psi : \mathbf{K}_+\,\mathrm{RES}[*] \otimes_{\mathbf{K}_+\,\Gamma^{\mathrm{fin}}[*]} \mathbf{K}_+\,\Gamma[*] \longrightarrow \mathbf{K}_+\,\mathrm{RV}[*], \tag{3.4}$$

which is a homomorphism of graded semirings. By the universal mapping property, groupifying a tensor product in the category of $\mathbf{K}_+\,\Gamma^{\mathrm{fin}}[*]$-semimodules is, up to isomorphism, the same as taking the corresponding tensor product in the category of $\mathbf{K}\,\Gamma^{\mathrm{fin}}[*]$-modules; the groupification of Ψ is still denoted by Ψ. Similarly, there is a $\mathbf{K}_+\,\mu\Gamma^{\mathrm{fin}}[*]$-linear map

$$\mu\Psi : \mathbf{K}_+\,\mu\mathrm{RES}[*] \otimes_{\mathbf{K}_+\,\mu\Gamma^{\mathrm{fin}}[*]} \mathbf{K}_+\,\mu\Gamma[*] \longrightarrow \mathbf{K}_+\,\mu\mathrm{RV}[*]. \tag{3.5}$$

We shall abbreviate $\otimes_{\mathbf{K}_+\,\Gamma^{\mathrm{fin}}[*]}$, $\otimes_{\mathbf{K}_+\,\mu\Gamma^{\mathrm{fin}}[*]}$ both as "$\otimes$" below when no confusion can arise.

Proposition 3.18 ([8, Corollary 10.3, Proposition 10.10(1)])**.** *Ψ and $\mu\Psi$ are both isomorphisms of graded semirings.*

Note that, however, [8, Proposition 10.10(2)] does not hold. One of the main motivations of this paper is to construct an alternative to be used in applications.