Lemma 2.28 ([18, Lemma 3.13])**.** *Let $A \subseteq \mathrm{VF}$ be an atomic open disc and $f : A \longrightarrow \mathrm{VF}$ a definable function. Then either f is constant or $f(A)$ is also an atomic open disc.*

Lemma 2.29. *Let $\mathfrak{a}$, f be as in Lemma 2.26 and suppose that f is not constant. Then there are $t_1, \ldots, t_k \in \mathrm{RV}_\infty$ and for each i, a t_i-definable function $f_i : \mathfrak{a} \longrightarrow t_i^\sharp$ such that $f = \bigcup_i f_i$ and each $f_i(\mathfrak{a})$ is either a point or a t_i-atomic open disc.*

Proof. By Lemma 2.26 and injectivity of rv on each finite fiber, f decomposes into finitely many t-definable single-valued branches $g : \mathfrak{a} \longrightarrow t^\sharp$. A branch with $t = \infty$ is the constant zero branch. Any constant branch already has the required form, so fix a nonconstant branch, for which $t \in \mathrm{RV}$. Lemma 2.24 makes $\mathfrak{a}$ atomic over t.

Every coordinate line of $\mathfrak{a}$ is atomic over its fixed coordinates and t. By Lemma 2.28, its image under g is a point or an open disc. Any two points of $\mathfrak{a}$ can be joined by changing one coordinate at a time. Starting at a point with image p and ending at a point with different image, the first nonconstant line image is an open neighborhood of p contained in $g(\mathfrak{a})$. Thus $g(\mathfrak{a})$ is open. It is a proper definable subset of VF, since it lies in $t^\sharp$.

By C-minimality, every point of a proper definable open subset of VF lies in a maximal open disc contained in that subset. Choose such a disc $D \subseteq g(\mathfrak{a})$ containing one value of g. An open line image that meets D is contained in D: intersecting open discs are nested, and maximality excludes a larger one inside $g(\mathfrak{a})$. A point image also stays in D when its initial value is in D. The coordinate chains therefore keep every value of g in D, so $g(\mathfrak{a}) = D$. The definable image of an atomic set is atomic over the same base, hence this open disc is t-atomic, as required.

□

Resolved referee R030 [Gap; revised] – A unary image lemma is used on a polydisc.
Original source line 918. For $n > 1$, Lemma 2.27 does not directly apply to $f : \mathfrak{a} \longrightarrow \mathrm{VF}$, since it has a unary domain. Supply the missing argument that $f(\mathfrak{a})$ has just one primitive component. A possible route is the canonical finite component decomposition of an atomic unary set: every coordinate slice of $\mathfrak{a}$ is atomic over its fixed coordinates, and its image is a point or an open disc by induction. Show that such an image must lie in one primitive component, then connect any two source points by changing coordinates one at a time. The same coordinate-chain argument excludes a closed disc or thin annulus as the whole image, because each slice image lies in a single maximal open subdisc. This fills the role of the inapplicable unary citation, once the component claim is spelled out.

Author resolution. The author approves the coordinate-chain proof. Unary coordinate-line images are points or open discs; maximal open discs in a proper definable open image then force the entire nonconstant image to be one open disc. This replaces the inapplicable unary citation on the polydisc. The tag ranges here and in Corollary 2.30 are also enlarged to RV_∞, retaining the constant zero branch.

Unapplied R031 proposal for author review. The full graph of a branch on $\mathfrak{a}$ names its domain by $\ulcorner \mathfrak{a} \urcorner$: even $f(x) = \{1\}$ on $A = \mathrm{VF}$ restricts to arbitrary polydiscs, while its tag $\mathrm{rv}(1)$ stays fixed. The green edits explicitly allow $(\ulcorner \mathfrak{a} \urcorner, t)$ parameters, cite the corrected multidimensional Lemma 2.29, and state the uniformly coded property used in compactness. These edits await approval.

Corollary 2.30. *Let $A \subseteq \mathrm{VF}^n$ be a definable set and $f : A \longrightarrow \mathcal{P}(\mathrm{VF})$ a definable finitary function as in Lemma 2.26. Then there are a definable finite partition $(A_i)_i$ of A and, for each i and all open polydiscs $\mathfrak{a} \subseteq A_i$, finitely many $(\ulcorner \mathfrak{a} \urcorner, t_{\mathfrak{a}j})$-definable functions $f_{\mathfrak{a}j} : \mathfrak{a} \longrightarrow t_{\mathfrak{a}j}^\sharp$, where $t_{\mathfrak{a}j} \in \mathrm{RV}_\infty$, such that $f \restriction \mathfrak{a} = \bigcup_j f_{\mathfrak{a}j}$ and each $f_{\mathfrak{a}j}(\mathfrak{a})$ is either a point or an open disc.*

Proof. By compactness, there is a uniform finite bound N on the cardinalities of the fibers $f(a)$. For an open-polydisc code d, write $\mathfrak{a}_d$ for its polydisc and let $P(d)$ say that $\mathrm{rv} \circ (f \restriction \mathfrak{a}_d)$ is constant

and that the image of each of its single-valued rv-tagged branches is a point or an open disc. This is a first-order condition: use the finite disjunction over the possible numbers $0 \leq k \leq N$ of distinct tags, and express a point or an open disc by its usual uniform codes. Injectivity of rv on each fiber gives uniqueness of each tagged branch.

For $a \in A$, let $D_a \subseteq A$ be the complete type-definable set containing a. By Lemma 2.25, every open polydisc $\mathfrak{a}_d \subseteq D_a$ is $\ulcorner \mathfrak{a} \urcorner_d$-atomic. Lemma 2.29, over the ground base together with this disc parameter, therefore gives $P(d)$. If no finite part of the type of a had this property for every polydisc contained in it, compactness would produce an open-polydisc code d with $\mathfrak{a}_d \subseteq D_a$ and $\neg P(d)$, a contradiction. Thus there is a ground-definable set $A_a \subseteq A$ containing a such that $P(d)$ holds whenever $\mathfrak{a}_d \subseteq A_a$. A second application of compactness gives finitely many such sets covering A; take the partition they generate.

For a polydisc $\mathfrak{a}$ in a partition piece, choose its finitely many distinct tags $t_{\mathfrak{a}j} \in \mathrm{RV}_\infty$. The full graph of its jth branch is

$$\{(x, y) : x \in \mathfrak{a},\ y \in f(x),\ \mathrm{rv}(y) = t_{\mathfrak{a}j}\}.$$

It is $(\ulcorner \mathfrak{a} \urcorner, t_{\mathfrak{a}j})$-definable, the branches reconstruct $f \upharpoonright \mathfrak{a}$, and $P(d)$ gives the asserted images. □

Referee R031 [Error] – The local branch functions need the disc parameter.
Original source line 925. The functions $f_{\mathfrak{a}j}$ must be allowed parameters $(\ulcorner \mathfrak{a} \urcorner, t_{\mathfrak{a}j})$, not just $t_{\mathfrak{a}j}$. Their domains are $\mathfrak{a}$, so literal $t_{\mathfrak{a}j}$-definability would imply that every such disc is $t_{\mathfrak{a}j}$-definable. Already for $f(a) = \{1\}$ on $A = \mathrm{VF}$, this is impossible for arbitrary subdiscs, since $t_{\mathfrak{a}j} = \mathrm{rv}(1)$ is fixed. The proof itself introduces $\ulcorner \mathfrak{a} \urcorner$, which explains the missing parameter. Also replace the appeal to the unary Lemma 2.28 by the corrected multidimensional Lemma 2.29, and explicitly formulate the uniform first-order property used in the compactness step.

Definition 2.31. Let $f : A \longrightarrow B$ be a bijection between two sets A and B, each with exactly one VF-coordinate. We say that f is *concentric* if, for every VF-fiber f_t of f and all open polydiscs $\mathfrak{a} \subseteq \mathrm{dom}(f_t)$, $f_t(\mathfrak{a})$ is also an open polydisc; if both f and f^{-1} are concentric then f has the *disc-to-disc property*.

Lemma 2.32 ([18, Proposition 3.19])**.** *Let $f : A \longrightarrow B$ be a definable bijection between two sets A and B, each with exactly one* VF*-coordinate. Then there is a definable finite partition $(A_i)_i$ of A such that every restriction $f \upharpoonright A_i$ has the disc-to-disc property.*

2.3. **Continuity and fiberwise properties.**

Definition 2.33. We set $\dim_{\mathrm{RV}}(\varnothing) = -\infty$. For a nonempty definable subset $U \subseteq \mathrm{RV}_\infty^m$, the RV*-dimension*, denoted by $\dim_{\mathrm{RV}}(U)$, is the smallest number k such that there is a definable finite-to-one function $f : U \longrightarrow \mathrm{RV}^k$ (RV^0 is taken to be the singleton $\{\infty\}$).

Resolved referee R164 [Clarification; new] – Dimension of the empty set.
Original source line 941. Specify $\dim_{\mathrm{VF}}(\varnothing) = \dim_{\mathrm{RV}}(\varnothing) = -\infty$ (or another negative convention) separately. The VF definition otherwise has no largest natural number for the empty set, whereas the literal RV definition gives it dimension 0. With that latter convention the boundary lemma fails when $n = 0$, since its boundary is empty but $0 < 0$ is false. Alternatively restrict that lemma to $n \geq 1$, while still specifying the empty VF dimension.

Author resolution. The author approves $\dim_{\mathrm{VF}}(\varnothing) = \dim_{\mathrm{RV}}(\varnothing) = -\infty$. Both definitions now state this convention separately and retain their original clauses for nonempty sets, including zero-coordinate cases.